

LIST OF FIGURES

Fig. No.	Figure
1,2	Light microphotograph of control gills of bivalve, <i>Lamellidens marginalis</i>
3,4	Light microphotograph of gills of bivalve, <i>Lamellidens marginalis</i> exposed at 0.5 % of tannery wastewater for 10 days exposure period
5,6	Light microphotograph of gills of bivalve, <i>Lamellidens marginalis</i> exposed at 0.5 % of tannery wastewater for 20 days exposure period
7,8	Light microphotograph of gills of bivalve, <i>Lamellidens marginalis</i> exposed at 2 % of tannery wastewater for 10 days exposure period
9, 10	Light microphotograph of gills of bivalve, <i>Lamellidens marginalis</i> exposed at 2 % of tannery wastewater for 20 days exposure period
11, 12	Light microphotograph of control hepatopancreas of bivalve, <i>Lamellidens marginalis</i>
13, 14	Light microphotograph of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed at 0.5 % of tannery wastewater for 10 days exposure period
15,16	Light microphotograph of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed at 0.5 % of tannery wastewater for 20 days exposure period
17, 18	Light microphotograph of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed at 2% of tannery wastewater for 10 days exposure period
19, 20	Light microphotograph of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed at 2% of tannery wastewater for 20 days exposure period