
BIBLIOGRAPHY

B I B L I O G R A P H Y

- ✓ Abdelhadi, A.A., Y.M. Elkheir and T. Hassan (1989) A Succinyl Choline - like action of an Ipomoea carnea spp. fistulosa Mart ex Choisy extract. Pharmacol Res 24(4) 431-437.
- ✓ Austin Danial F. (1977), Ipomoea carnea Jacq Vs Ipomoea fistulosa Mart, ex choisy. TAXON 26(2/3) : 235-238.
- ✓ Arnon, D.I. (1949), Copper enzymes in isolated chloroplasts, polyphenol oxidase in Beta Vulgaris plant physiol. 24 : 1-15.
- ✓ Bonset, K. (1993), Control of environmental factors on pattern of montane rain forest in Puerto Rico. Tropical Ecology 34 (1) : 53-63.
- ✕ (Banin, A. and Navrot, J.) (1992), Pattern of iron distribution in the soil plant system and its possible relation to iron chlorosis Comm. in Soil Science and Plant analysis 3 (3) : 177-82.
- ✓ Baskin J.M. and Cardl C. Baskin (1974). Some ecophysiological aspects of seed dormancy in Geranium carolinianum L. from Central Tennessee. OECOLOGIA (BERL) 16(3): 209-219.
- ✓ Baskin, J.M. and Cardl C. Baskin (1975). Ecophysiology of seed germination and germination in Torilis japonica in relation to its life cycle strategy. Bull. Torrey Bot. Club. 102 (2) : 67-72.

- ✓ Baskin, J.M. and Cardl C. Baskin (1983). Germination ecophysiology of eastern deciduous forest herbs. Hydrophyllum macrophyllum A.M. DIDL NAT. 109(1) : 63-71.
- ✓ Baskin, J.M. and Cardl, C. Baskin (1986). Germination ecophysiology of the mesic deciduous forest herb Isopyrum biternatum. Bot.Gaz. 147 (2) : 152-155.
- ✓ Bear, F.E.; A.L. Prince, S.J. Toth and E.R. Purvis (1951). Magnesium in plants and soils, New Jersey Agr.Expt. St. Bull. 260 : 3-24.
- ✓ Bhanot, R.K. and G.P. Bhawane (1992). Impact of different plant extractives on the digestive carbohydrates of adult Leucopholis lepidopnora. JADU,ZOO.13(1-2): 19-24.
- ✓ Bhattacharyya, P.K. (1988). The Vascular Cambia of doder and its allies. Bull.Bot.Surv.INDIA. 30 (1-4): 149-155.
- Bhattacharyya, P.K. and A.R. Midya (1980). Arboreal Ipomoea of the Indian Botanical garden Calcutta, India. Bull. Bot.Soc. Bengal, 33 (½) : 75-86.
- Bhattacharyya, P.K. (1976). A note on two species of Ipomoea namely I. carnea Jacq and fistulosa Mart. Ex. choisy. in eastern Asia. JBombay Nat.Hist.Soc.73 (3): 317-320.
- ✓ Bhosale L.J. (1974). Physiology of salt tolerance of plants. Ph.D. Thesis, submitted to Shivaji University, Kolhapur (India).
- Blackman, E.F. (1905). Optima and limiting factors. Ann.Bot. 19 : 281-295.

✓ Braun - Blanquet, J. (1932), Plant sociology. (English translation by G.D. Fuller and H.S. Conard), McGraw Hill Book Co. Inc, New York, p. 438.

✓ Borchert, Rolf. (1980). Phenology and ecophysiology of tropical trees. Erytrina poeppigiana. Ecologia, 61(5) : 1065-1074.

Chirputkar, M.B. (1969). Physiological studies in marine plants of Bombay Ph.D. Thesis University of Bombay, India.

✓ Datta, S.C. and S. Sinha - Ray (1995) APTI value of five different species of higher plants from the metropolis of Calcutta. Environ and Adaptive Biol of Plants spp. : 289-294.

Duhan, J.S., K. Lakshinarayana (1995), Allelopathic effect of Acacia nilotica on cereal and legume crops grown in fields. Allelopathy Journal, 2 (1): 93-98.

✓ Dawalkar M.P. and Joshi G.V. (1962). Effect of Saline environment on ion uptake and distribution in Ipomoea pescarpae J. Biol. Sci. 5(1) : 89-93.

D. Leela (1995). Allelopathic effects of purple nut sedge Cyperus rotundus L. tubers on growth of field crops Allelopathy journal, 2 (1) : 89-92.

D'Souza T.J. and K.B. Mistry (1979), Uptake, distribution and metabolic fate of ^{59}Fe , $^{58}_{\leftarrow}\text{Co}$ ^{50}Mn and ^{65}Zn in plants and their mobility and availability to crops in typical black and laterite soils, "Isotopes and Radiation in research on soil plant relationships". International Atomic Energy Age. Vienna.

- ✓ Elias Pavol (1949). Some ecophysiological features in leaves of plants in an Oak-Lorn bean forest. Folia. GeoBot. Phyta Taxon·14 (1) : 29-42.
- ✓ Elias PaVol (1981). Some ecophysiological leaf characteristics of components of spring synusium in temperate deciduous Forests Biologia (Bratisl) 36 (10): 841-850.
- ✓ Epstein, E. (1972). Mineral nutriton of plants : principles and perspectives John Wilgy and Sons. Inc., New York.
- ✓ Ferry, J.F. and H.S. Ward (1959). Fundamental of Plant Physiology MacMillan and Co. New York.
- ✓ Ford, C.W. and J.K. Wilson (1981). Changes in levels of Solutes during osmotic adjustment to water stress in leaves of four tropical pasture species. Aust. J. plant physiol.·8 : 77-91.
- ✓ Gold C.S., J.A. Wightmar and M.P. Pimers (1991). Effect of malches on Frnging behaviour of Microtermes obesi and odontolermes sps. in India. Insect SCI. APPL 12 (1-3) : 297-304.
- Growth Doris (1991) Morphological characterization of seeds and seedling of seven weed species of convolvulaceae occuring in agricultural field in Brazil. Iheri. NGIA SER BOTO (41) : 83-100.
- ✓ Gupta, O.C.D., Rajni Gupta and P.C.Gupta (1980). Chemical examination of Flowers of Ipomoea fistulosa, Planta. Med. 38 (2) : 147-150.

- ✓ Hardcastle W.S. (1978). The influence of temperature and acid scarification duration on Ipomoea obscura Hersk Seed germination. WEED RES. 18 (2) : 89-92.
- ✓ Hawk, P.B.; B.L. Oser and W.H. Sumerson (1948) Practical physiological chemistry b. The Black stone Co. USA Toronto.
- Hsiao, T.C. (1973). Plant response to water stress. Ann.Kev. Plant physiol. 24 : 519-570.
- ✓ Huzulak Joref and Pavol Elias (1975) Within-crown pattern of ecophysiological features in leaves of Acer cumpestor and Carpinus betulus. FOLIA GEOBOT. PHYTOTAXON. 10 (4) : 337-350.
- Jha, K.P., Dinesh Chandra and B.B. Nanda (1980). Yield and nitrogen uptake in rice as influenced by green leaf manuring with Ipomoea carnea, ORYZA II (1) : 18-23.
- ✓ Joseph, D.E. and J.N. Dube (1988). Patterns of phosphate solubilization by Aspergillus niger. PROC. NATL. ACAD. SCI. INDIA SECT. B. (Biolsci.) 58(2): 329-33.
- ✓ Jotsova, Baurenska, Nadezhda (1986). Ecological and physiological studies of Junglans regia in the region of Dolni-Lom. Mikhaylovg district (Balgeria) Rasteniev'd NAVKT 23 (1) : 91-96.
- ✓ Kanaujia, R.S. (1977). Observation on soil fungistasis VI Fungistasis of amended soils. Bangladesh J.Bot. 5(½) : 1-7.

- ✓ Karatela, Y.Y. and L.S. Gill (1985). Epidermal morphology and stomatal ontogeny in some West African Convolvulaceae species. Herba. Hung. 24(2/3): 11-18.
- ✓ Khan, A.M., V. Pandey, M.Yunus and K.J. Ahmed (1989). Plants as dust Scavengers : A case study Indian For. 115 (9) : 670-672.
- ✓ Kirk J.O.T. and R.L. Allen (1965). Dependence of Chloroplast pigment synthesis on protein synthesis. Effect of actidione, Arch. Biochem. Biophys. Res. Commun. 21: 523 - 530.
- Kulkarni, P.K. (1990). Some ecological and physiological aspects of Rhizophora propagules. Ph.D. Thesis, Shivaji University, Kolhapur, India.
- ✓ Lardizabal, Recard D. and P.G. Thompson (1990). Growth regulators combined with grafting increase flower number and seed production in sweet potato. Hort.Science 25(1) : 79-81.
- ✓ Lawlor, D.W. and O.F.J. Milford (1973). The effect of sodium on growth of water stressed sugarbeat Ann.Bot. 37 (151) : 597-604.
- Leal Nea MaCedo (1974). Anatomical characteristic of Ipomoea fistulosa mart. BOL INST BIOLBAHIA 18 (1) : 107-126.
- Leeras Eduardo (1976). Ecophysiological parameters in leaves I. Basic considerations. Acta Amazonica 6(4) : 409-416.

- Liasen, J.S. and A. Jensen (1971). Quantitative determination of carotinoids in photosynthetic tissues. In: Methods in Enzymology (Ed) San Pietro A. Academic Press Inc. publishers, New York, pp. 586-602.
- Mall, L.P., V.P. Singh; A. Garge and S.M. Pathak (1986). Mangrove forest of Andamans and some aspects of its ecology in The Mangroves : Proc.Natsymp. Biol. Util. Cons. mangroves (ed L.J. Bhosale) Nov. 1985: 49-59.
- Mallik Firoj and R.P. Parkayastip (1975). Notes on some Hyphomycetes of eastern India. Bull.Bot.Soc.Bengal 29 (1) : 53-55.
- Masarvicova Elenu and M. Duda (1976). Some ecophysiological aspects of quantitative changes of carotenoids in the leaves of Palmonaria officinalis L. spp. maculosa Biologia (Bratisl) 31(1) : 15-23.
- Mc donald, J. Andrew (1992). Evolutionary implications of typical and anomalous secondary growth in arborescent Ipomoea, Bull Torey Bot club 119 (3) : 262-267.
- M. Eyini, A.V. Mateswari, T. Chandra and M. Jayakumar (1996). Allelopathic effects of Leguminous plant leaf extracts on some weeds and corn. Allelopathy Journal 3(1) : 85-88.
- Nair G.G., M. Daniel and S.D. Sabnis (1986). Chemosystematics of Ipomoea Linn and some related taxa. Curr. Scy. (Bangalore) 55 (19) : 961-965.

- Narayanan, N. and P. Sivadas (1966). Studies on the intertidal macrofauna of the sandy beach at Kavaratii Atoll (Lakshadweep). Mahasagar Bull of NIO 19 (1) : 11-12.
- Nielsen K.F.; T. Caddy and W. Wood (1960). The influence of the extract of some crops and soil residues on germination and growth Can. J. Plant. Sci. 40 : 188-197.
- Osmond, C.B. (1966). Divalent cations absorption and Interaction in Aripex, Aust. J. Biol. Sci. 19 : 37-48.
- Pande, H.K. and P. Singh (1969). Effect of moisture and nitrogen on growth, yield and mineral content of rice. Exp. Agri. 5 (2) : 125-132.
- Pandey, N.D.; H. Shivrajsing and G.C. Tewari (1977). Use of some plant powders, oils and extracts as protectance against pulse beetle Callosobracho Chinensis Linn. Indian J. Entomol 38 (2) : 110-113.
- Pandey, N.D.; Lal Singh; Y.P. Singh and R.A. Tripathi (1987). Effect of certain plant extracts against Lipaphis carysi Kalt under laboratory condition. Indian J. Entomol. 49 (2) : 238-242.
- Patil, B.B. and R. De. (1976). Influence of antitranspirants on rape seed (Brassica campestris) plants under water shessed and not stressed conditions. Plant physiol. 59 : 941-943.
- Paul, Kukali and A. Mandal (1987). Effect of phenoxy acids on growth and development of Costus speciosus. Cobios. 14 (2-3) : 134-136.

Pavlovic S.; R. Jancis; A. Galin, E. Roznjestove and A.S. Elena (1980). Anatomical structure ecophysiological properties and active substances of the plant species. Peucedanum oreselinum. Acta. Biol. Med. Exp. 5 (1): 37-42.

✓ Prabha Kiran and S.C. Gupta (1984). Effects of floral age and flowering season on self-incompatibility vigor in Ipomoea fistulosa. BEITR. BIOL. PFLANZ 59(3): 359-366.

Perez Amadar M.C.; A. Gontale; J. Morquez; J. Bailin; F. Garciajimentz and O. Collera (1980). Chromatographic profiles of the seeds of some spp. of convolvulaceae Phyton. Rev. INT.BOT.Exp. 39 (1): 85-94.

✓ Price, C.A. (1988). Iron Compounds and plant nutrition. Ann. Rev. Plant physiol 19: 239-248.

✓ Putnam, A.R. and W.B. Duke (1974). Biological suppression of weeds evidences for all allelopathy in accessions of cucumber, Science 185: 310-372.

✓ Rahman, A.A.A.; A.F. Shalaby and M.O.E. Monayeri (1971). Effect of moisture stress on metabolic products and ions accumulation. Plant and Soil 34: 65-90.

✓ Rajan Bawa (1992). Effect of Eucalyptus globulus and Aesculus indica leaf leachates on growth of Glaucium flavum and Digitalis lanata seedlings. Indian J.Ecol. 19(1): 105-107.

Rao S.R.; Shanukha and M. Leela (1990). Leaf architecture in relation to taxonomy Ipomoea L. Feddes Report 101 (11/12) : 611-616.

- ✓ Rao, M.V. and P.S. Dubey (1992). Occurrence of heavy metals in air and their accumulation by tropical plants growing around in industrial area Sci.Total Environ. 126 (1-2) : 1-16.
- ✓ Rao, T.A. and K.R. Agrawal and A.R. Mukharjee (1966). Ecological studies of Saurashtra Coast and neighbouring island - IV Piram Island Bull.Bot.Surv. India 8: 60-67.
- ✓ Rai, V.N. and S.B. Agrawal (1985). Seed polymorphism and germination behaviour of Ipomoea muricata growing in cultivated Indian tropicdeserts. Acta. Bot.Indica 13 (1) : 1-5.
- ✓ Raza, S.H., K. Vikaya and R.M.S. Murthy (1985). Air pollution tolerance index of certain plants of Hyderabad Symp Biomonitoring State Environ. p. 243.
- ✓ Sadashivam S. and T. Balasabraminan (1987). Practical Manual Biochemistry Tamil Nadu Agricultural University, Coimbatore P. 14.
- ✓ Saxena Anurag and A.K. Sharma (1996). Allelopathic potential of Acacia tortilis in agroforestry systems of arid regions. Allelopathy Journal 31 (1): 81-84.
- ✓ Sekine T. (1965). Photoelectric Calorimetry Biochemistry (2nd Ed.) Nanchod Publishing Company, Tokyo pp. 242.
- ✓ Shahi, A.K., Singh and S. Chandra (1990). Growth analysis studies based on degree days and its significance to howest management practices on Cymbopogon jwarancasa Schult Curr. Agric. 14 : 13-16.

- ✓Shanmughavel, P. (1995). Seasonal primary production and energy accumulation in macrophytic vegetation of Banar lake Sathyamangalam forests. Ad. plant. Sci. 8(2) : 410-413.
- ✓Shamsi S.R.A. and F.H. Whitehead (1974), Comparative ecophysiology of Epilobium hirsutum L. and Lythrum Salicaria L. J. Ecol. 62 (2) : 631-645.
- Sharma B.D.; D. Rathore and R. Harsh. (1995). Phytochemistry in relation to ecology of Rajasthan pteridophytes. Environ. and Adaptive Biol. of plants pp. 37-40. (ed.) D.D. Chawan Scientific publishers. Jodhpur (India).
- ✓Sharma, S.K.; J.S. Saini; I.M. Mishra and M.P. Sharma (1989). Classification of woody biomass : Ipomoea fistulosa plant stem. Bio Wastes 28 (1) : 25-32.
- ✓Sharma, S.S. and D.N. Sen. (1978). Water relations in three species of the convolvulaceae in the Indian arid zone. Folia Geo Bot Phytotaxon 13 (2) : 165-173.
- Shelford V.E. (1952). Paired factors and master factors in environment relations. F Illinois Acad. Sci. Trans. 45 : 155-160.
- Shingh V., A.M. Wadhvani and B.M. Johri (1983) Dictionary of economic plants in India ed Rajendra Sing pp. 108.
- ✓Shingh, S.K. and D.N. Rao (1983). Evaluation of plants for their tolerance to air pollution. Proc. symp. Air Pollution control. Delhi p. 218.

- ✓ Shingh, A.; M. Madan and P. Vasudevan (1987). Increasing biomass yields of horby weeds through coppicing : Studies on Ipomoea fistulosa and Adhatoda Vasica with reference to wastland utilization. Biol.Westes 19 (1): 25-34.
- ✓ Shingh, S.P.; U.R. Pal and K. Luka. (1989). Allelopathic effect of three serious weeds of Nigerian Savana on germination, seedling vigor of Soyabean and Maize. Journal of Agronomy and Crop. Science 162 (4) : 236-240.
- ✓ Sinha, A.K.; D.K. Singh; A.A. Baruah and V.P. Sharma (1992). Seasonal variation of physicochemical properties of bottom sediment of Karwar like Begusa. J. Freshwater Bio-4 (4) : 249-254.
- Singh, A.; M. Madan and P. Vasudevan. (1988). Fertilizer replacement trials with Raphanus Sativa L. Trop. Agric. 65 (4) : 361-363.
- ✓ Stout, P.R. (1961). Micronutrients in Crop. Vigour. Proc. 9th Ann. Ealif fertilizer conf. pp. 21-23.
- ✓ Sturgess, P. and D. Atkinson (1993). The clear feeling of sand-dunes plantations : Soil and Vegetational processes in habitat restoration. Biological conservation 16 : 171-183.
- ✓ Subramanian A.N. and V.K. Venugopalan. (1983). Phasporus and iron distribution in two mangrove species in relation to environment. Mahasagar Bulletin of the National Institute of Oceanography, 16 (1) : 183-191.

- ✓ Suseelama and R.R. Venkata Raju (1994). Effect of Digera muricata L. extract on the germination and seedling growth of ground nut Allelopathy Journal 1 (1) : 53-57.
- Taptour, G.S.; E.I. Adam; H.M. Obeid and O.F. Idris (1974). Development of anaemia in goats fed with Ipomoea cornea BR. VET J. 130 (3) : 271-279.
- ✓ Thakur, P.S. and A. Thakur. (1990). Potential of extracted root forming factor from Ipomoea fistulosa on dormant populus stem cuttings. Exp. Biol 34 (4) : 385-386.
- Tirkey, K.; R.P. Yadava; T.K. Mandal and N.C. Banerjee (1988). Pharmacological study of Ipomoea carnea. INDIAN VET. J. 65 (3) : 206-210.
- ✓ Toth, S.J.; A.L. Prince; A. Wallace and S.D. Mikkelsen (1948). Rapid qualitative determination of eight mineral elements in plant tissues by systematic procedure involving use of flame photometer. Soil. Sci. 66 : 459-466.
- ✓ Turner, N.C. and P.B. Waggoner. (1968). Effect of changing stomatal width in a red pine forest on soil water content, leaf water potential, hole diameter and growth. Plant Physiol. 43 : 973-978.
- ✓ Ugborogho, R.E.; K.O. Ogunwenmo and O.T. Aina. (1992). Epidermal morphology of six taxa of Ipomoea in Nigeria - Feddes Report. 103 (7-8) : 543-550.
- ✓ Umar, S., J. Peter and M. Wichtl (1980). Isolation and Identification of agroclavine and dihydrolyseral from leaves of I.fistulosa Mart. plant Med. 40 (4) : 328-332.

- ✓ Van Ooststroom, S.J. (1953), Convolvulaceae. Flora Malesiuna
1 (4) : 388-512.
- Venture, J.C. (1946), Ecuador hig V. Med. trop. (Guaygui
Ecuador) 3 : 149-281.
- ✓ Wagner, H. (1932). The growth Course of different grains
especially of Oats. Z. pflanzernahr. Dung u. Bodenk
25, : 48-102.
- Weber J.M. and T.S. Ma (1976). Micro chemical investigations
of medicinal plants : XIII separation of the alkaloids
in the leaves of Ipomoea violecca using thin layer
chromatography and solid probe mass spectromety.
Microchim Acta 1 (2/3) : 227-242.
- Witters, W.L. (1975). Extraction and identification of clavine
and lysergic acid alkaloids from morning glories.
OHTO J Sci. 75 (4) : 198-201.
- Wolf, F. I. (1956). Changes in chlorophylls 'a' and 'b' in
autumn leaves. Am. J. Bot. 43 : 714-718.