

LITERATURE CITED

- Abdul Kareem, A. 1980. Evaluation of neem products on the control of gram pod borer. *Pesticides* 15 : 78
- Ajri D.S. Mali., A.R., Shelke S.S. Patil C.S. and Subhedar, A.J. 1986. A status paper on 'Problem of Whitefly, *Bemisia tabaci* (Gennadius) in cotton and other crops in western Maharashtra. Seminar on status of whitefly on cotton, Pune March 14, 1986.
- Amtonin, P. Baillod, M. Linder C. and Mittaz, C. 1997. Chemical and biological control of *Tetranychus urticae* in strawberry. *Revue Suisse – de – Viticulture, Arboriculture – et – d. Hort.*, 29(3) : 179 – 187.
- Anderson, P. 2001. History, current status, and collaborative research projects for *Bemisia tabaci*. *Crop. Prot.*, 20 : 709 – 723.
- Anonymous 1980 Evalution of different pesticides for the control of Whiteflies on cotton. Annual Report of Research work in Agriculturd Entomology. M.A.D., Parbhani, Maharashtra state, India, 25.
- Anonymous, 1990-91. Annual Report. Central Biological Control Station, Solan (Himachal Pradesh). pp. 28.
- Askaru,A .and zare N. 1976. Comparative study of the efficacy of insecticides and acaricides against two spotted spider mite on bean population. *Ind. J. Agri. Sci.* 46(6) : 257 – 259.
- Atwal, A .S; Choudhary, J.P.and sohi, B.S. 1969. Chemical control of red vegetables mite, *Tetranychus cucurbitae*. *Punjab Agri, University Res, J.* 214 – 219.
- Ayyangar, G.S.C. and P.G. Rao.1989. Neem (*Azadirachta indica* A.Juss) extract as larval, repellent and ovipositional deterrents to *Spodoptera litura* (Fab) *Indian J. Entomol.* 51 (2) : 121 –124.
- Bao – Li QIU; Shun – Ziang REN; N.S. Mandour and Shou – Young WEN. 2006. Population differenetation of *Bemisia tabaci* (Gennadius)

- (Hemiptera: Aleyrodidae) by DNA polymorphism in China. Journal of Entomological Research, 30 (1): 1 – 6.
- Bapu, P.C.S; Lakshmanan, P.Hanifa, A.N. and Gunasekaran, C.R. 1970. insecticidal control trials against mite, *Tetranychus telarius* on brinjal. South Ind.K.Hort 18 (1 & 2): 33 – 28.
- Barakat A.A., Shereef, G.M., Abdallah, S.S. Amer. S.S.A. (1984 / 85 Pub. 1986). Toxic action of some plant extracts against the *Tetranychus urticae* Koch. Bulettien of the Entomological society of Egypt, Economic series, 14 : 233-242.
- Basu, A.C. and Pramanik, L.M. 1965: Acaricidal tests of nine pesticides against *Tetranychus urtice* a pest of brinjal in west Benhal.J. Econ. Entoma, 61 (3) 768-770.
- Bedford, I.D., Briddon, R.W., Brown, J.K., Rosell, R.C. & Markham, P.G. (1994), Geminivirus transmission and biological characterization of *Bemisia tabaci* (Gennadius) biotypes from different geographic regions. *Annals of Applied Biology* 125: 311-325.
- Bhagat, K.C. and singh, W. 1999. Some observation on biology and behaviour of carmine spider mite. *Tetranychus annabarinus* – a pest of brinjal vegetable J. Advan., 20 (1) : 28 –31 .
- Bhokare, A.D.1986. Evaluation of some insecticides for the control of whitefly. *Bemisia tabaci* Geen. On Bringal *Solanum melongena* L. Ph. D Thesis submitted to Mahatma Phule Krishi vidyapeeth. Rahuri (M.S.)
- Bindra O.S. and Goyal N.P. 1966. Prdiminary studies in the control of vegetable mite on brinjal L.Res. Punjab agri University 3 : 174 – 177.
- Bindra, O.S. and Goyal, N.P. 1965. Preliminary studies in the control of vegetable mite on brinjal. J. Res, Punjab Agri. University. 3 : 174 – 177.

- Brown, J. K 2000. Genetic analysis of *Bemisia tabaci* Genn. (Hemiptera:Aleyrodidae) populations by isoelectric focusing electrophorisis. Biocham Genu,38 1/2) :14 –25.
- Brown J.K., D.R. Frohlich and R.C.Rosell, 1995. The sweet potato or silver leaf flies: Biotypes of *Bemisia tabaci* or a species complex ? Anna. Rev.Entomol. 40 : 511 – 534.
- Chandra, B.K.N., Rai,P.S. and shetty, B.V.1973. Protection of brinjal crop from attack of spider mite. Curr. Res., 2 (9): 75.
- Chen, H.G. 2000. Infestation by and of *Tetranychus*.in shanxi. Plant prot., 26 (2) : 29 –30 .
- Chinniah, C 1999. Evaluation of certain neem derivatives for their safety to predatory mites *Amblyseius sp.* (Acari : Phytoseiidae). Insect Environment 5 (1):38
- Chitra, vamsidhar 2000. Effect of *Annona squamosa* L. seed Extract on Protein Metabolism of *spodoptra litura* Fab. Insect Envirnment 6 (1) 39 – 40.
- Chitra. K.C. et al. (2000). Effect of *Annona squamosa* L. Seed Extract on protein metabolism of *Spodoptera litura* Fab. Imsect Environment. 6 (1) :39- 40.
- Cohen, S. (1990) Epidemiology of whitefly-transmitted Viruses. In Gerling, D.(ed.) *Whiteflies their bionomics, pest status and management, Intercept, Andover* pp. 211-226.
- Costa, H.S., Brown T.K., Sivasupramanism S., Bird J. 1993. Regional distribution, insecticide resistance and reciprocal cross between the 'A' and 'B' biotypes of *Bemisia tabaci*. Insect sci: Appl, 14: 255 – 266.

- De Barro P.J. Driver F. 1997. Use of RAPD PCR to distinguish the B biotype from other biotypes of *Bemisia tabaci* (Gennadius) (Hemiptera : Aleyrodidae). Austral. J. Entomol. 36: 149 – 152.
- Deshmukh, S.D. and Borle, M.N.1975. Studies on insecticidal properties of indigenous plant products. Indian J. Entomol. 37 (1): 11 –18
- Devraj Urs ; K.C. 1990. Efficacy of certain plant products in the control of brinjal red spider mite, *Tetranychus curticae* In : 4th national symp on Acarol, Kerala PP.68.
- Dhawan, A.K.Simwat, G.S.1992. Management of sucking pest of cotton with natural plant products. In proc. 1st National symp. Allelopathy in Agroecosystem, H.A.U.Hissar, PP.154 – 155.
- Dhoora, M.S. and Bindra, O.S.1980. Evaluation of different pesticides for control of *T.cinnabarinus* infesting brinjal in Punjab.Ind.J. Entomo. 42 (2) : 271 – 273.
- Dhooria, M.S. and Mann. M.S. 1980. Chemical control trials using ultra low volume and high valume sprays against spider mite, *Tetranychus neocaledonicus* infesting brinjal in Punjab. Ind. J.Entomo. 42(2): 269 – 270.
- Dhooria, M.S; Sandha, G.S.and Bhalla, J.S.1976. Effectiveness of different granular insecticides against mite, *Tetranychus neocaledonicus* infesting brinjal . Pesticides. 10 (12) : 47.
- Dilbagh – singh and Gill, C.K. 1981. effect of selective insecticides and fruit borer and mites in brinjal Punjab Veg. Grower. 15 –16 : 27 – 30.
- Dongare, T.K. and Rahalkar, G.W.1984. Ovipositional deterrent action in leaves of *Xanthium stramonium* (Compositae) tested against *E.vittella*. Indian J. Entomol. 46 (1) : 111 – 112 .

- Dongre, T.K. and Rahlkar, G.W. 1978. Antifeedant activity of *Xanthium Stramonium* against larvae of *Earias vittella* F. Proceedings of a symposium on "Nuclear Techniques on studies of Metabolism, Effect and Degradation of Pesticide", Department of Atomic Energy (In Press.)
- Fagoonee, I. 1987. Use of neem in vegetable crop protection in Maurius In: H. schmutterer and K.R.S. Ascher, Eds., Natural pesticides from the neem tree and other Tropical plants. GTZ, Eschborn, pp. 419 – 429.
- Gill R.J. 1992. Reviews of the sweet potato whitefly in southern California Pan-Pac. Entomol. 68 : 144 – 52.
- Gloria Nombela, Susana Pascual, Milagros Aviles, Emilif Guillard and Marians Muniz. 2005 Bemothiadiazole Induces Local Resistance to *Bemisia tabaci* (Hemiptera : Alyrodidae) in tomato plants. J.Econ. Entomol. 98 (6) : 226 – 227.
- Gopal, S. and Senguttuvan, T. 1997. Integrated management of tomato fruit borer with insecticides, neem products and virus. Madras Agril. J.84 (2) :82-84.
- Goyal, M. 1982. The spider mite, *Tetrangchus telarius* on glusshouse brinjal sci. and cult., 48 (6) : 220 – 221.
- Gujar, G.T. and Mehrotra, R.N. 1988. Biological activity of neem against red pumpkin bettle *Aulacophora foveiollis*, *Phytoparasitica* 16 (4) : 297- 302.
- Gupta S.K. 1985 Hand book of plant mites of India, Publication of Zoological survey of India, Calcutta pp. 106.
- Gupta, S.K; Sidhu, A.S. singh, G. and Dhooria, M.S. 1972. Evalution of new pesticides for the control of mite, *Tatracnychus carcurbitae*. Infesting brinjal in Punjab. Ind.J. Agri. Sci., 42 (10) : 921 – 942.
- Hari, Raghavendra. Rao A., Kameshwara Rao, and Mani, A. 1990. Management of gram pod borer *Heliothis armigera* (Hb) with newer insecticides in

- pigeon pea. Pp. 119-126. In: Heliothis Management Proceeding on National Workshop (Jayaraj, S. Uthamsamy, S., Gopalan, Mand Rabindra, R.J. Eds). Centre for plant protection studies, Tamil Nadu Agric. Univ., Coimbatore.
- Hiremath, I.G. Ahn Young Than and Kim Soonil 1997. insecticidal activity of Indian plant extracts against *Nilparvata lugens* (Homoptera), Delphacidae Appl. Entomol and Zool 32 (1): 156 – 166.
- Jayaraj, S. 1988. The problem of Heliothis management in India and its integrated management. Paper presented at Nat. Workshop on Heliothis management held at Coimbature, (India), on Feb. 18-19.
- Kashap, N. P., Gupata, V., Kaushal, A. N. 1974. *Mentha spicata* a promising protectant to stored wheat against *Sitaphilus oryzae* L.bull. Grain Technol. 12 (1) : 41-44.
- Kesularts, J.L.W. 1980. Effect of the vegetable leafminer, *Liriomyza sativa* Blanchard and the associated plant pathogens on yield and quality of the tomato, *Lycopersicon esculantum* Mill. Cv.Walter Ph.D.dissertation, Univ. of H. Florida, Gainesville, pp.153.
- Khan, Q., and Rao, V.P. 1960. Insect and mite pests. Cotton in India, Indian Central cotton committee, Bombay, 217 – 301.
- King, A.B.S. and Sawiski, R.M. 1990. Insecticide resistance of *Helicoverpa* and its management in chickpea in the nineties Proc. Ind International Workshop on chickpea improvement, ICRISAT Patancheru, India, pp. 195-201.
- Krishnaiah, K and Tandon, P.L. 1975. chemical control of *Tetranychus cucurbitae* infesting brinjal. Curr.Res. 4 (5) : 87 – 89.
- Kumar S.Jain, P.C.and kumawat, S.R. 1989. Field efficacy of various insecticides against the mite, *Tetranychus macfarlanei* Ind.J. Entomo., 51 (3) : 253 – 255.

- Martin, J.H., Aguiar, A.M.F. & Baufeld, P. (2001) *Crenidorsum aroidephagus* Martin & Aguiar sp. nov. (Sternorrhyncha:Aleyrodidae), a new
- Mathur, K.C. 1970. Four new records of parasites attacking *H. armigera* in Himachal Pradesh, India. Curr. Sci. 39(7) : 167.
- Mehrotra, K.N. and Phokela, 1991. *H. armigera* resistance to insecticides in India. Resistant Pest Management 3 (2) : 15-16.
- Mehta P.J.A. Wyman, M.K. Nakhla and D.P. Maxwell 1994. transmission of tomato yellow leaf curl germinivorous by *Bemisia tabaci*. J.Econ.Entomol. 87 : 1291 – 1297.
- Men, U.B. Sarode, S.B., 1999. Vertical Distribution of Whifefiles, *Bemisia tabaci* Genn. On Sunflower Insect Environment, Vol. 5(3) 111-112
- Menth spicata apromising protectant to stored wheat against *Sitophilus oryzae* L. Ball. Grain Technol. 12 (1): 41 – 44.
- Metabolism, Effect and Degradution of Pesticide", Department of Atomic Energy (In Press.)
- More , G.D. Kadu,N.R. Honda, H and Yamamoto I 1990. Pesticidal active substance in tropical plants insecticided substances from seeds of Annonaceae. J.agric Sci. Tokyo 34 (4) 253 – 258.
- More, G.D., Kadu, N.R. and sakhare, S.D. 1989. Evaluation of insecticidal properties of indigenous plant prodects. Mahazine, collage of Agri., Nagpur, PP. 49 – 59 .
- Mote, U.N., Dawkhar, R.V. AND Lolage G.R. 1995. Efficacy of imidaclorid as seed treartment against intial sucking pestsof cotton. Pestolgy xix (1) : 5-8.
- Murugan, K., B abu, R. and Sivaramakrishnan, S.1999. Toxic effect of plants on *Spodoptera litura* Fab. Insect environment 4 (4) 135.

- Murungan K. and Jeyabalan, D. 1998. Neem enhance the activity of Microbial Pesticides. Insect environment 4 (1) : 3 – 4. Narasimhan, V.and Mariappan, V. 1988. Effect of plant derivatives on green leafhooper and rice tungru transmission. Intern. Rice Res. News Letter. 13 (1) : 28-29.
- Narasimhan, V.A AND Mariappan, V. 1988. Effect of plant derivatives on green leafhooper and rice tungru transmission Intern. Rice Res. News.Letter. 13 (1) : 28 –29.
- Natarajan, K.1984. The cotton Whitefly problem and progress. Paper presented in South Zone panel meeting of AIICCP held at Dharwad. June 6 – 7, 1985.
- Oatman, E.R. and Planter, C. R. 1971. Biological control of tomato fruit worm on processing tomatoes in South California using mass release of *T. pretiasum*. J. Econ. Ent. 64 : 501-506.
- Oliveria : Henneberry and Anderson P. 2001. History, current status and collaborative research projects for *Bemisia tabaci* crop Prot. 20 : 709 – 723.
- Oudhia, P. 2000 a. Effects of leaf extracts on metallic coloured tortoise beetle, *Aspidomorpha miliaris* F. Insect Environment 5 (4) : 165.
- Oudiha, P. 2000 b. Toxic effect of *Parthenium* leaf extracts on *Apidomorph miliaris* F. and *Zonabaris pustulata* Thunb. Insect Environment. 5 (4) : 168.
- Pande, Y.D. and Sharma, A.K.1980. Evaluation of some pesticides for the control of *Tetranychus neocaledonicus* (ed. G.P.Channa – Basavanna). Contributions to acarol in in India. 207 – 208.
- Parihar, S.B.S, thapaliyal, P.and Ramkishor. 1999 Effect of botanicals on peach aphid, *Myzus persicra* (Sutzer). Insect Environment 5(2) : 84.
- Patanik, H.P. 1997. studies on neem formulations against serpentine leaf miner and tobacco cutterpillar on Insect Enviroment, 3 (1) : 10.

- Patel, B.K. Patel, R.C., Shah, A.H. and Mehta, N.P. 1987. Incidence of Whitefly on cotton in Gujurat state and action taken for its management. AICCIP workshop, Hyderabad, January, 1987.
- Patel, C.B. Shah, A.H. and Patel, S.H. 1982. Effect of insecticides on the resurgence of two spotted spider mites, *T. telarius* in brinjal. Ind. J. Agri. Sci. 52 (11) : 774 – 776.
- Patel, C.B., Rai, A.B., Patel M.B. Patel, A.J and shah, A.A. 1993. Acaricidal tests of botanicals in compurison to conventional insecticides jacarocodes against red spider mites on okra, brinjal and Indian bean. Ind.J. Entom., 55 (2) : 184 – 190.
- Patel, J.R., Borad, P. K., Shah, B.R. and Ratanapara, H.C. 1991. Need based control of *H. armigera* on tomato with endosulfan. Indian J. PL. Prot. 19 (1) : 79-80.
- Pawar, D.B., Lawande, K. E. and Warade, S.D. Maharashtra Agric. Univ. 21 (1) : 165-166.
- Pawar, J. G. and Rathore, V.S. 1959. Mites are big thieves of crops . Ind. Farm., 9 : 13-16.
- Perring T.M., 2001. The *Bemisia tabaci* species complex. Crop Prot. 20: 725 – 737.
- Perring, T.M., Cooper A.D., Rodriguez, R.J. Farrar C.A., Bellows TS. 1993. Identification of a whiteflies species by genomic and behavioral studies, Science. 259 : 74 – 77.
- Powell, R.G. 1989 Higher plant as a source of new insecticidal compounds. Pesticide Science 27(2) : 228 – 229.
- Prasad, V. (1974), “ A catalogue of mites of India”. Indian A car. Pub. House Ludhiana , pp 320.
- Puri, S.N. Dhanarkar, B.K. Bilopate, G.G. and Daware D.G. 1986. Present status of cotton whitefly. Seminar on status of whitefly on cotton. Pune, March 14.1986.

- Puttaradrich, M.. 1974. Field control of the leaf mite of Jola. Mysore Agri. J., 26:17-19.
- Rahman, K.A. 1939. Insect pest numbers, Lyallpur. Punjab Agri. College Magazine 7: 1-82.
- Rao, S.M., Chitra, K.C., Gunesekhar, D. and Rao, P.K. 1999. Antifeedent properties of certain plant extracts against second stage larvae of *Henosepilanhna vigintioctopuntata* Indian J. Entomol, 52 (4) : 681-685.
- Reddy, A.S. Rosaiat B. Rao, T.A. Rao R.B. and Rao, V.N 1986. The problem of whitefly on cotton. Paper presented in seminar on 'Problems' of whitefly on cotton, held at collage of Agriculture, Pune on March 14 , 1986.
- Rupprecht, J.K., Alkofahi, A., Anderson, J.L. McLaughlin, K.L. Mikolajczak and B.A. Scoot (1990). Search for New Pesticides from Higher Plants. ACS Symp. Ser.387. Amercial Chemical Society, Washington, D.C. PP. 25-43
- Russell L.M. 1951. Synonyms of *Bemisia tabaci* (Gennadius) (Hemopopatera : Aleyrodidae) Bull. Brooklyn. Entomol.Soc. 52: 122 –123.
- Sahayaraj,K. and Gabriel Pavraj. M.1998. Effect of neem leaf extract on *Amsacta albistriga* walker. Insect Environment. 4(2) : 42 – 43 .
- Saito M. L. Oliveira, F. Fell, D. and Takematsu, A.P. 1989. Evaluation of the insecticidal activity of some Brazilian plants. Arquiovos do instituto Biologico, Brazil 56(1/2) : 53 – 59.
- Sandhu G.S. & Verma, G.G., Indian. J.PI. Prot., 3 (2), (1975). 207-08
- Sandhu, G.S. & Darshan Singh, Indian J. PI. prot., 3(2). (197-5). 177-08
- Santharam, G., Blasubramanian, M. and Chelliah, S.1981. Control of H.armigera on Red gram 9Cajanus cajan) with a NPV and insecticide. Madras Agric. J. 68 (5) : 417-420.

- Sardana, H.R. and Kumar, N.K.K. 1991. Effectiveness of plant oils against leafhooper and shoot and fruit borer on okra. Indian J. Entoml. 53 (1) : 57.
- Saxena, R.C. Liquido, N.J. & Juito, H.D. Neem Seed oil – Aitfedent growon plant hooper – paper presented at Annual conference control, council of Philippines, Mania. 1979
- Shah, A.H., Prajapati, B.G. and Patel, C.B. 1989. Field evaluation of acaricides against mites of brinjal, red gram and Indian bean.. In : G.P. Channa Basavanna and C.A. virktamath: Progress in Acarol, Oxford and IBH Pub. Co, New Delhi 2 : 432 – 437.
- Shin – foon, C. 1987. Experiments on the practical application of china berry, *Melia azedarch* and other naturally occurring insecticides in China. In: H Schmutterre and K.R.S. Ascher, Eds. Natural pesticides from the neem tree and other tropical plants. GTZ. Eschborn, pp. 661 – 688.
- Shrinivas, K and Krishnamoorthy, P.N. 1993. pest management in solanaceous vegetables. Report of Research Activites Published in Annual Report of Research Activites Published in Annual Report – 1992-1993. Indian Institute of Horticultural Research. Hessarghatta, Bangalore 7.2.1 : 59.
- Siddhia, S.A. 1987. A proposed pest management programme including neem treatments combating potato pest in the sudan. In:H Schmuttere and K.R.S. Ascher, Eds, Natural pesticides from the neem tree and other Tropical plants. GTZ., Eschborn, pp. 449 – 459
- Sidhu, A.S. and Singh, G. 1971. chemical control of *Tetranychus cucarbitae* on brinjal. Punjab Hort. J. 11 (1&2) : 95 – 97.
- Singh, R.P. Devakumat, C and Dhingra, S. 1988. Activing of neem (*Azadirachta indica A.Juss*) seed extracts against the mustard aphid, *Lipaphis erysimi* *Phytoparasitica* 16(3): 225-230

- Singh, R: verm, A.N and verma, N.D. 1975. chemical control of *Tetranychus cacurbitae* on brinjal. Acarol., 17 (1) : 92 – 94.
- Singh, S. and Guram, M. 1960. Cotrol of vegetable mites in Punjab and Uttar Pradesh. Sci. and Cult., 32 : 51 –53.Kaneria, V.S. and R. K. Bharodia. 1991. Efficacy , relative toxicity and residual persistent toxicity of different pesticides to adults of *Tetranychus cinnabarinus* infesting brinjal crop Gujarat, Agri. University RES., J. 17 (1) : 54- 59.
- Sithanantham, S.1987 Insect pests of pigeonpea and chickpea and their management. In : Plant Protection in Field crops. Pp. 159-173.
Veerbhadr Rao, M. and Sithanantham (Eds.) Pl. Prot. Assoc. India, Hyderabad, India.
- Sood, A.K., P.K.Mehta and S. Parmar 2000. Evaluation of plant extract against cabbage aphid, *Brevicoryne brassicae* (linn). Indect Environment, 6 (2) 87 – 88.
- Stinner, R.E., Ridgway, R.L. and Morrison, R.K.1974a. Parasitsm of *Helithis* eggs after field release of *T. pretiosum* in cotton Environ. Ent. 3 : 498 – 500.
- Stinner, R.E., Ridgway, R.L. and Morrison, R.K.1974b. Longevity, fecundily and searching capacity of T. PRETIOSUM BY 3 METHODS environ. Ent. 3 : 558
- Subba Rao, P.V; Krishnamurthy, M.M; Punnaiah, K.C.Reddy, G. P.V; Mayuravalli, V.V.L. and Subbaiah, K.R.1984 Relative efficacy of certain insecticides for controlling brinjal pests. Andhra Agri., J. 31 (3) : 242 – 243.
- Sudhakar, K; Punnaldh, R.C. and krishnayya, P.V. 1998. efficacy of cerlatin selected insecticides on the sucking pest complex of brinjal. Ind.J Entomo, 60 (3):241 –244.

- Swain T1977. Secondary compounds as protective agents Arev. Pl. Physiol. 28 : 479 – 501.
- Thakare, H.S., Gawande, R.B., Satpute U.S., Saroda, D.B. and Aherkar, S. K. 1986. Status paper on “ Problem of whitefly on cotton, in Vidharbha region” seminar on status of whitefly on cotton Pune, March 14 ,1986.
- Vaneria, V.S. and R.K. Bharodia. 1991. Effecacy, relative toxicity and residual persistent toxicity of different pesticides to adults of *Tetranychus cinnabarinus* infecting brinjal crop. Gujrat Agri. University Res, J. 17 (1) : 54 – 59.
- Verma, S. 1992. Persistence of insecticides against insect and non – insect pest complex of brinjal. Ind. J. Entomo, 54 (4) : 415 – 149.
- Vevai E.J. and Talgeri, G.M. 1948. Bombay crop pest Calender and a seasonal schedule of their control by Modern insecticides J.Bombay Natu Hist Soc V 48 : 725.
- Walung, A.R. and Mote, U.N. 1995. Evaluation of imidacloprid against thrips and whitefly on tomato Pestology XIX (11) : 21-23.
- Webb, Ralphi E. Mark A. Hinebaugh, Richard, K. Lindquist and Martin Jucobson 1983. Evaluation of aqueous solution of neem seed extract against *Liriomyza sativa* and *L. trifolii*. (Diptera : Agromyzidae) . Econ. Emt. 76 : 357 – 362.
- Yadav, G.S. Anand, R.K. and Yadav, P.R. 1989. Bioefficacy of some chemicals against red spider mites on brinjal. Ind.J. Entomol. 49 (4) : 582 – 584.