

Chapter VI

Bibliography

Abdul-Baki., A. A. and Anderson, J. D. (1973). Vigour determination in Soybean seed by multiple criteria. *crop sci.* **13**: 630–633.

Agarwal, P. C., Dev, U., Singh, B., Indra, R. and Khetarpal, R. K. (2006). Seed-borne fungi detected in consignments of soybean seeds (*Glycine max*) imported into India. *Bulletin OEPP*. **36**(1): 53-58.

Agarwal, P. K. and Menon, S. K. (1973). Note on appearance of cracking and wrinkling on seed coat during seed development in some spring grown varieties of soybean (*Glycine max* (L.) Merrill). *Seed research*. **1**: 98-100.

Agarwal, R. L. (1980). *Seed technology*. Oxford and IBH Publishing Co. New Delhi. 449-507.

Agarwal, S. and Singh, T. (2002). Plant latices as biopesticide against seed-borne fungi of okra. *Journal of Mycology and Plant Pathology*. **32**(1): 135.

Agarwal, V. K. and Joshi, A. B. (1971). A preliminary note on the purple stain disease of soybean. *Indian phytopathology*. **24**: 810–812.

Agarwal, V. K. and Sinclair, J. B. (1993). *Principles of Seed pathology*. CBS publisher and distributors, Delhi. 1-158.

Agarwal, V. K. and Singh, O. V. (1974). Fungi associated with Sunflower seeds. *Indian Phytopathology*. **27** (2): 240-241.

Barnett, H. L. and Hunter, B. B. (1973). *Illustrated Genera of Imperfect Fungi*. Burgess Publishing Company. Minneapolis, Minn., USA 2nd ed., 1-241.

Basak, A. B. and Woong, L. M. (2002). Prevalence and transmission of seed-borne fungi of maize grown in a farm of Korea. *Mycobiology*. 30(1): 47-50.

Bedi, P. S. and Kapoor, K. S. (1974). Mycoflora of pea and its associative effect on the crop. *J. Res. Pb. Agri. Univ.* 11: 284-288.

Beebi Rezeena, P. M. and Ahmad, R. (2007). Control of seed borne fungi of okra with *Pseudomonas fluorescens* and aqueous leaf extract of Heena (*Lawsonia inermis*, L.) *Journal of Mycology and Plant Pathology*. 37(3): 485-487.

Bhadraiah, B. and Ramarao, P. (1987). Fungi from stored sorghum seeds. *Indian Phytopathology*. 40: 401- 404.

Bhatia, A., Singh T. and Singh D. (1995). Seed mycoflora of guar in Rajasthan and its phytopathological effects. *Journal of the Indian Botanical Society*. 74: 329-330.

Bhowal, S., Borah, P. K., Senapaty, D. and Rabha, B. K. (2006). Seed-borne microflora in groundnut and their management. *Indian phytopathology*. 59(3): 398.

Bilgrami, K. S., Jamaluddin, T. P. and Roy, A. K. (1976). Studies on the deterioration of some pulses by fungi. *Indian Phytopathology*. 29 (4): 374-377.

- Boroduline, U. S. (1941). Development of the natural microflora of the seed coat on the roots of the cultivated plants and its effects on the bacterization of seeds, *Microbiology. USSR*, **10**: 924.
- Ch. Ramesh. and Avitha, K. M. (2005). Presence of external and internal mycoflora on sunflower seeds. *J. Mycol. Pl. Pathol.* **35**(2): 362-364.
- Chandrani, A. N. and Paul, Y. S. (1993). Biodeterioration of French bean seed due to seed microflora. *Indian Phytopathology.* **46**(3): 283
- Chandrani, A. N. and Paul, Y. S. (1993). Detection of seed borne microflora of French bean (*Phaseolus vulgaris L.*) in Himachal Pradesh. *Indian Phytopathology.* **46**(3): 283
- Chandvaria, M. S. (1991). *Acreage Response of Oil Seeds in India.* Radha Publications New Delhi. 1-198.
- Charya, M. A. S. and Reddy, S. M. (1980). Effect of *Cajanus cajan* seed coat leachates on germination of some seed borne fungi. *Indian Phytopathology.* **33**: 112.
- Chen, C. C. (1920). Cf. *Principles of seed pathology.* Agarwal, V. K., Sinclair, J. B., Vol. I, CBS publisher and distributors. Delhi. 2p, 1993.
- Chenulu, V. V., Singh, A. and Joshi, L. M. (1970). Cf. *Seed Pathology.* Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 9p, 1977.
- Chiarappa, L. and Gombogi, P. (1986). Seed pathology and food production. *FOA Plant Prot. Bull.* **34**: 167-185.

Flannigan, B. (1970). Comparision of seed-borne mycoflora of barley, oats and wheat. *Trans. Brit. Mycol. Soc.*, **55**(2): 267-276.

Frank. A. B. (1885). Cf. *Principles of seed pathology*. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS publisher and distributors. Delhi. 3p, 1993.

Gachande, B. D., Mandge, S. V. and Mukadam, D. S. (2007). Nutritional impact on extracellular protease activity of some seed-borne fungi in Legume. *Geobios*. **34**(1): 229-232.

Gawai, P. B., Solanke, R. B., Deosarkar, D. B. and Jawale, L. N. (2000). Studies on seed mycoflora of maize and its impact on seed quality. *Journal of Maharashtra Agricultural Universities*. **25**(2): 222-223.

Goulart, Augusto. C. P., Paiva, Fernando. De. A., Andrade, Paulino J. M. (1995). Control of soybean seed-borne fungi by fungicide treatments. *Summa Phytopathologica*. **21**(3-4): 239-244.

Grover, R. K. and Bansal, R. D. (1971). Seed-borne nature of *Collectotrichum capsici* in chilli seeds and its control by seed dressing fungicide. *Indian Pythopathology*. **24**: 664-668.

Gupta, A. (2001). Role of seed mycoflora in deterioration of soybean seed during storage under ambient condition. *Indian phytopathology*. **54**(4): 503.

Gupta, A., and Aneja, K. R. (2001). Mycofloral spectrum during storage and its effect on seed viability of soybean (*Glycine max* (L.) Merrill) seeds under ambient conditions. *Proceedings of the National Academy of Sciences India Section B (Biological-Sciences)*. **71**(3-4): 245-253.

Gupta, D. K. and Chaudhary, K. C. and Basu. (1995). Seed borne fungi of bhindi, brinjal and chillies grown in Sikkim. *Indian Journal of Mycology and Plant Pathology*. **25**(3): 282-283.

Gupta, V. K. and Chouhan, J. S. (1970). Seed-borne fungi and seed health testing in relation to seedling diseases of groundnut. *Indian Phytopathology*. **23** (4): 622–625.

Gupta, V. K. and Chouhan, J. S. (1970). Seed-borne fungi and seed health testing .in relation to seedling disease of groundnut. *Indian Phytopathology*. **23**: 622– 625.

Hashmi, F. M. H. and Ghaffar, A. (2006). Seed-borne mycoflora of wheat, sorghum and barley. *Pakistan Journal of Botany*. **38**(1): 185-192.

Hydar-Ali, M. M., Fakir, G. A. (1993). Control of seed-borne fungi of wheat with fungicides. *Bangladesh Journal of Botany*. **22**(2): 135-141.

ISTA. (1966). International Rules for Seed Testing. *Seed Sci Technol*. **24**: 335–337.

Jain, J. P. and Patel, P. N. (1969). Seed mycoflora of guar, their role in emergence and vigour of seedling and efficiency of fungicide. *Indian Phytopathology*. **22**(2): 245–250.

Jha, D. K. (1993). *A Text book of Seed Pathology*. Vikas Publishing House Pvt. Ltd.1-72.

- Joshi, B. C., Srivastava, S. K. and Prajapti, S. K. (1987). Studies on detection of mycoflora and mycotoxins in paddy harvested from rain affected crop in Haryana. *Bull. Grain tech.* **25**: 188-190.
- Jot, M. B., Mahajan, P. D., Gaikwad, D. G. and Ekbote, M. V. (1991). Documentation of different seed-borne fungi of sorghum and their effect on seed germination. *Indian phytopathology*. **44**: 17.
- Kadian, O. P. and Suryanarayana, D. (1971). Studies on seed microflora of oilseed crops. *Indian Phytopathology*. **24** (3): 487-490.
- Kamat, M. N. (1959). *Hand Book Of Mycology*. Part I, *Phycomycetes* and *Ascomycetes*. Prakash Publishing House, Poona.1-183.
- Karmakar, S. S., Subuddhi, S. and Bandopadhyay, B. (1980). Studies of seed microflora of soyabean. *Indian Jour. Microbiol.* **20** (3): 236-238.
- Karwasra, S. S. and Singh, M. (1982). Seed mycoflora of cluster bean in Haryana and their control by seed treatments. *Indian Phytopathology*. **35** (3): 501-502.
- Kavita rani, M. M. and Mukergi, K. G. (1978). Studies on seed borne fungi, I. Occurrence of three pathogenic fungi on sorghum seeds. *Seed Research*. **6** (I): 38-42.
- Khayum, A. S., Gopal. K., Prasad baby, G. and Munikrishnaiah. (2005). Fungicidal control of seed-borne fungi in Soybean. *Geobios*. **32**(4): 225- 228.
- Konde, B. K. and Pokharkar, B. R. (1979). Seed borne fungi of Sorghum. *Seed Research*.**7**(I): 54-57.

Krishnapp, M. and Deepak, R. (2004). Studies on seed borne fungal diseases of French bean (*Phaseolus vulgaris*, L.) in Karnataka. *Journal of Mycology and Plant Pathology*. 34(3): 982-983.

Kumar, I. K., Prasad, U., Manoharachary, C. and Rao, R. P. (1991). A survey of grain moulds of sorghum. *Indian phytopathology*. 44: 10.

Kumar, M. and Agarwal, V. K. (1998). Location of seed-borne fungi associated with discolored maize seeds. *Indian phytopathology*. 51 (3): 247-250.

Kumar, P., Saifulla, N. M. and Byre Gowda, M. (2004). Management of seed mycoflora of pigeon pea. *Journal of Mycology and Plant Pathology*. 34(3): 980-981.

Lal, S. P. and Kapoor, J. N. (1979). Succession of fungi in wheat and maize during storage. *Indian Phytopathology*. 32 (1): 101-104.

Lalithakumari, D., Govindaswamy, C. V. and Vidhyasekaran, P. (1971). Cf. *Seed Pathology*. Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 24p, 1977.

Lokesh, M. S. and Hiremath, R. V. (1993). Effect of relative humidity on seed mycoflora nutritive value of red gram (*Cajanus cajan* Linn). *Mysore J. Agri. Sci.* 27(3): 268-271.

Mall, O. P., Pateria, H. M. and Chauhan, S. K. (1986). Microflora and Aflatoxin in wet harvest Sorghum. *Indian Phytopathology*. 39 (3): 409-413.

Mathur, R. L. and Sharma, L. C. (1971). Field assay of seed dressing fungicide for groundnut seeds. *Madras Agric. J.* **58**: 199–202.

Mathur, S. K., Mathur, S. B. and Neergaard, P. (1975). Detection of seed – borne fungi in sorghum and location of *Fusarium moniliforme* in seed. *Seed Sci. and Technol.* **3**: 783–790.

Medidar, H. S., Joi, M. B. and Kshirsagar, C. R. (1991). Isolation of seed-borne fungi of wheat and their chemical control. *Indian phytopathology*. **44**: 106.

Mehta, P. R., Singh, B., Mathur, S. C. and Singh, S. B. (1953). Cf. *Seed Pathology*. Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 18p, 1977.

Meiri, H. A. (1973). Infection of chickpea seed by *Ascochyta ruberi* in Isreal. *Pl. Dis. Repr.*, **54**: 442–445.

Meiri, H. A. and Cohen, R. (1992). Mycoflora of sunflower seeds. *Hassadeh (Isreal)* **72** (12): 1453-1456.

Mishra, R. R. and Kanaujia. (1973). Studies on certain aspects of seed-borne fungi - II. seed-borne fungi of certain oil-seeds. *Indian Pythopathology*. **26** (2): 284 –294.

Mittal, R. K. and Mathur, S. B. (1993). *Pathology*. Part I, ICAR, New Delhi, India and Danish Government Institute of Seed Pathology, Denmark, 178p.

Mukergi, K. G., Bhatnagar A. K., Tripathi, S. C., Bansal, M., Saxena, M. (1992). *Current concept in seed biology*. Naya Prokash Culcutta. 213p.

Muley, S. M. and Baig, M. F. (2007). Study on mycoflora of oil seeds. *Bioinfolet*. 4 (2): 146–150.

Murthy, G. P., Raghavendra, V. B. (2007). Effectiveness of fungicides on seed borne mycoflora, seedling vigour and bio-chemical constituents during early emergence of bhendi (*Abelmoschus esculentus*, L. Moench) cv. Kiran. *Advances in Plant Sciences*. 20(1): 261-263.

Muskett, A. E. and Melone, J. P. (1941). Cf. *A Text book of Seed Pathology*. Jha, D. K . CBS publisher and distributors. Delhi. 57p. 1993.

Nahar, S., Mushtaq, M., and Hashmi, M. H. (2005). Seed-borne mycoflora of sunflower (*Helianthus annuus* L.). *Pakistan Journal of Botany*. 37(2): 451-457.

Nahar, S., Mushtaq, M., Pathan, I. H. (2004). Seed-borne mycoflora of *Capsicum annuum* imported from India. *Pakistan Journal of Botany*. 36(1): 191-197.

Nair, N. N. and Arora, R. (1994). Cultural characterization of some seed-borne fungal pathogens of legume crops. *Geobios*. 21(2): 105-108.

Narasimham, K. S. and Rangaswami, G. (1968). Effect of some fumigants on the microflora of sorghum seeds. *Indian Pythopathology Soc. Bull.* 4: 57–63.

Padule, D. N., Mahajan, P. D., Perane, R. R. and Patil, R. B. (1999). Efficacy of fungicides for increasing storability of grain moulds infected seed of sorghum hybrid CSH 14. *Seed Research.* 27(1): 95-99.

Pandey, U. (1976). Fungi associated with seeds of chillies grown in Kumaon Hills. *Indian Phytopathology.* 29: 472.

Pandey, V., Kumar, N. and Tripathi, N. N. (2007). Inhibition of fungal deterioration of pigeon pea seeds by Cuminum cymimum oil. *Indian phytopathology.* 60(3): 306-312.

Parveen, S., Kumar, V. R. (2000). Seed-borne mycoflora of wheat collected from Rajasthan with special reference to *Alternaria* species. *Journal of Phytological Research.* 13(2): 183-186.

Paul, M. C. and Mishra, R. R. (1991). Evaluation of fungicides against seed borne fungi of maize. *Indian phytopathology.* 44: 13-14.

Paul, M. C. and Mishra, R. R. (1992). Studies on seed mycoflora of maize: I. Seasonal variation in mycoflora. *Crop Research (Hisar).* 5(SUPPL.): 225-232.

Paul, Y. S. (2002). Biodeterioration of Frenchbean seed and its management. *Journal of Mycology and Plant Pathology.* 32(2): 167-173.

Pawar, N. P., Solanke, R. B., Mirza, F. N., Baig and Somwanshi, S. D. (2005). Studies on seed-borne fungi of chilli (*Capsicum annuum L.*). *Indian phytopathology.* 58(3): 357.

Pensalwar, S. N., Solanke, R. B. and Kore, S. S. (1996). Seedborne fungi of Sunflower. *Phytopathology.* 10: 18-25.

Randhawa, H. S. and Aulakh, K. S. (1994). The effect of maturity on the incidence of seed borne fungi of wheat. *Indian phytopathology*. **47**(3): 308.

Ranf, B. A., Bhatti, M. H. R. and Ahmed, I. (1997). Effect of seed – borne fungi on oil content and fatty acid profile in sunflower. *Helia* 20, Nr. 27. pp 67-72.

Rangaswami, G. and Rao, A. V. (1957). Alternaria blight of cluster beans. *Indian Phytopathology*. **10**: 18- 25.

Rao, P. H. and Reddy, S. M. (1987). Influence of humidity on seed mycoflora and deterioration of sorghum seed during storage. *Indian Journal of Mycology and Plant Pathology*. **17**(1): 6-10.

Rao, P. Hareesh Vardhan and Reddy, S. M. (1987). Influence of humidity on seed mycoflora and deterioration of sorghum seed during storage. *Indian J. Mycol. Pl. Path.* **17**: 6-10.

Rao, P. Hareesh vardhan., Giridhar, P. and Reddy, S. M. (2000). Fungal succession on Sorghum seed harvested at different stages of maturity. *Indian phytopathology*. **53**(1): 102-104.

Rao, P. Hareesh Vardhan., Giridhar, P. and Reddy, S. M. (2001). Effect of culture filtrates of some seed borne-fungi on seed germination and seedling growth of Sorghum. *Proceedings of the National Academy of Sciences India Section B (Biological-Sciences)*. **71**(1): 63-65.

Rao, P. Hareesh vardhan., Giridhar. P. and Reddy, S. M. (1996). Effect of volatile compound on seed mycoflora of sorghum. *Indian phytopathology*. **49**(1): 37–40.

Rathour, R. and Paul, Y. S. (2004). Pathogenicity and Management of seed mycoflora of pea. *Journal of Mycology and Plant Pathology*. **34**(2): 456–460.

Reddy, A. and Muhammad, S. (2004). Seed mycoflora of pigeonpea. *Indian phytopathology*. **57**(3): 386.

Reddy, S. H., Sugunakar, M. and Reddy, B. M. (1998). Effect of seed treatment with fungicides and insecticides on seed borne fungi, storage insect pest, seed viability and seedling vigour of groundnut. *Seed Research*. **26**(1): 62-72.

Richardson, M. J. (1979). Cf. *Principles of seed pathology*. Agarwal, V. K., Sinclair, J. B., CBS publisher and distributors. Delhi. 3p. 1993.

Rizvi, G. (2006). Deterioration of seed quality in relation to vigour by seed mycoflora in forage Sorghum. *Indian phytopathology*. **59**(4): 515–517.

Rodriguez, C., Robertson A. E. and Kanobe, C. (2007). Incidence of seed-borne *Fusarium* spp. in commercial maize (*Zea mays*) seed lots. *Phytopathology*. **97**(7): S100.

Rout, B. K. and Rath, G. C. (1972). Note on seed borne diseases of chilli (*Capsicum annuum*). *Indian Phytopathology*. **25**: 597–598.

Sandikar, S. N., Gaikwad, D. B. and Jamge, D. H. (1994). Impact of various storage methods on the composition of seed mycoflora in soybean (*Glycine max*). *Indian phytopathology*. **47**(3): 294.

Satyanarayana, B., Reddy, B. M. and Hussaini, S. H. (1993). Effect of incidence of seed-borne fungi on seed size and germination in sorghum. *Seed Research*. **21**(1): 55-58.

Sawhney, G. S. and Aulakh, K. S. (1981). Fungi associated with different grades of pea seeds and their pathogenic potential. *J. Res. Pb. Agri. Univ.* **18**: 37-40.

Schneidar, R. W., Thapliyal, P. N. and Sinclair, J. B. (1971). Fungi associated with soybean seed from India. *Indian phytopathology*. **24** (4): 792-793.

Shabbir, M. S. and Rajasab, A. H. (2004). Diversity of *Fusarium* species on sorghum grain. *Indian phytopathology*. **57**(4): 450-453.

Shafique, S., Shafique, S. and Arshad, J. (2005). Fungitoxicity of aqueous extracts of allelopathic plants against seed-born mycoflora of maize. *Mycopath*. **3**(1-2): 23-26.

Shailbala. and Tripathi. (2004). Seed treatment with fungicides and biocontrol agent on pathogens in urdbean seeds. *Journal of Mycology and Plant Pathology*. **34**(3): 851-852.

Sharma, J. R. and Basuchaudhary, K. C. (1975). Assessment of seed mycoflora of pearl millet and their control. *Indian Pythopathology*. **28** (3): 388-390.

Sharma, P. N., Basandrai, A. K. and Sharma, O. P. (1992). Seed mycoflora of buck wheat and its pathogenic potential. *Seed Research.* **20**(2): 168-169.

Sharma, P., Singh S. D. and Gour, H. N. (2004). Pathogenic potential and management of seed microflora of pea (*Pisum sativum*, L.). *Journal of Mycology and Plant Pathology.* **34**(3): 965.

Sharma, R. B. and Roy, A. N. (1991). Seed-borne fungi of Urdbean. *Indian phytopathology.* **44**(1): 64-67.

Singh, K. and Singh, A. K. (2004). Detection of seed mycoflora of gram (*Cicer arietinum*). *Indian phytopathology.* **57**(3): 361.

Singh, K. T., and Singh, D. (1987). Seed mycoflora of corn from Trible areas of Rajasthan, *Indian Jour. of Mycology and Plant Pthology.* **30** (1): 70-72.

Singh, O. K., Agarwal, M. N. and Nene, Y. L. (1973). Seed health studies in soybean raised in the Nainital, Tarai. *Indian Pythopathology.* **26** (2): 260– 267.

Singh, R. P., Singh, A. K. and Singh, K. (2006). Effect of fungicidal seed treatment on seed-borne fungi and germination of chilli. *Indian phytopathology.* **59**(3): 388-389.

Singh, R. P., Singh, A. K. and Singh, K. (2007). Seed-borne mycoflora of Chilli and their management. *Review of Plant Pathology.* **86**(3): 322.

- Singh, S. D., Rawal, P. and Bhargava, N. K. (2004). Pathogenic potential and control of seed mycoflora of Groundnut (*Arachis hypogaea*). *Journal of Mycology and Plant Pathology.* **34**(2): 687–690.
- Singh, T. (1993). Effect of mixed fungal infection in soybean seed. *Indian phytopathology.* **46** (3): 313.
- Solanke, R. B. and Kore, S. S. (1991). Seed-borne fungi of sunflower and response of *Alternaria alternata* to various concentration of fungicidal seed treatment. *Indian phytopathology.* **44**: 103.
- Solanke, R. B., Kore, S. S., Pensalwar, S. N. (1997). Effect of fungicidal seed treatment on seed viability and mycoflora during storage of sorghum. *Journal of Mycology and Plant Pathology.* **27**(3): 297-298.
- Solanke, R. B., Kore, S. S., Sudewad, S. M. (1998). Detection of soybean seed borne pathogens and effect of fungicides. *Journal of Maharashtra Agricultural Universities.* **22**(2): 168-170.
- Sowell, Jr. G. (1965). The effect of seed treatment on seed-borne pathogen of guar, *Plant Dis. Rep.* **49**: 895–897.
- Subbaiah, P. V., Shetty, H. S. and Safeeulla, R. M. (1982). Incidence of seed-borne fungi in maize (*Zea mays L.*) and their significance. *Indian Jour. Microbiol.* **22** (1): 57–60.
- Subramanian, C. V. (1961). *Hypomycetes*. ICAR Publication New Delhi.1-917 (Thom and Raper key included in this book)

- Sud, D., Sharma, O. P. and Sharma, P. N. (2005). Seed mycoflora in kidney bean (*Phaseolus vulgaris* L.) in Himachal Pradesh. *Seed Research.* **33**(1): 103-107.
- Suhag, L. S. and Suryanarayana, D. (1975). Some aspect of seed health testing with respect to seed-borne fungi of pulse crops grown in Haryana. *Indian Journal of Mycology and Plant Pathology.* **6:** 32-36.
- Suryanarayana, D., and Bhombe, B. B. (1961). Studies on fungal flora of some vegetable seeds. *Indian Phytopathology.* **14:** 30-41
- Swarup, J. and Mathur, R. S. (1972). Seed microflora of some Umbelliferous species. *Indian Phytopathology.* **25** (1): 12.
- Tamuli, P., Nath, M. J. (2007). Seed mycoflora of wheat and its control. *Geobios.* **34**(1): 93-94.
- Tandon, R. N. (1968). *Mucorales of India.* Indian Council Of Agricultural Research New Delhi. 1-120.
- Tariq, M., Dawar, S. and Mehdi, F. S. (2005). Effect of different moisture and storage temperature on seedborne mycoflora of soybean. *International Journal of Biology and Biotechnology.* **2**(4): 947-950.
- Tariq, M., Dawar, S., Abid, M. and Shaukat, S. S. (2005). Seed borne mycoflora of soybean. *International Journal of Biology and Biotechnology.* **2**(3): 711-713.

Thakur, M. P. (1999). Effect of fertilizer doses on the seedborne mycoflora of soybean. *Journal of Mycology and Plant Pathology*. **29**(3): 393-397.

Thippeswamy, T. and Lokesh, S. (1997). Efficacy of different seed dressing chemicals on seed mycoflora, seed germination and seedling vigour of sunflower (*Helianthus annuus* L.). *Seed Research*. **25**(1): 64-67.

Thoke, R. B. (1989). *Fungi of Maharashtra*. Ph.D. Thesis. Pune university. Poona.

Tillet, D. M. (1775). Cf. *Principles of seed pathology*. Vol I, Agarwal, V. K., Sinclair, J. B., CBS publisher and distributors. Delhi. 2p, 1993.

Toorray, N. K., Verma, K. P., Thakur, M. P. and Sinha, A. K. (2005). Evaluation of chickpea accessions by standard blotter method. *Advances in Plant Sciences*. **18**(1): 31-37.

Varshney, J. L. (1990). Seed borne diseases of Wheat– their impact in relation to production and productivity. *Int. J. Trop. Dis.* **8**: 173 – 192.

Varshney, J. L., Ganar, H. D., Gaur, A. and Sharm, S. P. (2002). Impact of plant product on quality of carry-over seeds of sorghum under storage. *Indian phytopathology*. **55**(3): 362.

Verma, V. (1984). *A Text Book of Economic Botany*. Emkay Publications Delhi. 1-196.

Vidhyasekaran, P., Thulasidas, G., Ramaswamy, K. R. and Kandasway, T. K. (1980). Preservation of viability of sorghum seeds by controlling seed-borne fungi. *Indian Phytopathology*. **33**: 225–230.

Vishawanathan, R. (1996). Seed mycoflora composition in sunflower. *Indian phytopathology*. **49**(3): 286–289.

Wallen, V. R. (1965). Cf. *Seed Pathology*. Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 832p. 1977.

Yadav, D. S. (1992). *Pulse Crops*. Kalyani Publishers New Delhi. 1-90.

Zaidi., B. I. S., Khan, M. I. and Saxena, S. K. (1991). Effect of fungicide on mycoflora of Chickpea seeds. *Indian phytopathology* **44** (3): 394–395.
