

---

**CHAPTER - VI**

**BIBLIOGRAPHY**

---

## BIBLIOGRAPHY

Abott, W.S. 1925. A method of computing effectiveness of an insecticide, J. Econ. Entomol. 18 (1) : 265 – 267.

Acharya. A 1985 a control of *Melodogyne incarnate* on betelvine with neem coke and sawdust under field conditions Proc. IV. Nematology symposium . Mahanpal Sukhadra Univ. Udupur (India)

Acharya. A 1985 pl. Pathogenicity of *Melodogyne incognita* on betelvine Proc. IV Sukhadra Univ. Udupur (India)

Acharya. A., and Padhi N.S. 1988. Effect of neem oil cake saw dust against root knot nematode, *Melodogyne incognita* on betelvine (*Piper betle* L). India J. Nematol., 18 : 105 – 106.

Agnihotria, V.R., Sen, C and Srivastava, 1975. Role of Fungitoxicants in the control of sclerotium root rot of sugar beet *Beta Vulgaris* L. Indian J. Ex PH. Biol. 13 : 89-91.

Alam, M.M., 1989., Control of root-knot and stunt nematodes with horn meal, bore meal and oil seed cakes, Indian J. Nematol. (9(2)) : 166-170.

Albuquerque, F.C. 1966. Foot rot of black Pepper (*Piper nigrum*) caused by *Phytophthora polmivora* (Butl.) (in Spanish ). Anais. Inst. Mico. 3: 468-491 (RAM 46: 142. 1867)

Ali, S.S. 1987. Effect of three systemic nematicides against root knot nematodes in a cardamom nursery. Nematologic Mediterrania., 15: 155-158.

Altomor. C., W.A. Narvell, T. Bjorkman, and G.E. Harmon, 1999. Solubilization of phosphates and micronutrients by the plant growth promoting and biocontrol fungus *Trichoderma harzianum*. Applied and Environment microbiology, 65(7) : 2926-2933.

Alzarez, M.R., and King D.O. (1969). Peroxidase localization, activity and isozyme patterns in the developing seedlings of Vanda (Orchidaceae). Amer. J. Bot. 56, 180.

Anandaraj, M. and Sarma, Y. R. 1994. Biological control of black pepper disease. Indian cocoa. Areca nut and spices J. 18: 22-23.

Anonymous, (1992) Growth characters of betelvine Ann. Rept. All India Coordinated Research Project on Betelvine 1981 – 92. O.U.A.T., Bhubaneshwar pp.59.

Anonymous, 1982. Annual Report of J.N. Krishi Viswavidyalaya, Jabalpur, for 1982 : 18 p.

Anonymous, 1983. Proceedings of the Forth All India Workshop on Biological Control, 22-24 February 1983, Bangalore, India.

Anonymous, 1985. Percentage of the Fifth All India Workshop on Biological Control, 13-16 September 1985, Coimbatore, India.

Anonymous, 1985. All India Coordinated Research Project on Betelvine. Velur (Tamil Nadu). Ann. Rept.

Anonymous, 1987. AICRP on betelvine. Annual Report, Technical Document No. 6. Bidhan Chandra Krishi Viswavidyalaya (BCKV), Kalyani.

Anonymous, 1988. Microbial Control of *Heliothis armigera*. Technical Bulletin No. 6., Indian Institute of Horticultural Research, Bangalore- 560089.

Anonymous, 1993. Effect of Azospirillum on brinjal and Caleus. In: Proceedings of Scientific Workers Conference in Horticulture, Tamil Nadu Agricultural University. Pp. 112-119

Anonymous, 1997. Pesticides Production – Current status. Pesticides Information 22(4): 36-38.

Anonymous, 1997-98. Annual Report. All India Coordinate Research Project on Betelvine. Indian Institute of Horticulture Research, Bangalore, pp. 82-89.

Anonymous, 1998. National IPM programme in India, Pestology. Annual 1998. p. 413-416.

Anonymous, 1998-99. All India Coordinate Research Project on Betelvine. Indian Institute of Horticulture Research, Bangalore, pp. 91-99.

Anonymous, 1999-2000. Annual Report. All India Coordinate Research Project on Betelvine. National Research Centre for Medicinal and Aromatic Plants, Borivali, Anand. pp. 108-117.

Anonymous, 2000-2001 Annual Report. All India Coordinate Research Project on Betelvine. National Research Centre for Medicinal and Aromatic Plants, Borivali, Anand. pp. 75-86.

Arnon, D. J. Copper enzymes in isolate chloroplasts polyphenol oxidase in *Beta Vulgaris*. Pl. Physiol. 24 : 1-5

Arora, R. Battu, G.S., and Bath, D.S. 1997. Comparative evaluation of some native isolates of a nuclear polyhydrosis virus against *Heliothis armigera* Hubner. Journal of Entomological Research , 21 : 2, 183-186.

Arya, R.S. and Dutta, C.P. 1982. Cultivation of betelvine. Indian Fmg. 31 (12) 15-16.

Ashthana, R.P.. "Disease of *Piper betle* and their control" ,Indian Farming, 8 : 394-397.

Ashthana, R.P.and K.P. Mahmud (1945), "Vegetative propagation of *Piper betle* in the central provinces", Proc. Indian Acad. Sci., 22(B): 70-75.

Asokan, R. 1998. Plant derived polyflavanoids as "screens" for NPVs. Insect Environment. 4 (3) 65-66.

Asokan, R., K. S. Mohan and C. Gopalakrishnan (1996). Effect of commercial formulations of *Bacillus thuringiensis* Berliner on yield of cabbage. Insect Environment, 2 (2) : 58-59.

Ayyar T.V.R. (1963) Pest of Betelvine. Hand book of economic entomology for South India, Madras. 329-331.

Balasubrahmanyam R. , Chaurasia R.S. and Rawat A.K.S. 1984. A foliar analysis survey of betelvine plantations in Mahoba. In : Proc. Natl. Workshop on Betelvine, PP 65.

Balasubramanyan V.R. , Chaurasia R.S. and Singh, K.K. 1989. A foliar analysis survey of betelvine plantations in parts of Uttar Pradesh and Madhya Pradesh J. Plantn. Crops 17 : 90 – 95.

Balasubramani V. and swamiappan, M. 1998, Safety at Ho NPV and SI NPV to greenlace wing crysoiperia carnea stephens- Insect environment 4 (3) 70-71.

Balasubramanyam V.R., Johri, J.K., Rawat A.K.S., Tripathi, R.D., Chaurasia, R.S. 1994. Betelvine (*Piper betle* L.) National Botanical Research Institute, Lucknow.

Balikai, R.A. and sathgi, H.N. 2000. Management of sunflower head  
barer by suing bioinsecticide alone and in combination  
with insecticide. Agriculture science Digest. 20 (2) 110-  
111.

Balsubrahmanyam, V.R.B., Johri, J.K. Rawat, A.K.S., Chaurasia, R.S.  
and Tripathi, R.D., In Betelvine (*piper betle L.*) Economic  
Botany Service, NBRI, Lucknow, 1995.

Barber, C.A. 1903 Pepper disease in the Wynad. Trap. Agric (Colombo)  
22: 206

Barret – Lennard, E.D., Robson, A.D. and Greenway, H. (1982), Effect of  
phosphorus deficiency and water defecation phosphate  
activity from wheat Leaves J. Exp. Bot. 33 : 682-693.

Behera, T.H., Panda, S.K. and Patra, H. K. (1999). `Chromium ion  
induced lipid peroxidation in developing wheat seedlings :  
Role of growth hormone. Indian J. Plant Physiol. 4: 236-  
238

Bhangale, G.T. (2005) Research work carried out at Betelvine Research  
Station, Nimkhedi road, Jalgaon presented in Betelvine  
Group Discussion (26-2-2005) held at MPKV, Rahuri.

Bhattacharjee, Anjan, Bero, P. Das, A.K., Chowdhuri, A., Dasgupta , B. and Sengupta, K. 1989. Studies on carbendazim residues occurring in betelvine leaves. *Pesticides* 23: 41-43.

Bieleski, R.L. and Ferguson, I. B. (1983). Physiology and Metabolism of Phosphate Compounds (Eds. A. Lauchi and R.L. Bieleski) In L Inorganic Plant Nutrition. Encyclopedia of Plant Physiology, New Series, 15. pp. 422-449. Springer – Verlog New York.

Bielesko , R.L. and Ferguson, I.B.(1983). Physiology and metabolism of phosphate compounds (Eds. A. Louchi and R.L. Bieleski) In: morganc plant Nutrition Encyclopedia of plant physiology, New series, 15. pp. 422. Sringer-veriag New York.

Birecka H. and Garraway, M.O. (1975). Corn leaf isoperoxidase reaction to mechanical injury and infection with *Helminthosporium maydis*. *Plant physiol.* 61 : 561-566.

Bis Wol G. and A Narain (1992), Toxic effect of culture filtrates of *Aspergillus flavus* on seed germination and seedling growth of jute, ragi and mustard plants, orissa J. Agric. Res. 5 (1-2) : 136-139.

- Biswal G. and A. Narain (1992). Toxic effect of culture filtrates of *Aspergillus flavus* on seed germination and seedling growth of jute, rogi and mustard plants. Orissa Agric. Res. 5 (1-2) :136-139.
- Biswas, T.D. 1971. Cumulative effect of different levels of manures on the physiological properties of soil. J.
- Black W. (1960), "Races of *phytophthora infestans* and resistance problem in potatoes", scott. Pl. Breeding sta. Rep., pp. 29-38.
- Black, W. (1960), "Races of *Phytophthora infestans* and resistance problem in paratoes", Scott. Pl. Breeding sra. Rep., pp. 29-38.
- Black, W., C. Mastenbroek, W.R. Mills and L.C. Pearson(1953), "A proposal for an international nomenclature or races of *Phytophthora infestans* and of genes controlling immunity of *solanum demissum* derivatives", Euphytica, 2:173-179.
- Black, W.C. Mastenbroek, W.R. Mills and L.C. Pearson (1953), "A proposal for an international nomenclature of races of *phytophthora infestans* and of genes controlling immunity if *solanum demissum* derivatives", Euphytcs, 2:173-179.
- Bohert, H.J., Nelson, D.E. and Jensen, R.G. (1995). Adoptations to environmental stresses. Plant Cell 7 : 1099-1111.
- Bose T.K., Mukhopadhyay, T.P. and Basu, T.K. (1984). A note on the effect of ascorbic acid and IBA on rooba in cuttings. Hort. 54: 207.

Boskaron, R.K.M. Venugopal, M.S. and Mahadevan, N.R. (1997 Publ. 1998). Optical brightners as UV protections and their influence on the virulence of NPV of *S. litura*. Journal of Biological Control II (1-2) 17-22.

Boskaron, R.K.M. Venugopal, M.S. and Mahadevan, N.R. 1996. Effect of host plants on the infectivity and yield of nuclear polyhydrosis virus on spodoptera litura. Journal of Biological Control. 10 : 1-2, 73-78.

Boskaron, R.K.M. Venugopal, M.S. and Mahadevan, N.R. 1999. Effect of certain botanical on biological activity of NPV of tobacco caterpillar (*S. litura*). Indian Journal of Agricultural Sciences. 69 (3) 224 – 226.

Bould, C. 1955. J. Hort. Sci. 30:183, as quoted from Technical Bulletin No. 15. Determination of the nitrogen requirements of crops by analysis. pp. 89-96.

Brady C.N. (1996). The nature and properties of soils. Prentice – Hall of India Private Limited, New Delhi, pp. 620.

Brasier, C.M. 1969 a. The effect of light and temperature on reproduction in vitro in two tropical species of *Phytophthora*. Trans. Brit. Mycol. Soc. 52: 105-133.

Brasier, C.M. 1969 b. Formation of oospores in vivo by *Phytophthora palmivara*. Trans. Brit. Mycol. Soc. 52: 273-279.

Bray, H.G. and Thorpe, W.V. 1954. Analysis of phenolic compounds of interest in metabolism Meth. Biochem. Anal. 1:27-52.

Brefman, A. 1940. Cultivation and trade of Pepper ( *P. nigrum* ) on the island of Bangka Meded. Van der Dienst V.D. Landb., NO. 16.

Brian, F.W. and Hemming, H. G. 1950. Some nutritional conditions affecting spare production by *Trichoderma viride* pers. Ex. Fries. Trans. Brit. Mycol. Soc. 33: 132-141

Briton – Jones, H.R. 1940. The disease of the coconut palm Baillere, Tindal & Co., London.

Broadbent P. and Baker K.F., 1974. Behaviour of *Phytophthora Cinnamomi* in soils suppressive and conducive to root. Aust. J. Agric. Res. 25: 121-137.

Brundrett MC, Kendrick B, Peterson CA, 1991. Efficient lipid staining in plant material with Sudan Red 73 or Fluorol Yellow 088 in polythene glycol glycerol. Biotechnic and Histochemistry 66, 111-116.

Burkhill, I.H. (1964) A dictionary of the economic products of the Malayan peninsula, Vol. II Min. of Agric, Malayasia : 1762 – 72.

Butler, E. J. 1906. The wilt disease of pigeon pea and pepper. Agric J. India. 1:25.

Butler, E. J. 1918. Fungi and Disease in plants. 598 pp. Thacker and Spink, Calcutta.

Chang, Y. C., Y. Baker, R., Kleifeid, O. and Chet, I. 1986. Increased growth of plant induced by the biological control agent, *Trichoderma harzianum*, Plant Dis. 70: 145-148.

Chattopadhaya, N.C., Das, A. K., Mukherjees, S. Das, S. and Mishra, A.K. 1961. Phytochemistry of betel leaves and some aspects of storage. Proc. Gr. Discuss. NBRI. Feb. 1981. pp.139-145.

Chauhan, L. S. and Jyoti Singh(1983): Thread blight and rhizome rot of Ginger and its control; pesticides 17 (8): 24-25.

Chaurasia, R. K. 1993. In : Advances in Betelvine Production Technology (Eds. R. K. Chaurasia, M.L. Nayak and M. N. Khare), 56-57.

Chaurasia, J. P., Vyas, K.M. 1997. In vivo evaluation of some homeopathic drugs against betelvine *Phytophthora* disease. Indian Phytopathology 50 (4) : 542-547. Curr. Res. Univer. Of Agr. Sci.Bangalore 19 (10) 174-175.

Chee, K.H. 1969 Hosts of *Phytophthora palmivaro*, Rev. Appl. Mycol. 48: 337-344.

Chee, K.H. 1973. Production, germination and survival of chlamydospores of *Phytophthora* from *Hevea brasiliensis*. Trans. Brit. Mycol. Soc. 61: 21-26.

Chelliah, S. and K. Srinivasan (1986). Bioecology and Management of diamond back moth in India. Proc. J. Int. Workshop on Diamond back moth Management. Pp. 63-76. Eds. Talekar, N.S. and Gr: 999, T.D. The Asian Vegetable Research and Development Centre, Shanhua, Taiwan.

Chopra, R.N., Nayar, S.L. and Chopra, I.C. 1956. Glossary of Indian Medical Plants *Piper betle* Linn. CSIR, New Delhi, pp. 194.

Chouasia, S.C. (1976), "Studies o Foot Root and Leaf Rot Disease of pan (*Piper betle* L) with special reference to pathogenesis and Control Measures". Ph. D. Thesis, Univ. of Sagar.

Choudhari, H.C. (1954), "spray tests for control of potato blight in the hills of West Bengal", Am. Potato J., 31: 263-272.

Choudhari, H.C. (1959), "spray tests for control of potato blight in the hills of West Bengal", Europ. Potato J., 2: 89-99.

Chowdhary, S. (1944), "Disease of pan (*Piper betle* L.) Sylhet Assam – I, II and III : Effect of manuring on the incidence of *Phytophthora* foot rot and leaf rot disease", proc. Indian. A Cad.

Christie, J.R. 1957. The Yellows disease of Pepper and spreading decline of citrus. Plant. Dis. Repr. 41: 267-268.

Clark, R.B. (1984),Physiological aspects of calcium ,magnesium and molybdenum deficiencies in Plants. Agronomy Monograph (Second Eds.),Madison,U.S.A. 992-996.

Clifford, C., Yang, Y. and Chung, F., In Current Drug Metabolism, Bentham Science Publishers, 2002, pp. 233-255.

Cobb, N.A. (1916). Estimating the nema population of Soil Agric. Tech. Grc. Bur. Pl. Ind US. Dep. Agric. 1: 49.pp.

Crowdy, S.H. 1977. Translocation. Pp. 93-114, In Systemic Fungicides ed. Marsh, R. W. Longman Group Ltd., London.

D'Souza, A. Roy, J.K.,Mahonty, B. and asgupta, B.2001. Screening of *Trichoderma harzianum* against major fungal pathogens of betelvine. Indian Phytopathal. 54:340-345.

Dake, G.N. and S. Edison (1989) : Association of Pathogens with rhizomes rot of Ginger in Kerala. Indian phytopathology. 42 (I) : 116-119.

Danielson, R.M. and Dovey, C.B. 1973, Non nutritional factors affecting the growth of *Trichoderma* in Culture. Soil Biol. Biochem., 5:495-504.

Dastur,J.F. (1935),"Disease of pan in central provinces", proc. Indian.ACod.Sci,1(B):778-815.

De Bruyn,H.L.G.(1926),"The saprophytic life of *Phytophthora* in soil." Mededeeling von de Londfonwhooge school en van de Dareen Verbonden Instituten,24 (paper 4):1-38.

DeJong, D.W. (1967). An investigation of the role of Plant peroxidase in cell wall development by the histachemical method. J. Histachem. Cytochem. 15,335.

Desai, B.G.and V.M. Pathak (1969),"some studies on soft rot disease of Luffa cylindrical Roem",Phytopath.Mediterranea,8:238-239.

Deshmukh, M.J. and K.N. Chhibber (1960), "Field resistance to blight (*Phytophthora Colacasiae* Roc.) in colocasia antiquorum Schott", Curr.Sci.,29: 320-321.

Deshmukh,N..S. 1978. Economics of production and marketing betelvines  
in Amravati district. M.Sc. Thesis unpub.,PKV.Akola.

Dff, M.G.S., Sarath, G. and Plaxton, W.C. 91994) . The role of acid  
Phosphatase in Plant Phosphorus metabolism, Physio. Plant.  
90:791-800.

Dhande, G.W. and sulaiman, M. (1961) occurrence of root-knot nematodes  
on betelvine in Maharashtra, Curr Sci. 30 : 351-352.

Dhandopani, N.,Kalyanasundram, M.,Swamiappan, M.,Sundara Babu,  
P.C.Jayaraj, S.1992. Experiment or management of major  
pests of cotton with biocontrol agents in India. Journal of  
Applied Entomology. 114:1, 52-56.

Duff M.G.S., Sarath G. and plaxton W.C. (1994) The role of acid  
phosphatase in plant phosphorus metabolism, physio plant.90  
:791-800.

Dutt,B.L.(1962).,"Late blight of potato in the hills and its control",Indian  
potato J.,2: 27-33.

Dutta, P. and Das, B.C.1997. Biomass production of *Trichoderma*

*harzianum* using various common wastes, New Agriculturist, 8  
(1) : 85-88.

El – Flouly, M.M. and Jung, J. (1970), changes in enzyme activity in wheat seedlings grown under soil stress conditions. Plant Physiol. 46 (suppl.): 60.

El-Flouly, M.M. and Jung, J. (1970) changes in enzyme activity in wheat seedlings grown under soil stress conditions, Plant Physiol. 46 (suppl.) : 60.

Elhance, D.N., 1984. Fundamentals of Statistics, Kitabmahal. 15. Thornhill Road, Allahabad, India 1000 PP.

El-keblawy, A and Al – Rawai, A. (2005). Effects of salinity temperature and light germination behavior of the invasive *Prosopis juliflora*. J. Arid. Environ. 61:555-565.

El-Sharkawi, H.M., Forghal; K.A. and Sayed, S.A. (1977). Trifactorial interactive effects of nutrients, water potential and temperature on carbohydrate allocation to the embryonic axis of desert plant seeds Journal of Arid Environments 35:655-664.

Epstein, E. (1969). Mineral metabolism. In "Plant Biochemistry" Banner S.J. and Varner J.E. (Eds.) Academic press, London and Orlando 438-466.

Epstein, E. and Robsinson, R.R.1965. A rapid method for determining leaf area in potato plants. Agron. J. 57 (5): 115-116.

ESpelie KE, Franceschi VR,Kolattukudy PE.1986. Innmunocyto chemical localization and time course of appearance of an anionic peroxidase associated with suberization in wound-healing potato tuber tissue. Plant physiology 81\_487-492.

Espelie KE,Kalativkudy PE.1985. purification & characterization of an abscisic acid – inducible anionic peroxidase associated with suberization in potato (*Solanum tuberosum*). Archives of Biochemistry and Biophysics 240, 539-545.

Evans, G.C.1972. The qualitative Analysis of Plant growth. Blackwell Science Publication, Oxford.

Evans, H.J. and Sorger, G.L. (1966). Role of mineral elements with emphasis on univalent cations, Ann. Rev.Plant. Physial. 17:47-76.

Fahmad, R., Yadava, C.P. and Lal, S.S. 1999. Efficacy of NPV for the management of *H. Armigera* (Hubner) infesting chickpea. Indian Journal of Pulses Research 12 (I) 92-96.

Fair, P., Tew, J. and Creswell, C.F. (1973). Enzyme activities associated with CO<sub>2</sub> exchange in illuminated leaves of *Hordeum vulgare* L. Effect of light period, leaf age and position on CO<sub>2</sub> compensation point. Ann. Bot. 37: 831-844.

Farkode, B.C. Ahire, R.D. Patonge, N.R. and Ahire P.S. 1999. Extent of adaption of biological pest control in cotton. Journal of soils and crops g (i) 108-110.

Fernando, L.C.P., Wickramananda, I.R. and Aratchigo, N.S. (2000). Studies on coconut Mite Acerio Guerreronis in Sri Lanka. In: International Workshop on coconut mite Aceria Guerreronis Coconut Research Institute, Lunwilo, Sri Lanka.

Fielding, J.L. AND HALL, J.L. (1978). A biochemical and cytochemical study of peroxide activity in roots of *Pisum sativum*. J. Exptl. Bot. 29, 469.

Fincham, J.R. and P.R. Day (1969), Fungal Genetics, (Second edition), Blackwell, Oxford, PP. 326.

Finney, D.J.1971. Probit Analysis 3<sup>rd</sup> ed.Combridge Univ.Press,  
London,PP.318

Ganguly, P. and Gupta, D.1973. Identity of betel leaf (*Piper betle* Linn.) in  
West Bengal market – Part-I. Indian Agric.,17:251-55.

Gangwar, Santhos k. and Dasgupta, B.1989. Rizolex 50 WP – a new  
fungicide to control the foot rot of betelvine (*Piper betle* L.)  
caused by S.Colfsii. Proc. Indian Acad. Sci. (Plant Sci.) :  
99:265-269.

Gitanjali Jayachandran.Choudhari, S. and Ramakrishna. N.1999. Age  
related responses of sponloptera litura (Fabricious) to NPV.  
Journal of Entomological Research 23(3) 247-260.

Gogoi, N.K., A.K. Phookan, and B.D. Narzary,2004. Effect of *Trichoderma harzianum* and fungicides on plant growth and intensity of collar rot of elephant foot Yam (*Amarphophallus paeoniifolius*). Journal of Root Crops, 30 (I) : 76-79.

Gomoa, A.O. and F.G. Mohamed,2000 Effect of some biocontrol agents  
and agricultural chemicals comparing with Rizolex T50 in  
controlling the Sclerotium disease and productivity of Dahlia  
plant. Annals of Ggricultral Science Cairo Special,1725-  
1748.

Greenberge, J.,Gorgen R.,and Riov,J. (1975). The role of cellulase and polygalacturonase in abscission of young and mature Shamouti Orange fruits. *Physiol.Plant.* 34%.

Grinstein, A.,Katan, J.,Abdul Rogak, A.,Zeidon, O and Elod,Y. (1979). Control of *S. rolfsii* and weeds in peanuts by solar heating of soil *Plant Dis. Repr.* 63:1056-1059.

Guha, P.K.,Halder, P. Dasgupta, B.,Sengupta, K. and Bhattacharya A. 1990. Studies on the dissipation of Corbendazim residue in/On betelvine leaves. *Pestology* 24:23-26.

Gunnasekaran, N.,Shanmugavelue, K.G. and Rajeskaran, K.L.1983. Effect of triacontanol on nutrient uptake in tomato. *Proc.Natl.sem.prod.Tech.Tomato and chilli.* 72-74.

Hadar, Y.; Chet, I. and Henis, Y.1979. Biological control of *Rhizoctonia solani* damping off with wheat bran culture *Trichoderma harzanium*. *Phytopathology*, 69:60-68.

Hall, R.A. HHussey, N.W. and Mariau, D. (1980). Results of a survey of biological control agents of the coconut mite *Eriophyes Guerreranis*. *Oleagineux*. 35, 395-398.

Hawker, J.S. and Jenner, C.F. 91993). High temperature affects the activity of enzymes in the committed pathway of starch synthesis in developing wheat endosperm Aust. J. Plant physiol. 20:197-209.

Hazarika, L.K. and puzari, K.C. 2000. Biopesticides for the management of tea pests. PP. 36-43. In : Entrepreneurs workshop on Ecofriendly Technologies of Bio-pesticides and Bio-fertilizers for crop management, conducted by Department of Biotechnology, Ministry of Science & Technology, Government of India, New Delhi – 110 003.

Health, R.L. and packer, L. (1968). Photoperoxidation isolated chloroplast. I. kinetics and stoichiometry of fatty acid peroxidation. Arch. Biochem. Biophys. 124 : 189-198,

Heigaord J. Peterson JF, veitch NC, Pedersen BJ , Welinder KG.1991 Three cationic peroxidases of barley grain. Isolation and preliminary characterization. In : Labarzewski J, Greppin H, penel C, Gaspar Th,eds. Biochemical, Molecular and physiological aspects of Plant Peroxidases. University of Geneva, 49-53.

Hendrix, F.F. Jr.,Powell and J.H.Owen (1966). Relation of root necrosis caused by *Pythium species* to peach tree decline,"Phytopath; 56:1229-1232.

Hendrix, F.F.Jr. and W.A."Campbell (1970), Distribution of *Phytophthora* and *Pythium Species* in Soil in the Continental United States", Con.J. of Bot.,48:377-384.

Hickmon, C.J. (1958),"phytophthora, plant destroyer" Trans. Brit. Mycol. Soc.,41: 1-13.

Hofstee, B.H.G.(1959).Fatty acid esterases of low serine sensitivity and related enzymes. In : P.D.H. Boyer (ed.), Enzymes PP. 485-500.

Horoon Usmani, S.M. and Gaffor, A (1983). Relative efficacy of poly – ethylene mulching in reducing viability of scleratia of *S. oryzae* in soil. Ecology and Management of soil borne pathogens. American Phytopathological society, St.Paul, Minnesota, U.S.A. PP. 285-288.

Hoslop, E.C. and Stahmann, M.A. (1971) peroxidase and ethylene production by barley leaves infected with *Erysiphe Graminis* of Sp. *Hordei*. Physiol. Pl. Path. 1.-297.

Hrnid, Y. and Popavizas G.C. (1983). Factors affecting germinability and susceptibility to attack of *S.rollsii* by *T.harianum* in field Soil. Phytopathology 73: 1469-1479.

Imberty A, Goldberg R.Catesson A.M.1985. Isolation and characterization of *Populus* isoperoxidases involved in the last step of lignin formation.*Planta* 164,221-226.

Iyengar, N.V.R., Srivastava, H.C..Date, W.B. and Mathur, P.B.1956. Studies on pocking and cold storage of betel leaves (*Piper betle* L.) *F.Ull.cent.Fd.Technol.Res.Inst.5*:307-310.

aanthi, P.D.K. and Padmovothatamma, K.1996. Cross infectivity and safe of NPU, *B.thuringiensis* sub sp. *Kurlaki Berliner* and *Beawceria hassiana* (Balsumo) Vuille to pest of Groundnut (*Arachis hypogaea* Linn) and their natural enemies. *Journal of Entomological Research* 20 (3).211-215.

Jackson, M.L.11973. Soil chemical Analysis, Peentice Hall of Plant India (Pvt.) Ltd., New Delhi.

Jackson, M.L.1973. So.I chemical Analysis, prentice Hall of plant India (Pvt.) Ltd.,New Delhi.

Jacob, A. and subromanian, T.R.1972. Nuclear polhedrosis on some Lepidoptera. *Curr Sic.*,41:536.

Jagdale, G.B. Pawar, A.B. and Darekar, K.S. 19850. Effect of organic amendments on root knot nematode infesting betelvine. Int. Nematol. Network News I.,1:4-5.

Jagdale, G.B., Pawar, A.B., Madhane, N.S. and Narkhae, T.N.1984. Managing betelvine insect pests. Farmers J. 4 :50-511.

Jagdale, G.B.,Pawar, A.B.and Darekar,K.S.1985, Studies on control of root knot nematode on betelvine. Ind.Nematol.Network Newsstr.,2:10-13.

Jagdale, G.B.Pawar, A.B.and Darekar,K.S.1984, control of root knot nematode on betelvine with systemic nemoticides. Int. Nematol. Network newsl.,1:4-5.

Jagdale,G.G.,Pawar,A.B. and Darekar, K.S.1985 b.Pathogenicity of *Meloidogyne incognita* an betelvine.Indian J.Nematol.,15:244

Jamal,A.(1982).Observations on the interaction of *Sphaerotheca fuliginea* and *M.incagnita* on *Lagenaria Leucanltia*. Comparative Phsiology and Ecology, 4:86-88.

James A.Spencer and H.J. Walters (1962). Stud of variation of *Copnespara cassiicola*. Phtopatholog 52:28.

Janathan, E.I.; Gajandran, G.; Arulmozhiyan, R. 1997. Interaction of *Rotylenchulus reniformis* and *Phytophthora palmivora* on betelvine, *Nematologia Mediterranea* 25:191-193.

Jeeva, S. 1987. Studies on the effect of *Azospirillum* on the growth and development of banana Cv. Poovan (AAB). M.Sc. (Hort) Thesis. Tamil Nadu Agricultural University, Coimbatore.

Jyothi, L.J., MANI, A.K. and PAPPAIAH, C.M. 1993. Influence of NPK and Azospirillum on the yield of Cabbage. S. Indian Hort. 41 (5) : 270-272.

Jeng, T.L. and Sung, J.M. (1994). Hydration effects on peroxidation and  $\text{H}_2\text{O}_2$  – scavenging enzyme activity of artificially aged Peanut. Seed SSC. Technal. 22:531-539.

Jensen, W.A. (1955). The histochemical localization of peroxidase in roots and its induction by indole acetic acid. Plant. Physiol. 30, 426.

Jeppson, L.R., Keifer, H.H. and Baker, E.W. (1975). Mites Injurious to Economic plants. University of California Press, Berkeley, California, USA. PP. 614.

Johan, A. 1998 use of fertilizers to increase the yield of Pann (*Piper betle* L.). Hamdard Medium 41:103-106.

Johnson, A.M. and Campbell Q.HG.1980. Managing nematode population densities of tomato using crop rotation and nematicide J. Nematol. 12:6-19.

Johri, J.K.,Santho Devi,1998. Ultra-structural studies on *Phytophthora palmivora* infection on betelvine (*Piper betle* L). Archieves of Phytopathology and plant protection 31 (3) : 233 – 240.

Kadam, N.D., Patil, G.D.,Cougule, B.A. and Kadam, S.S. 1988. Effect of folior application of vipul on yield, soluble protein, fatal protein and nitrate reductase activity in spinach. Indian J. Plant physiol. 31 (I):123-125.

Kalia, V.Choudhari, S. and Gujor, G.T.2001. Optimization of production of Nucleapalhedrosis of *H.armigera* through larval stages. Phytoparasitica 29 (I) 23-28.

Kar, S.N., S.K. Nog, S, Pal, B. dasgupta and Aditya Choudhary, N.(1996). Persistence of copper fungicides on betelvine (*Piper betel* L.) leaves & soil following application of Bordeaux mixture, pesticide Res. Journal 8 (I) : 71-74.

Kathriesan, K., Ravishankar, G.A. and Venkataraman, L.V. 1990, Auxin phenol induced rooting in a mangrove *Rhizophora apiculata* Blume. Curr.Sci. 59:8,43-432.

Kaung,Z.(1965),"persistence and movementof *Phytophthora infestans* in soil", Trns. Brit. Mycol. Soc., 39:385.

Khan, M.W., Khan, A.M. and sexena, S.K.1974. Rhizospheric fungi and nematodes of egg plant as influnced by oil-case amendments. Indian phytopathol.,27:480-484.

Khan, M.W., Khan.A.M. and saxena, S.K.1973. Influence of certain oil-cake amendments on nematodes and fungi in tomato field Acto Botanica India. 1:49-54.

Khanna, R.N. and Vishwadhar (1973),"A new race of *Phytophthora infestans* from khasi Hills",JIPA, 2:46-47.

Khanna, R.N., V.K.Bahal and Viiishwadhar (1977),"Identification of some high specirum races of *Phytophthora infestans* in khasi Hills",JIPA,4 (I) : 18-21.

Khatri, R.K., Keshwal, R.I., Nayak, M.L. 1997. Biochemical changes induced by *Xanthomonas campestris* pv. *Betlicola* causing leaf spot betelvine, plant Disease Research 12:25-27.

Khoshoo, T.N. 1984. Keynote address. Proceedings of the National Workshop on Betel vine 1982, pp.1-12.S.K.Rayet al. (Eds). The Agricultural society of India, Calcutta.

Knauss, J.F. 1978. CGA 48988, a new systemic fungicide for control of *Phytophthora nicotinae*. Vax. Parasitica on *peperamia obtusifolia*. Proc. Fla. State Hortic. Soc. 90:330-332.

Kousalya, G. and Jeyarajan, R. 1988. Techniques for mass multiplication of *Trichoderma spp.* Pers Fr. and *T. harianum* Rifoi. Abstract of papers presented in National Seminar on management of crop Disease with plant products/Biological Agents, 10-12 January, 1988, Agricultural College and Research Institute, Madhurai. PP.32-33 Kousalya, G. and Jeyoraman, r. 1990. Mass multiplication of *Trichoderma spp.* J. Biol. Control, 4 (I):70-71.

Krishankumar, N.K., K.Srinivasan, P.R. Ramachandar and C.L. Suman (1986). Optimum control strategy of cabbage pests from a

chemical control trial. Prog. Hort., 18:104-110.

Kulkarni, G.G. and Hugar, P.S.1999. Effect of host plants on POBS yield and mortality of *Spodoptera litura* larvae infected with NPV. Karnataka Journal of Agriculture Science 12 (1-4) 60-63.

Kulkarni, G.G. and Hugar, P.S.2000 standardization of inoculum dose for mass multiplication of spodoptera litura nuclear polyhedrosis virus (SI NPV) on different host plants. Karnataka Journal of Agricultural Sciences 13 (I) 51-54.

Kulkarni, G.G.and Hugar, P.S.1999. Persistence of NPV on different host plants of S.1:tura Haryana Agricultural University Journal of Research 29 (3-4) 95-99.

Kulkarni, G.S. (1913),"Observations on downy milde Sclerospora graminicola (Sacc).Schroet.of bajara and jawar" Mem.Deptt.Agr. Bot. Ser., 5:268-274.

Kumar, N.S.1999. Impact of NPV and Azadiractia : an the digestive enzyme activityand biochemical composition of the gut of *Harmigera* (Hubner) Tropical Agricultural Research II, 393-407.

La Berge DE, Kruger JE.1976. Changes in barley Peroxides activity during kernel development cereal Chemistry 53, 762-767.

La Berge DE.1975. Anatomical distribution of Peroxidase isozymes in barley Kernels. Canadian Journal of Plant Science 55, 661-666.

Lagrimini LM, Joly RJ, Dunlop JR, Liv TTY.1997. The consegence of Prroxidase over expression in transgenic plants on roots growth and development plant molecular Biology.33, 887-895

Lakshmanan P.L. 1985 Control of *Helicotylenchus dihystera* on tomato by organic amendments. Proc. IV. Nematology symposium. Mohanlal Sukhadia Univ.,Udiopur.

Lashin, S.M., H.S. Elnasr; M.A. Elnogar nd M.A. Nofal (1989), Biological control of *Aspergillus niger*, the causal organism of peanut rot by *Trichaderma harzianum*.Annals of Agricultural Sciences, 34 (2): 795-803.

Leather R, R.I.1967. The occurrence of a *Phytophthora* root and leaf disease of black piper in Jamaica. F.A.O.PI. Prof. Bull. 15:15-16.

Lekha, V.S. Bidappa, C.C. and kamalakshi Amma, P.G. (1999). Organic manures and their importance in coconut cultivation. Indian coconut J.30 (3), 7-8.

Leonion, L.H. (1934), "Identification of *Phytophthora* sp." (Agri. Exp. St. Coll. Of Agr.) West Virginia Univ. Bull.,262.

Li,C.W.(1981). The origin, evolution, taxonomy and hybridization of chines cabbage. In Chinese Cabbage: proc. First Int. Symp. Ed. N.S. Talekar, T.D. Griggs, PP.3-11. Shanhua, Taiwan : Asian vegetable Research and Development Center.

Liszkay S, Kenk B, Schopfer P.2003. Evidence for the involvement of cell wall peroxidase in the generation of hydroxyl radicals mediating extension growth planto 217, 658-667.

Madane, N.P. and Naket R.V. (1992). All India co-ordinated Research Project on Betelvine Project on Betelvine Progress Report 1992-93.

Maehly, A.C. (1955). Plant peroxidase. In : P.S. Cdowick and N.O. Koplan (eds.), Methods in Enzymology, Vol. II, pp.271-278. Academi cress, New York.

Mahanty B., Roy, J.K., Dasgupta, B. and Sen. C.2000. Relative efficacy of promising fungicides and biocontrol agent *Trichoderma* in the management of foot rot of betelvine J. plant. Crops.,28 (3) : 179-184.

Mahmud, K.A. (1940)," Report of pan cultivation Scheme in central provinces for 1940-41," Govt. printing press, central provinces and Berar, Nagpur, p.55.

Maiti, S. and Biswas, S.R.,1991. Sexual dimorphism in betelvine (*Piper betle* L.) Indian J.Agric. Sci. 61:656-58.

Maiti S. and Chaudhary, S. 1995 Effect of some systemic fungicides on sclerotial germination and growth of *Sclerotium rolfsii* in vitro. Z.Pflanzenschutz. 82:233-235.

Maiti, S. and Raghavendra Rao, N.N.1992, Cultivation and crop improvement of piper betle in India. In Recent Advance in Medicinal, Aromatic and Spice Crops (Raychaudhuri, S.P.Ed.).vol.2pp.409-12

Maiti, S., Khatua, D.C. and Sen., C. 1978. Chemical Control of two major foliage diseases of beteline. Pesticides 12:45-47.

Maitil, S. 1977. Disease of *Piper betle* (L) with emphasis on the foot rot caused by *Sclerotium rolfsii* Sacc. And its Control Ph.D. Thesis, B.C.Krishi Viswavidyalaa, W.Bengal, India 181 PP.

Maiti, S. Biswas, S.R. and Raghavendra Rao, N.N. 1992. Divergence in Sexual dimorphism among betelvine (*Piper betle*) clones. Indian J. Agric Sci., 62:780-782.

Maiti, S. and Biswas, S.R. 1991 Indian J. Agric. Set 61:656-658. Aaaitil, S. Biswas, S.R. and Rao N.N.R. 1992 Indian J. Agric Sci. 62:780-782.

Mammer, K.V. (1975). On a wilt disease of betelvine in Kerala caused by root – knot nematodes. Agric. Res. J. Kerala 12:76.

Mandal, B.K., Choudhary, S.K. Sengupta A, K. and Dasgupta, B. 1994. Response of betelvine (*Piper betle*) varieties to organic manure and prilled urea. India J. agric. Sci. 64 (5) : 297-301.

Marfmuthu, T., Rabindran, R. and Purushothaman, D. 1986. Association of Azospirillum spp. In the roots of betelvine (*Piper betle* L.). In : Prac. Natl. Sem. Microb. Ecol., Jan.23-24. Tamil Nadu Agricultural University, Coimbatore.

Marmiuthu, T. 1985, Past, Present and future status of *Phytophthora* wilt of betelvine in Tamil Nadu Worksho, AICORP on Betelvine, Bapatla.

Mathews, V.D. (1931) . Studies on genus *pythium*, Univ. of North Carolina Press, Chope Hill, pp.136.

Mathur Eneh; B.B.Lal Thakare and R.B. Singh (1984): Effect of different fungicides on Ginger rhizome rot pathogen and their effect on germination and rotting of rhizomes. Indian Journal of Mycology and plant pathology. 14 (2) : 155-157

Mayer, A.M., and Harel, E. (1979). Polyphenal oxidases in Plants. Phytochemistry 18, 193.

Mc Donough, E.S. (1938), "Host parasite relationship of *Sclerospera graminicola* on species of Setaria," phytopath; 28:846-852.

Mehetre, S.T. and Solankhe, G.N.1998. Cinoarature efficacy of Trichogrammas preteosum Riley and Ha NPV with endosultan in Controlling *H.Ormigera* (Hubner) Infesting tomato. Journal of Maharashtra Agricultural Universities 23 (2) 175-176.

Mehrotra R.S. (1961). "Studies on soil fungi from *Piper betle* orchards with special reference to the disease caused by *Phytophthora parasitica* var. *piperina* and their control." Ph.D. Thesis Univ. of Sagar.

Mehrotra, R.S. and D.P. Tiwari, (1976). Organic amendments and control of foot rot of *Piper betel* caused by Ann. Microbiol. (Institute of pasture) 127 A : 15-21.

Mishra, B. (1992). Studies on major disease problems of Betelvine in Orissa, Indian J. Mycol and Pl. Pathology 16(2) : 200-202.

Mishra, R.P. and Chand, J.N. 1970. Efficacy of different fungicides against foot and root rots caused by *Sclerotium rolfsii*. PANs. 16: 327-330.

Mishra, S.D. and GAUR, B.K. 1977. Role of petiole in protein metabolism of senescing betel (*Piper betle* L.) leaves, plant physiol. 57:961-964.

Mishra, S.D. and Prasad. S.K. 1974 Effect of soil amendments on nematodes and crop yield : 1. Oil-seed cakes organic matter, inorganic fertilizers and growth regulators on nematodes associated with wheat and their residual effect on moong. Indian J. Nematol., 4:1-19.

Mishra, S.D. and Gaur. B.K. 1972. Control of senescence in betel leaves by depholiation Ex.p. Geront. 7:31-35.

Misra, M.P. Pawar, A.D., Ram, N. 1991, use of Nev in Management of the inset pests, *Helianthis ormigera* (Hubner) in gram, Journal of the Andoman Science Asso.

Mithila, J., K.S. Shivashankara and Maiti S. 1999. Growth and morphology of vegetative and reproductive branches in betelvine (*Piper betle* L.). Journol of plantation crops (Accepted).

Mohanty N.N. and P.K. Mohapatra (1968). A new leaf spot disease of betelvine caused by *Corynespora Cassiicola*. Indian Phytopath. 21:275=80.

Mohanty N.N. Narain, A, Dash, S.C. Nayak, B.K. Mahapatra, P.K. and Mohanty, S.R. 91981). Betelvine disease in Orissa. Improvement of betelvine cultivation proceedings of group discussion. February 27-28, 1981, NBRI, Lucknow, pp – 30-32.

Mohanty, B.J.K. Roy and C. Sen. (2000). Two major fungal diseases of betelvine. In : Diseases of Plantation Crops, Spices, betelvine and mulberry (Ed.,M.K. Dasgupta) pp. 133-137.

Mohanty N.N. and P>K> Mohapatra (1968). A new leaf spot disease of betelvine caused by *Corynespora cassiicola*. Indian phytopath. 21:275-280.

Mote U.N.1977. Incidence and chemical control of Chilli mites, phytophagotorsonemus latus Banks J. Maharashtra Agric. Univ. 2 (I) : 73-74.

Mukhopadhyay, A.N. 1987. Biological control of soil borne plant pathogens by *Trichoderma* spp. Indian J. Mycol. Pl. Pathol.,17:1-10.

Muller, A.M. and Comphell Q.H. 1980. Managing amendments in nematode control on examination of the literature, *Nematropica*, 12:319-326.

Naik, M.K. Hiremath, P.C. and Hiremath, S.V. 1988. Post inflectional changes in the betel vine lenres infected with *Colletotrichum gloeosporioides*. Indian Phytopath. 41 (3) : 370-372.

Nakana, Y. and Asoda, K. (1981). Hydrogen peroxide is scavenged by ascorbate – Sepcific peroxidase in spinach cloroplasts, plant cell physiol. 22:867-880.

Nambiar, K.K. N. and Sarma, V.R. 1976. Quick wilt (foot rot) disease of Pepper( *Piper nigrum* L.). Arecanut & spices Bull. 7 : 89-91.

Nambiar, E.P., Nair, T.J., and MoneyY, N.S. 1965. Preliminary Studies on the incidence of wilt disease of Pepper and its relationship to the nitrogen and base status of the soil. Indian J. Agric. Sci. 35 : 276-281.

Narasimhan, V.1964. Some studies on wilt disease of betel vine ( *Piper betle* L.) M.Sc. (Ag.) Thesis 67 pp. Madras University, Madras.

Narasiamhn, V., Venkata Rao, A., Subramanian, K.S, and Vydy -  
Sekharan, P.1976. Fungicidal control of betel vine wilt.  
Pesticides 10 (4) : 34-35.

Narayan, K.1980. Effect of NPV of *Heliothis armigera* (Hubn.) and  
*Spodoptera litura* (F.) on parasites, predators and Silk-works.  
Paper presented at 3<sup>rd</sup> Annual Workshop on 'Biological  
Control of crop pests and weeds; at ludhiana, Punjab, Oct.27  
to 30<sup>th</sup> ,1980.

Narayanan, K. Subramanian, T.R. and Jayaraj. 1986. Effect of nucleo  
polyhedrasis virus on carbohydrate, glycogen and fat  
Contents of *Heliothis ormigera* (Hubner) (Noctuidae :  
Lepidoptera) National Sympasium on Insect Physiology,  
ecology and behaviour. Trivendrum. Abstract No.3 Page 2.

Narayanan, K.1980. Studies on the interaction between host *Heliothis  
armigera* (Hubn.). Puthogen (Nuclear polyhedrosis virus ) and  
parasite. *Eucelatoria* sp. R. *armigera* (Cog.). paper presented  
at 3<sup>rd</sup> Annual Workshop on "Biological Control of Crop pests  
and weeds" at ludhiana, Punjab, Oct.,27 to 30<sup>th</sup> 1980.

Nimbalkar, S.A., Kulat, S.S., Dhurve, S.D. and Tambe V.J. 1996.

Bioefficacy of insecticides and biocides for the bollworm management of Cotton. Journal of soils and Crops. 6:2, 162-164.

Nisha, M.S. Sarika, A. Sara, and M.Rai,2003. Synergistic effect of *piriformospora indica*, *Trichoderma viride* and *Pseudomonas fluorescens* in growth promotion of *Withania somerifer* in nursery. JNKVV Research Journal,37 (I) : 67-71.

N overoske, R.I.1975. Dowco (R) 269. a new systemic fungicide for control of *Phytophthora parasitica* of tobacco phytopathology 65: 22-27.

Ocana, G. and P.H. Tsao (1966), "A selective agar medium for the direct isolation and enumeration of *Phytophthora parasitica* in soil." (Abs.), Phytopath., 56:893.

Padwic, G.W. (1940). Report of Imp. Mycologist – Socio. Rep. Agric. Inst. New Delhi, 1938-39 : 103-105.

Pandev, A and Sushil Kumar, K. 1989, Potential of Azotabacter and Azospirillum and bio fertilizers for upland agriculture : A review. J.Sci. Ind. Res. 48 (3) : 134-144.

Papavizas, G.C. 1985. Trichoderma and Gliocladium : Biology, ecology and potential for biocontrol. Ann. Rev. Phytopath; 23:23-54.

Patil M.R. Waghe, S.V., Wangikar, P.D. and khune, N.N. (1982). Chemical Control of wilt of betelvine (Abst.) National workshop on betelvine. March, 4-6,1982. West Bengal Sci. and Tech. Committee, Calcutta, 37 pp.

Pawar, V.P., Magdum S.G., Jadhav R.G., Chougule B.A., and Shirke M.S., Krishna pan (Dpb-6) MPKV. Bulletin No.67 opened in Betelvine Group Discussion. (26/2/2005) held at MPKV Rahuri at the hands Hon. Minister of Agriculture, Govt. Maharashtra.

Pfanz H. 1993. Oxidation of IAA by extracellular peroxidases In : Welinder KG, Rasmussen SK, penel C, Greppin H. eds. Plant peroxidases : biochemistry and physiology. University of Geneva, 169-174.

Poovaiah; B.W. and Rasmussen, H.P. (1973). Peroxidase activity in the obscission zone of bean leaves during abscission. Pl. physiol. 52-263.

121

Pradhan, M.K. (1985). Pathogenic association of *Colletotrichum sp.* Woth  
M. incognita in betelvine, M.Sc. Ag. Thesis submitted to O  
VAT.,Bhubaneswar 64 pp.

Rabindran, R.,Marimuthu , T., and Rajukhann K.1988. Chemical control  
of anthrocnoise of betel vine caused by *Colletotrichum capsici*.  
Indian phytopath., 41:599-602.

Raghovendra RaO, N.N. and Maiti, S.1989. Occurrence of flowering in  
betel vine (*Piper betel*) Indian J. Agric. Sci., 59:605-606.

Rajurkar, R.G., Gade, R.M., Pastawar, A.N., Chauke, R.P.1998.  
Management of betel vine with through cultural and biological  
methods Journal of soils and crops. 8 (2) : 176-178.

Ramachandran, N.1990. Bioefficacy of systemic fungicides against  
*Phytophthora infestans* in black pepper, Ph.D. Thesis,  
University of Calicut, Calicu,138 pp.

Ramchandran, N. and Sharma, Y.R. 1985. Efficacy of three systemic  
fungicides in controlling *Phytophthora infectians* in block  
pepper. Indian phytopathology, 38:160-162.

Ramchandran, N., Sarma, Y.R. and Anandraji, M. 1991. Management of *Phytophthora infestans* in black pepper. In V.R. Sarma and T. Premkumar (Eds.) Diseases of black pepper, National Research for spices, Calicut, India, pp. 158-174.

Rao, M.B.N., Satyanarayam, G., Rameswar, A., Shiv RajAJ, A. and Padmanbham, V. 1988. Bio-chemical basis for root-regeneration in ringed shoot cuttings of cashew (*Anacardium occidentale* L.) plants of different ages – cofactor activity and total phenol content. J. Plant. Crops. 17:65-68.

Rawat, A.K.S., Shama, M.L., Yunus, M. and Balasubrahmanyam, V.P. 1984. Identity of five betelvine (*Piper betel* L.) cultivars. Proc. National Workshop on Betel vine. Eds. S.D. Khanduja and V.R. Balasubrahmanyam, National Botanical Research Institute, Lucknow. pp. 101-104.

Research Journal of Agriculture and Biological Sciences 1 (4):325-328,2005

Saksena, S.B. (1977). *Phytophthora parasitica* the scourge of 'pan' Indian phyopathol. 30. 1-16.

Sardhambal K.K., Singh, S.P., Prakash, S. and Naik, M.S. (1978). Effect of bacterial blight on the activities of nitrate reductase and peroxidase in rice plants. Indian J. Biochem. Biophys. 15 : 105-107.

Sarode, S.V., Jumde, V.S., Deokate, R.O. and Thakare, H.S. 1996. Bioefficacy of Neem Seed Kernel Extract (NSKE) and Ha NPV against *H. armigera* (Hubner) on Cottan. Pkv. Research Journal 20 (I) 96-97.

Scottorraig JS, Kerby KB, Stein BD, Somerville SC. 1995. Expression of an extracellular peroxidase that is induced in barley (*Hordeum Vulgare*) by the powdery midew pathogen (*Erysiphe graminis* f.sp. *hordei*). Physiological and Molecular plant physiology 47, 407-418.

Selvaraj, C. 1966. Studies on the will disease of betel vine (*Piper betel* L.) H. Effect of soil temperature on disease development. M.Sc. (Ag.) Thesis 87 pp. Madras University, Madras.

Sen, A.R. 1967. Responses of mature tea to phosphate and potash in North East India. Exptl. Agric. 3:55-62. Stephan. Z.A.E., Ebhodi, A.a., Zahron, H.M., Antan. B.G. and Geogress, M.S. 1996. Control of root-knot nematode and wilt disease complex

on tomato plants (Research Note) *Dirasat Series B. pure and Applied Sci.*,23:13-16.

Sen, V.1993. Ecology of *Trichoderma harzianum* and *Gliocladium Virens* : two potential biocontrol agents of soil – borne plant pathogens, Final Report, ICAR Project, 115p.

Shree, M.P., Shivakumar, P. Shanthi, K.R. and Nataraja, S. (1997) Influence of powdery midew and leaf spot disease on the enzyme activity in mulberry verities. *Geobias* 24:98-108.

Singh P.P.,Monabruallah. M. and Singin.B.1999. Field efficacy in some microbial pesticides against gram pod borer H. *Ormigera (H)* in Chickpea. *Shashpa* 6 (I) 63-66.

Singh, B.P. and Chand, J.N. (1972). Assessment of losses of betel vine (*Piper betle*) caused by *S. rolfsii* in Jabalpur (M.P.) *Sci. and cult.* 38:526-527.

Sing, K.K. and Balasubrahmanyam, V.R. 1988. Standardization of leaf sampling techniques in betel vine for chemical analysis. *J. Plant. Crops.* 16:105-108.

Singh, K.K., Tripathi, R.D.; Balasubrahmanyam, V.R. and Kochhar, V.K. 1988. Inhibition of changes in chlorophyll, protein, phenolics and cell permeability during senescence of betel leaves by benzyladenine. *Beit.r. Biol.Pflanzen*.62 : 415-427.

Singh, K.K.V.R. Balsubrahmanyam and V.R. Kochhar (1990). Effect of different packing methods, temperature conditions, treatment with Chemicals on the senescence and storage behaviour of betel leaves (*Piper betel* L.) J. of planation crops, 6:10432.

Singh, R.A. and Shankar, G.1971. Some parasitic fungi on *Piper betle* L. in Varanasi, V.P. Mycapath Mycol. Appl. 73 :109-115.

Singh, R.S. 1978. Introduction of plant pathology, oxford and IBH Pub. New Delhi pp. 210-217.

Singh, V.P. (1997). Interaction of temperature and microsomal peroxidase in aflotoxin degradation by *Aspergillus flavus* 102566. Curr. Sci. 72:529-532.

Srinivasan, G. Sundara Babu, P.C. Sathiah, N. and Balasubramanian, G.1994. Field efficacy of Ha NPV alone and in combination with delfin for the control of Gram Pod borer, *H. armigera* (H) on chickpea. Pest Management and Economic Zoology Z (1) 45-48.

Stevan, A. and J. Chet (1989) : The possible role of competition between *Trichoderma harzianum* and *Fusarium oxysporum* on rhizosphere colonization. Phytopathology. 79 (2) : 198-209.

Stitt, M., Van Schaewen, A. and Willmitzer, L. (1991). 'Sink; regulation of photosynthetic metabolism in transgenic tobacco plants expressing yeast invertase in their cell wall involves a decreases of the calvin Cycle enzymes and increases of Glycolytic enzymes. *Planta* 183:40-50.

Stonier, T., Stasions, S., and Murthy, K.B.S. (1979). The masking of peroxidase catalyse oxidation of IAA in Vigna. *Phyto. Chemistry* 18, 25.

Subramaniam, D. (1998). Effect of aluminium on growth lipid peroxidation, superoxide dismutase and peroxidase activities in rice bean and fresh bean seedlings. *Indian J. Plant physiol.* 3:240-242.

Summour, R.H. (1989). Effect of aging on carbohydrates, proteins and their related enzymes in natural aged seeds of flax. *Pakistan J. Biochem.* 22 (2) : 81-87.

Sumer, A.T. (1968). A substrate – film method for histochemical localization of cellulase. *Histachemie*, 13, 160.

Sung, F.J.M. and Chiu, C.C. (1995). Lipid peroxidation and peroxide – Scavenging enzymes associated with accelerated ageing of peanut seed. *Physiol. Plant.* 91 :51-55

Sung, J.M. (1996). Lipid peroxidation and peroxidase scavenging in sayabean seeds during aging. *Physiol. Plant.* 97 : 85-89.

Swaminathan, V., Sambanda – Moorthv, S. and Appavu, K.A.1993, Effects of *Azospirillum* on the yield and quality of *Amaranthus spp.* Golden Jubilee symposium. Horticultural Research – changing Scenario, Bangalore, May 24-28, India.

Takahama U. Oniki T.1992. Regulation of peroxidase oxidation of phenolics in apoplast of spinach leaves by ascorbate. *Plant and cell physiology* 33, 379-387.

Talekar, N.S. and Shelton, A.M.1993. Biology,ecology and management of the diamond back moth. *Ann. Rev. Entomol* 38:275-301.

Thomas, K.M. and T.S.Ramakrishnan (1948), Studies in the genus *Phytophthora-II*", Porac. Indian Acad.,Sci.,27 (B):55-73.

Tiwari, D.P. and R.S. Mehrotra (1968) studies on foot rot and leaf rot diseases of (*Piper betel* L.) *Bull. Indian phytopathol. Soc.* 4; 79-89.

Tiwari, D.P. (1968), Studies on the Foot Rot and Leaf Rot Disease of *Pipe betle* Linn. With special reference to Microbiology and Measures of Control," Ph.D. Thesis, Univ. of Sagar.

Tiwari, M.M. and H.C. Arya (1966), "Studies on green ear disease of bajara (*Fennisetum Hyphoideum* Stapf, and Hubb.) Caused by *Scierospara graminicola* axenic Cullure," Science, New York, 143:291-293.

Trujillo, E.E. and R.B. Hine (1965), "The rate of papaya residues in papoya root rot caused by *Pythium aphaniderma* tum and *Phytophthora parasitica*" Phytopath., 55:1293-1298.

Trujillo, E.E.(1965),"The effects of humidity and temperature, on *Phytophthora* blight of Tara,"Phytopath, 55:183-188.

Trujillo, E.Edvardo (1963): Fusorium yellows and rhizomes rot of qinger. Phytopathology. 53:1370-1371.

Tucker, C.M. (1931), "Taxonomy of the genus phytophthora de Barj", Missown Agr. Exp. Sta. Res. Bull.,153=208.

Turner, P.D. (1969), "Phytophthora palmivora from *Piper betle* in Sarawak,"  
Trans. Brit. Mycol. Soc., 52:411-418.

Uppal, B.N. and M.K. Desai (1932), "Physiologic specialization in  
*scleraspara graminicala* (Sacc.) Schroet., "Indian J. Agr. Sci.,  
2:667-678.

Van Fleet, D.S. (1947). The distribution of peroxidase in differentiating  
tissues of Vascular plants. Biodynamica 6,125.

Vasudeva, R.S. 1962. Fungi of India suppl. I, PP/206, ICAR New Delhi.

Vasudeva, R.S. and R.N. Azad (1962), "Efficacy of certain fungicides,  
against potato late blight and assessment of loss due to the  
disease, : Am. Potato J.,29:61-71.

Veech, J.A. (1969). Localization of peroxidase in infected tobacco  
susceptible and resistance to black shank.  
Phytopathology.59-566.

Vyas, K.M. and Chourasia, S.C. (1978). Leaf surface microflora in relation  
to *Phytophthora* leaf rot of pan. *Phytophthora* News letter  
No.6 : 41-42.

Vyas, K.M. and S.C. Chaurasia (1976), "Effect of fungicides on the oxidative metabolism of *Phytophthora parasitica* var. *piperina*

Vyas, K.M. Chile, S.K. and Vyas, R.S. 1981. Effect of Culture filtrate of various fungi on sporulation of betel vine *Phytophthora*. Hindustan Antib. Bull. 23: 25-26.

Wastie, R.L. (1961), "Factors affecting competitive saprophytic colonizer of the agar plate by various root infesting fungi," Trans. Brit. Mycol. Soc., 44:145-159.

Waterhouse, G.M. (1956), "The genus *Phytophthora*, diagnoses or descriptions and figures from the original papers," No.12. Misc. Publ., C.M.I., Kew, Surrey England.

Webster, B., Dunlap, W.T., and Craig, E.M. (1976). Ultrastructural Studies of abscission in *Phaseolus* : Localization of peroxidase. Amer. J. Bot. 63,759.

Weimberg, R. (1975) Effect of growth in highly salinized media on the enzymes of the photosynthetic apparatus in Pea Seedlings. Plant Physiol. 56:8-12.

Weiss, D.2000. Regulation of flower pigmentation and growth: multiple signaling pathways controls anthocyanin synthesis in expanding petals, *physiology plantarum*, 110:152-157.

Wood, R.K.S. (1967), *Physiological Plant Pathology* Oxford Endingbugg.  
Blackwell Scientific publ.

Wutscher, H.K.,C. hardesty (1979). Zinc and phenolics in the wood and incidence of citrus blight in a grove containing eight selections of 'Hamlin Orange' proc. Fla. Stake Hortic. Soc. 92 : 70-72.

Wyckoff, H....W., Handschumacher, M.,Murthy,H.M.K. and Sewdski, J.M. (1983). The three dimentional structure of alkaline phosphatase from *E. coli* In:*Advances in enzymology*, New York, Chichester, Brisbane, Taranto, Singapore. 435-480.

Yarkins, D.M. (1941). Activity of Amylase and catalase in Wheat leaves infected with rust *Puccinia tritici* under Irrigated condition.  
DOCKL, Veses, ACAD. S. kt. Nauk V.I. Leuina : 29-32.

Yarwood, C.E. and L. Jacobson (1955). Accumulation of Chemicals in diseased areas of leaves *Phytopathol.* 55:43-48.

Zheng X, Van Huystee RB. 1992. Peroxidase – regulated elongation of segments from peanuts hypocotyls. *Plant Science* 81, 47-56.