

## LIST OF TABLES

TABLE No.	TITLE	PAGE No.
3.1	Calculated $H_x$ values for $N = 0$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>65</b>
3.2	Calculated $H_x$ values for $N = 1$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>65</b>
3.3	Calculated $H_x$ values for $N = 2$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>66</b>
3.4	Calculated $H_x$ values for $N = 0$ mode [ $2\rho = 200 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>67</b>
3.5	Calculated $H_x$ values for $N = 1$ mode [ $2\rho = 200 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>67</b>
3.6	Calculated $H_x$ values for $N = 2$ mode [ $2\rho = 200 \mu\text{m}$ , $\lambda = 0.85 \mu\text{m}$ ]	<b>68</b>
3.7	Calculated $H_x$ values for $N = 0$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 1.3 \mu\text{m}$ ]	<b>69</b>
3.8	Calculated $H_x$ values for $N = 1$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 1.3 \mu\text{m}$ ]	<b>69</b>
3.9	Calculated $H_x$ values for $N = 2$ mode [ $2\rho = 100 \mu\text{m}$ , $\lambda = 1.3 \mu\text{m}$ ]	<b>70</b>
4.1	Calculated values of $\epsilon_2^{1/2}  E(z) $ as a function of $z/\lambda$	<b>99</b>
4.2	Critical power of the surface wave for the linear and a positive nonlinear medium as a function of $d/\lambda$	<b>100</b>