

***BIBLIOGRAPHY***

**BIBLIOGRAPHY**

- Abbasi, S.A. and R. Soni (1984) : Toxicity of lower than permissible levels of chromium (VI) to the freshwater teleost *Nuria denricus*. Environ. Pollut. Ser. A. Ecol. Biol. 36(1) : 75-82.
- Abbasi, S.A., V. Baji, K. Madhavan, R. Soni (1991) : Impact of chromium (VI) on catfish *Wallago attu*. Indian J. Environ. Hlth. 23(3), 336-390.
- Abbasi, S.S., T. Kunhahmed, P.C. Nipaney and R. Soni (1995); Influence of the acidity on chromium toxicity. A study with the teleost, *Nuria denricus* as model. Poll. Res. Vol.14(3): 317-323.
- Adams, S. Marshall, C.A. Burtis and J.J. Beauchamp (1985) : Integrated and individual biochemical responses of rainbow trout (*Salmo gairdneri*) to varying duration of acidification stress. Comp. Biochem. Physiol. C. Comp. Pharmacol. Toxicol. 82(2) : 301-310.
- Agarwal, A., Sharma, R. and Sharma, K. (1982): State of India's Environment A Citizen Report, Centre for Science on Environment, New Delhi.
- Agarwal, S.L. and A.K. Gautam (1988) : Chronic toxicity of distillery effluent to freshwater fish *Channa punctatus* (Bl). Acta. Ecologica. 10(1) – 9-18.

- Ajimal, M. Raziuddin and A.U. Khan (1985): Physico-chemical Aspect of pollution in Kali Nadi. IAW PC Tech. Annual, XII.106-114.
- Akhilender, N. K., N.K. Abhinender and R. Ramamurthy (1983): Histological observation in gills of the teleost fish, *Sarotherodon mossambicus* with reference to mercury toxicity. *Ecotoxicol and Environ. Safety* Vol.5:455-462.
- Alis, A.A. and J.K. Shamsi (1988) : Behaviour responses imputed by contamination of water with mercuric chloride and zinc sulphate. *J. Biol. Sci. Res.* 19, 949-956.
- Ambrose, T., L. Cyril, A. Kumar, S. Vixent and R. Lambert (1994): Biochemical responses of *Cyprinus carpio* (Communis) to toxicity of tannery effluent. *Journal of ecobiology* 6(3): 213-216.
- Amudha P., R. Nagendran and S. Mahalingam (1997) : Studies on the effects of dairy effluents on the behaviour of *Cyprinus carpio* (Cyprinidae). *J. Environ. Biol.* 18(4): 415-418.
- Amudha, P. and S. Mahalingam (1999): Effect of dairy effluent on the body composition of *Cyprinus carpio* (Cyprinidae). *J. Ecotoxicol. and Environ. Monit.* 9(1): 03-08.
- Amudha, P., S. Mahalingam (1999): Studies on the effect of dairy effluents on survival and feeding energetics in *Cyprinus carpio*. *J. Environ. Biol.* 20(3): 275-278.

- Anjaneyulu, C., P. Reddy, S. J. Tulasi, J.V. Ramana Rao and P.U.M. Reddy (1987-1988) : Toxic effect of chromium on certain metabolites of *Anabus scandens* as a function of pH. J. Aqua. Biol. Vol. 5(1-2), 1987-88 : 57-60.
- APHA. AWWA and WPCF (1985): Standard methods for the examination of water and waste water. American Public health Association, American water works association and Water pollution control federation. 16<sup>th</sup> Edn. American Public Health Association, Washington, D.C.
- Arilo, A.C., F. Melodia, P. Melodia and P. Menst (1983): In : Influence of sex and season. Chemosphere, 11: 47-57.
- Awari, S.B. and S.A. Gaikwad (1990): Cadmium toxicity on *Ambassis ranga* (Ham.). Proc. of the Symp. Trends in Ecotoxicolgy. Ed. R.C. Dalela pp.89-92.
- Badrinath, S.D., V. Raman and V. Agumugam (1983): Upgrading An Existing Waste Water Treatment for Textile Industry. A Case Study. IAWPC Tech. Annual, X: 55-73.
- Balaji, A. and S. Chockalingam (1990) : Effect of sublethal concentration of dairy effluent on an air breathing fish, *Channa punctatus*. Proc. (Vol.2) 10<sup>th</sup> Ann. Sess. Acad. Environ. Biol. And seminar "Environ. Impact Biosystems" Madras, Dec. 13-17 (1989), 75-80.
- Balavenkatesubbaiah, M., A. Usha Rani, K. Geethanjali, K.R. Purushottam and R. Ramamurthy (1984) : Effect of cupric chloride on oxidative metabolism in freshwater teleost fish, *Tilapia mossambica*. Ecotoxicol. And Environ. Safety, Vol. 8(3) : 289-291.

- Barnes, H. and Black-stock (1973): Estimation of lipid in marine animals, detail investigation of sulphophospho vanillin method. *J. Exp. Mar. Biol. Ecol.*, 12:103-118.
- Bath, K.S. and H. Kaur (1999): Physico-chemical characteristic of water of Buddha-Nallah (Ludhiana, Panjab), *Ind. J. Environ. Sci.* 3(1): 27-30.
- Bengeri, K.V. and H.S. Patil (1986) : Respiration, liver glycogen and bioaccumulation in *Labeo rohita* exposed to zinc. *Indian J. Comp. Animal Physiology* 4(2) : 79-84.
- Bengeri, K.V. and H.S. Patil (1995) : Uptake of zinc by two freshwater fishes *Lepidocephalichthes guntea* and *Labeo rohita*. 3<sup>rd</sup> Cong. On Toxicol. in Developing Countries Cairo, Egypt.
- Bengtesson, B.E. (1974): Vertebral damage to minnows, *Phoxinus phoxinus* exposed to zinc. *Oikos* 25(3): 134-139.
- Bengtesson, B.E. (1975): Vertebral damage in fish induced by pollutant. In: Sub lethal effect of toxic chemical on aquatic animal. (Eds.: J.H. Kolman and J.J.T.W.A. Strik) Amsterdam: Elsevier, pp. 23-30.
- Benoit, D.A. (1975) : Chronic effect of copper on survival, growth and reproduction of the blue gill (*Lepomis macrochirus*). *Trans. Am. Fish. Soc.* 104(2) : 353-358.
- Benoit, D.A. (1976); Toxic effects of hexavalent chromium on brook trout, *Salvelinus fontinalis* and rainbow trout, *Salmo gairdneri*. *Water. Res.* Vol.10: 497-500.

- Bhaskar, M. and S. Govindappa (1985): Tissue compensatory metabolic profiles in *Tilapia mossambicus* on acclimation to sublethal acidic and alkaline media : Gill glycogen metabolism. Arch. Int. Physiol. Biochem. 93(2) : 59-64.
- Bhaskar, M. and S. Govindappa (1986): Effect of environmental acidity and alkalinity on the physiology of *Tilapia mossambica* during acclimation. Biochem. Syst. Ecol. 14(4): 439-444.
- Bhaskar, M., P. Sobha Rani, M.V. Chalapathy Rao, P. Reddanna and S. Govindappa (1986) : Hepatic tissue lipid profile in freshwater fish *Oreochromis mossambicus* acclimated to acidic and alkaline media. GEOBIOS, 12(5) : 200-203.
- Bhattacharya, S. and K. Mukherjee (1976 ) : Activity of hepatopancreatic protease and esterase in fish exposed to industrial pollutants. Comp. Physiol. Ecol. 1, 45-50.
- Blinski, E. and R.E.E. Jonas (1973) : Effect of cadmium and copper on the oxidation of lactate by rainbow trout (*Salmo gairdneri*) gills. Water Res. 10, 37-43.
- Bradley, R.W. and J.B. Sprague (1985) : The influence of pH, water hardness and alkalinity on the acute lethality of zinc to rainbow trout (*Salmo gairdneri*). Can. J. Fish. Aquat. Sci. 42(4) : 731-736.
- Bradley, R.W. and J.B. Sprague (1985): Accumulation of zinc by rainbow trout (*Salmo gairdneri*) as influenced by pH, water hardness and fish size. Environ. Toxicol. Chem. 4(5): 685-694.

- Burton, O.A. and J. Cairns (1972): Acute zinc toxicity of rainbow trout *Salmo gairdneri* confirmation of hypothesis that death related to tissue hypoxia J. Fish. Res. Bd. Canada 29: 14-63.
- Cairns, J., A.G. Heath and B.C. Parker (1975): Temperature influence on chemical toxicity to aquatic organism. J. Wat. Poll. Cont. Fed. Vol. 47(2): 267-280.
- CDC (1982); Centre for Disease Control Surveillance of Childhood lead poisoning- United States, MMWR 31:132-134.
- Chandravathy, V.M. and S.L.N. Reddy (1994): In vivo recovery of protein metabolism in gill and brain of a freshwater fish, *Anabus scandens* after exposure to lead nitrate. J. Environ. Biol. 15(1): 75-82.
- Charles, A., W. Nalland and L.D. Betz (1945): Determination of nitrates in Bioler water by Brucine. Indul. and Eng. Chem. (Anly. Edn.) 17 (7): 426-427.
- ✓ Chauhan, S.V.S., J. Kumar and T. Singh (1993): Physico-chemical characteristics of effluents from tanneries. Acta. Ecologica. 15(1): 39-42.
- Chauhan, R.R.S., and K.K. Saxena (1992): Effect of temperature on the toxicity of some metals to *Heteropneustes fossilis* (Bl.) Polln. Res. 11(3): 131-134.
- Chitra, K.C. (1983) : Induction of biosynthesis of proteins and amino acids in the tissue of *Sarotherodon mossambicus* under ambient urea stress. Indian J. Comp. Anin. Physiol. 1(2): 37-39.

- CIFRI (1981) : Central Inland Fisheries Research Institute. Annual Report Project No.2011, CIFRI Barrackpur, India.
- Coughlan, J. David, S. P. Gloss and J. Kubota (1986) : Acute and sub-chronic toxicity of lead to the early life stages of small mouth bass (*Micropterus dolomieu*). Water, Air, Soil Pollut. 28 (3/4) : 265-276.
- Crasser, M. and A. Edwards (1987): Acidification of fresh water Cambridge University, Press, New York.
- Crespo, S. (1984): An in Vitro study of the effects of Zinc on osmoregulatory processes. Mar. Pollut. Bull. 15(9): 341-342.
- Crook, J.G. (1964): Hand book of textile fibre. Merrow Publishing Co. Ltd., Herts, England.
- Das, S.M. (1978): High Pollution in lake Nainital, U.P. as evidenced by biological indicators. Sci. and Cult. 44, 236-237.
- Datta, S.P.S. and N. Zutshi (1988) : Effect of sewage effluents on the egg of *Cyprinus carpio* (Linn). National Academy of Sciences, India. 58<sup>th</sup> Annual Session Jammu, Nov. 5-7.
- Dave and Goren (1985) : the influence of pH on the toxicity of Al, Cd, and Fe to eggs and larvae of the zebra fish *Brachydanio rerio*. Ecotoxicol. Environ. Saf. 10(2) : 253-267.
- David, A. and P. Ray (1966) : Studies on the pollution of river Daha (N. Bihar) by Sugar and distillary waste. Indian J. Environ. Hlth. 8, 6-35.



- Delamarre, O.P. and M. Truchet (1985) : Comparative study of mercury and cadmium exposure in a freshwater fish (*Brachydonio rerio*) : Histological microanalysis of cadmium induced mineral concentrations. VIE MILIEU 34 (2/3) : 79-86.
- DeZwaan and D.I. Zandee (1972): Body distribution and seasonal changes in glycogen content of the common sea mussel, *Mytilus edulis*. Comp. Biochem. Physiol. 43: 53-55.
- Dhande, R.R. and G.P. Patil (1998) : Effect of heavy metal pollutants on the endocrine kidney of the fish *Channa punctatus* (Bloch). J. Environ. Biol. 7(2), 149-154.
- Dhanpal, S. and A.A. Sivakumar (1988) ; Effect of tannery effluent on the water quality of the river Cauvery (Tamil Nadu). Himalayan J. Env. Zoo. 2 (2): 142-145.
- Dharwadkar, M. M. and Y.H. Deshpande (1990) : Heavy metal in industrial and domestic sewage sludge. Proc. Symp. Trends in Ecotoxicology, 13-17 Dec. (1989) Madras, 127-130 AEB.
- Doudoroff, P. and M. Katz (1953) : Critical review of literature on the toxicity of industrial waste and their components to fish II. The metals as salts. Sewage Ind. Wastes. 25, 802-832.
- Elizabeth, T.K., N.K. Balasubramaniam and P.A. John (1981) : Effect of alkaline and acidic fraction of industrial effluents on some lymphoid cells of the fish, *Rasbora daniconius*. Environ. Pollut. Ser. A., Ecol. Biol. 25(2) : 135-138.

- Ellgaard, E.G. and J. Y. Gilmore, III (1984) : Effects of different acids on the blue gill sunfish *Lepomis macrochirus*. J. Fish. Biol. 25 (2) 133-138.
- Eyanoer, F. Harwinta, E.S. Upatham, M. Duangsawasdi and S. Tridech. (1985): Effect of water hardness and temperature on the toxicity of detergents to the fresh water fish, *Puntius gonionotus*. J. Sci. Thail. 11(2): 67-78.
- Eye, J.D. and L. Lawrence (1971) : Treatment of wastewater from a sole leather tanner. J. Water Pollut. Cont. Fed. 43. 2291-2303.
- Fraser, J.A. L. and E.R. Clark (1984) : The effect of a settled industrial domestic sewage works effluent from percolating filters on the embryo viability and hatching success of rainbow trout *Salmo gairdneri*. J. Fish. Biol. 25(4): 393-404.
- Gandheeswari, P., A. Bernice and M. Selvanayagan (1989) : Effect of tannery effluent on rotifer population in relation to physico-chemical factor. Environmental Impact on Biosystem 267-274. Proc. Of the 10<sup>th</sup> Ann. Ses. of the AEB. Madras.
- Ghosh, T.K. and S.K. Konar (1992) : Effect of sugar mill complex effluent on fish growth and reproduction. Env. Eco., 10(4): 794-803.
- Gill, T.S. and J.C. Pant (1985) : Effect of organomercurial poisoning on the peripheral blood and metabolite level of a freshwater fish. Ecotoxicol. Environ. Saf. 10(2) : 150-158.

- Gill, T.S. and J.C. Pant (1985) : Erythrocytic and Leukocytic responses to cadmium poisoning in a freshwater fish *Puntius conchoni* (Ham). Environ. Res. 36(2) : 327-337.
- Goel, K.A. and S.D. Sharma (1987) : Some haemato-chemical characteristics of *Clarias batrachus* under metallic stress of Arsenic. Comp. Physiol. Ecol. 12(2) : 63-66.
- Gornall, A.G., J. C. Bardawill and M. M. Devid (1949): Determination of serum protein by means of the biurate reagent. J. Biol. Chem. 177:751-766.
- Govindan, V.S. and V.S. Sundaralingam (1979) : Studies on the treatment of textile mill waste by Stabilization Pond method. Ind. J. Environ. Hlth. 21 (4): 321-331.
- Guru Prasada Rao, M. and N.V. Nanda Kumar (1981): Analysis of irrigation reservoir contaminated by tannery effluent. Ind. J. Environ. Hlth. 23(3), 239-241.
- Haesloop, UWE and M. Schirmer (1985) : Accumulation of orally administered cadmium by the eel (*Anguilla anguilla*). Chemosphere 14(10) : 1627-1634.
- Haniffa, M.A. and S.A. Selvan (1991) : Relative toxicity of bleaching, dyeing and mixed textile mill effluent to fresh water fish *Oreochromis mossambicus* (Trewaves). J. Environ. Biol. 12(4): 359-362.
- Helmy, M.M., A.E. Lemke, P.G. Jacob and Y.Y. Alsultan (1979): Hematological changes in Kuwait mullet, *Liza macrolepis* induced by heavy metals. Indian J. Mar. Sci. 8: 278-281.

- Hingorani, N.G., A.O. Diwan, N. Naidu Chandrashekharan (1979) :  
Oxygen consumption on the fish *Labeo rohita* during  
exposure to different concentrations of industrial effluent.  
Comp. Physiol. Eco. 4(4), 272-276.
- Hiraoka, Yukio, I. Shoichi, K. Toshihiko and O. Hasanori (1985) :  
Acute toxicity of 14 different kinds of metals affecting  
medaka (*Oryzias latipes*) fry. Hiroshima J. Med. Sci. 34(3) :  
327-330.
- Howells, G., T.R.K. Dalziel, J. P. Reader and J.F. Solbe (1990) :  
EIFAC Water quality criteria for European freshwater fish :  
Report on aluminium. CHEM. ECOL. 4(3) : 117-173.
- Hughes, M., A.M. Margatto, S. Morrison and W. Berry (1989): An  
evaluation of a short term chronic effluent toxicity test using  
sheephead minnow larvae. Environ. Pollut. 60 (1/2), 1-14.
- Hutchinson, N.J. and J.B. Srague (1986) : Toxicity of trace metal  
mixtures to American flagfish, *Jordoella floridae* in soft  
acidic water, implication for cultural acidification. Can. J.  
Aquat. Sci. 43 : 646-655.
- IRPTC (1984); Chemical safety International Registrar of  
Potentially Toxic chemicals. UNEP-Geneva-pp.269.
- Iyr, H. (1977): Anti fungal compounds, M.R. Seigle and H.D. Sister  
Eds., Vol. 2. Decker, New York, 301-332.
- James, R., K. Sampath, V. Sivakumar, S. Babu and P.  
Shaniganadam (1995): Toxic effects of copper and mercury  
on food intake, growth and proximate chemical composition  
in *Heteropneustes fossilis*. J. Environ. Biol. 16(1): 1-6.

- Jana, S. and S. Bandyopadhyaya (1987) : Effect of heavy metal on some biochemical parameters in the freshwater fish *Channa punctatus*. Environ. and Ecol. 5, 488-493.
- Jana, S., S.S Shhana, M.A. Choudhari and D.K. Choudhari (1986) : Heavy metal pollutant induced changes in some biochemical parameters in the freshwater fish *Clarias batrachus*. Acta. Physiol. Hung 68(1) : 39-44.
- Janiyani, K.L., S.R. Wate and S.R. Joshi (1993): Physico-chemical characteristics of oil sludges from various sources of refinery. Proc. Acad. Environ. Biol. 2(2): 149-155.
- Jesudass, L. and R. Akila (1995) : Studies on the physico-chemical characteristics of the sugar factory effluents. IJEP 16(11): 808-810.
- Jha, B.S. (1991): Alteration in the protein and lipid content of intestine, liver and gonads in the lead exposed freshwater fish *Channa punctatus* (Bloch). J. Ecobiol. 3:29-34.
- Jha, B.S. (1999): Impact of chronic exposure of the household detergents, Surf and Key on tissue biochemistry of the freshwater fish *Clarias batrachus* (Linn). Indian J. Environ. Ecoplan. 2(3); 281-284.
- Jha, B.S. and A.K. Jha (1995a): Biochemical profiles of liver, muscle and gonads of the freshwater fish *Heteropneustes fossilis* under chromium stress. In : Toxicity and monitoring of Xenobiotics (Eds. R. Prakash and P. Sood). Venus Publishing House, New Delhi pp. 127-137.

- Jha, B.S. and M.M. Jha (1995b): Biochemical effects of nickel chloride on the liver and gonads of freshwater climbing perch *Anabus testudineus* (Bloch). Proc. Nat. Acad. Sci. India 65 B : 39-46.
- Jha, B.S. and S. Pandey (1989) : Alteration in the total Carbohydrate level of intestine, liver and gonads induced by lead nitrate in the fish, *Channa punctatus*. In: Environ. Risk Assessment (Eds.V.N. Sahai, P.N. Deshmukh, T.A. Mathai and K.S. Pillai). The Academy of Environmental Biology, Muzaffarnagar, India pp.207-211.
- Johnsen, K., K. Mattsson, J. Tana, T.R. Stuthridge, J. Hemming and K.J. Lehtinen (1995) : Uptake and elimination of resin acids and physiological responses in rainbow trout exposed to total mill effluent from an integrated newsprint mill. Environ. Toxicol. Chem. 14(9) : 1561-1568.
- Joseph, T.A. (1992) : Behavioural alterations induced by nickel and chromium in common carp *Cyprinus carpio*. Env. Ecol. 10(4) : 911-913.
- Joseph, T.A., S.M. Sebastin and S. Raja (1992): Toxicity of nickel on protein content in tissue of *Cyprinus carpio* (Linn). Indian J. Environ. Hlth. 34(3): 236-238.
- Kallanagaudar, Y.P. and H.S. Patil (1997): Influence of water hardness on copper, zinc and nickel toxicity to *Gambusia affinis* (B & G). J. Environ.. Biol. 18(4): 409-413.

- Kane, A. Donald and R.F. Charles (1987); Effects of aluminium and pH on the early life stages of smallmouth bass (*Microterus dolomieui*). Water. Res. 21(6): 633-640.
- Katti, S.R. and A.G. Sathyanesan (1984) : Changes in tissue lipid and cholesterol content in the catfish *Clarias batrachus* (L) exposed to cadmium chloride. Bull. Environ. Contam. Toxicol. 32, 486-490.
- Kaur, K. and K. Bajwa (1987) : Effect of zinc and cadmium on early life stages of common carp, *Cyprinus carpio*. Cnn. Biol. 3, 28-33.
- Kaviraj, A. (1989) : Heavy metal concentration of fish and prawn collected from Hooghly estuary. Sci. Cult. 55, 257-259.
- Kaviraj, A. and S.K. Konar (1983): Lethal effects of mixture of mercury, chromium and cadmium on fish. Environ. Ecol. 1:153-158.
- Kennedy, C.J., R.M. Sweeting, J.A. Honasen, A.P. Forrell and B.A. Mckeown (1995): Acute effects of chlorinated resin acid exposure on juvenile rainbow trout *Oncorhynchus mykiss*. Environ. Toxicol. Chem. 14(6): 977-982.
- Khambe, S.D. (1992) : Effects of industrial effluents on edible fishes and fauna of Krishna river near Sangli. Ph.D. thesis submitted to Shivaji University, Kolhapur.
- Khambe, S.D. (1992) : Effects of industrial effluents on edible fishers and fauna of Kirshna river near Sangli. Ph.D. thesis submitted to Shivaji University, Kolhapur.

- Khan, E.A., M.P. Sinha, N. Saxena and P.N. Mehrotra (1992): Biochemical effect of cadmium toxicity on a hill stream teleost *Garra mullya* (sykes) during egg maturation II cholesterol and glycogen. *Poll. Research* 11(3): 163-167.
- Khangarot, B.S. and R.C. Somani (1980): Toxic effect of mercury on the gills of freshwater teleost, *Punctius sophore* (Ham) *Curr. Sci.* 49(21)L 832-834.
- Khangarot, B.S. and V.K. Rajbanshi (1979) : Toxicity of zinc to a freshwater teleost, *Rasbora daniconius*. *Hydrobiologia* 65 (2) : 141-144.
- Khanna, S.K. (1991): Toxicity, carcinogenic potential and chronical epidemiological studies on dye and intermediates. *J. Sci. Ind. Res.* 50, 965-975.
- Kothandaraman, V.P., Murahari Rao and C.A. Sastry (1972) : Characteristic of water from three typical tanneries in Madras. In : Treatment and disposal of tannery and slaughter hosue waste (Eds. V.S. Krishnamurthy, C.A. Sastry and R. Bhaskaran). CLRT, Madras 1-17.
- Kumar, A. and A.K. Sharma (1999): Physico-chemical characteristics of river Sengar in district-Etawah (U.P.). *Ind. J. Environ. Sci.* 3(2): pp. 201-203.
- Kumar, H.D. , G.P. Bisaria, L.M. Bhandar, B.C. Rana and V. Sharma (1974): Ecological studies on algae isolated from the effluents of an oil refinery, a fertilizer factory and a brewery. *Ind. J. Environ. Hlth.* 10(3); 247-255.



- Kumar, N. J., K.P. Krishnamoorthy and D.M. Rao (1986): Toxicity and biochemical responses of carp (*Cyprinus carpio*) to dinitrobenzene plant effluent. *Water Air Soil. Pollut.* 28(1/2) : 117-126.
- Labat, R., C. Roqueplo, J.M. Ricard and M. Burgat (1977): Ecotoxicological action of some metals (Cu, Zn, Pb, Cd) on fresh water fish in the river. *Lot. Ann. Limnol.* 13(2): 191-208.
- Labat, R., J. Pequignot and A. Chatelet (1974): Toxic action of copper on the gills of carp (*Cyprinus carpio*) *Ann. Limnol.* 10(1); 109-114.
- Lawe-J. L. and A.J. Nivmi (1984): Short term and long term effects of cadmium on glycogen reserves and liver size in rainbow trout (*Salmo gairdnerii* Richardson). *Arch. Environ. Contam. Toxicol.* 13, 759-53.
- Lehtinen, K. Johan, M. Notini and L. Landner (1984) : Tissue damage and parasite frequency in flounders, *Platichthys flesus*, chronically exposed to bleached kraft pulp mill effluent. *Ann.Zool. Fenn.* 21(1) : 23-28.
- Lohar, P.S. (2000): Comparative toxicity of four heavy metals to freshwater fishes. *J. Aquat. Biol.* Vol.15(1&2): 95-98.
- Magre, S.R. (2000): Influence of acidity on zinc toxicity a study with the freshwater fish, *Cirrhinus mrigala*. *J. Aquat. Biol.* Vol.15(1&2): 101-104.

- Mahapatra, S. P. (1987) : Heavy metal concentration in the industrial effluents discharge to the Thane Creek. Indian Environ. Prot. 7 (4), 284-286.
- Mane, P. S. (1988) ; Studies on the effects of aquatic pollution on the physiology of digestion of commercially important common edible fish in the Panchaganga river Ph.D. Thesis, Shivaji University, Kolhapur.
- Manson, C.F. (1987) : A survey of mercury, lead and cadmium in muscle of British (UK) freshwater fish. Chemosphere 16(4) : 901-906.
- Martin, T.R. and D.M. Holdrich (1986): The acute lethal toxicity of heavy metals to peracarid crustaceans with particular reference to freshwater acellids and gammarids. Wat. Res. 20: 1137-1147.
- Martinez, T., G. Viscor and J. Palmomeque (1988) : Effect of temperature oxygen and carbondioxide on osmotic fragility of carp, *Cyprinus carpio* L., erythrocytes. J. Fish. Biol. 32(2) : 247-252.
- Maruthi, Y.A. and M.V. Subba Rao (2000): Effect of distillery effluent on biochemical parameters of fish, *Channa punctatus* (Bloch). Journal of Environment and pollution 7(2): 2000, 111-113.
- Mason, C.F (1981) : Biology of freshwater pollution. Longman Group Ltd.

- Matthiessen, P. and A. Brafield (1973): The effect of dissolved zinc on the gill of the stickleback *Gasterosteus aculeatus* J. Fish. Biol. 5; 607-613.
- McKee, T. and D. Wolf (1971) : American Sewage Practice. Vol. III. Disposal of sewage McGraw Hill Book Company.
- Meenakshi, V., N. Indira and K. Pugazhendhi (1998) : Influence of zinc sulphate on the freshwater edible fish *Labeo rohita* fingerlings. Uttar Pradesh J. Zool. 18(3) : 181-183.
- Metlev, V.V., A.L. Kanaev and N.G. Xzasokhava (1974) : In water toxicology, Translation of Vodnaya Toksikologiya, Kolos Publishers, Moscow (American Publishing Co. Pvt. Ltd.) New Delhi.
- Michibata, H., S. Sahara and M.K. Kojima (1986) : Effect of calcium and magnesium ions on the toxicity of cadmium to the eggs of the teleost, *Oryzias latipes*. Environ. Res. 40(1) : 110-114.
- Miller, T.G. and W.C. Mackay (1980): Effect of hardness, alkalinity and pH of test water on the toxicity of Cu to rainbow trout (*Salmo gairdneri*). Water Res. 14(2): 129-134.
- Mishra, S., D. Sharma and V.Melkania (1988) : Physico-chemical characteristics of effluents from a textile mill at Rampur and their effect on fish mortality. IXth Annual Session of and National Symp. on Fish. Assess. and Management of Acad. Env. Biol. India at Valvada (Gujrat).
- Mishra, S.R. and D.N. Saksena (1988) : Industrial effluent pollution at Birlanagar, Gwalior. IXth Annu. Session and

National Symp. on Fish. Assessment and Management of Acad. Env. Biol. India at Valvada (Gujrat).

Mishra, S.R. and D.N. Saksena (1989): Industrial effluent pollution at Birla Nagar, Gwalior. Poll. Res. No. 8(2): pp 77-86.

Moore, J.W. and S. Ramamoorthy (1984) : Copper. In : Heavy metal in natural waters – Applied monitoring and impact assessment (Ed.) R.S. Desanto Springer – Verlag, New York, Berlin, pp. 77-79.

Mount, D.I. (1966) : The effect of total hardness and pH on acute toxicity of Zinc to fish. Int. J. Air and Water Poll. 10:49-56.

Mukhopadhyay, M.K. and S.K. Konar (1984) : Toxicity of copper, zinc and iron to fish (*Tilapia mossambica*) plankton (*Cytopsis viridis*) and worm (*Brachium sowerbyi*). Geobios 11(5) : 204-207.

✓ Munshi, J.S.D. (1980): Structure and function of the respiratory organs of air breathing fishes in India, Proc. 67<sup>th</sup> Ind. Sci. Congress, Association.

Murphy, J. and J. Riley (1962): A modified single solution, method for determination of phosphate in natural waters. Anal. Chem. Acta. 27(31-36).

Murthy, K.V., M. Bhaskar and S. Govindappa (1994): Studies on lipid profile of fish liver on acclimation to acidic medium. J. Environ. Biol. 15(4); 269-273.

Murugeson, A.G. and M.A. Haniffa (1992) : Histopathological and histochemical changes in the oocytes of the air-breathing

fish *Heteropneustes fossilis* (Bloch) exposed to textile mill effluent. Bull. Environ. Contam. Toxicol. 48(6): 929-936.

Nanda P. (1997): Nickel toxicity to liver and brain of catfish *Heteropneustes fossilis* (Bloch). Env. Eco. 15(2): 329-331.

Narbonne, J.F. (1973) : Toxicity of lead nitrate in carp metabolism. C.R. Soc. Biol. 4, 167-172.

Natrajan A.V. (1989): Effects of tannery effluent on the muscle and liver glycogen in a fish *Sarotherodon mossambicus*. The Indian Zoologist. 13(1 & 3); 147-151.

Nebeker, V. Alan, C. Savonen and D. G. Steven (1985) : Sensitivity of rainbow trout (*Salmo gairdneri*) early life stage to nickel chloride. Environ. Toxicol. Chem. 4(2) : 233-240.

Neville, C.M. (1979): Sublethal effects of Environmental acidification on a rainbow trout (*Salmo gairdneri*) J. Fish. Res. Bd. Can. 36, 84-87.

Nichol, C.A. and F. Rosen (1963): In: Advances in enzyme regulation (Ed.G.Weber) Vol.I Pergamon Press, Oxford, pp. 341.

Nikam, S.R. (1986) : Studies on the impact of distillery effluent (Spent wash) on two freshwater fishes *Tilapia mossambica* (Peters) and *Rasbora daniconius* (Ham.) M.Phil. dissertation, Shivaji University, Kolhapur (India).

Norin, H., V. Marie, C. Alexandros, S. Margareta (1985) : Concentration of inorganic and total arsenic in fish from industrially polluted water. Chemosphere 14(314) : 325-334.

- Odum, E.P. (1971): Fundamentals of ecology, 3<sup>rd</sup> ed. W.B. Saunders Company, Philadelphia, 574 pp.
- Painter, H.A. (1971) : Chemical, physical and biological characteristic of waste and waste effluent. In : Water and water pollution Handbook (Ed. L.L. Ciaccio) Marcel and Dekker, Inc. New York. 329-360.
- Paka, S. and N. Rao (1997): Inter-relationship of physico-chemical factors of a pond. J. Environ. Biol. 18(1): 67-72.
- Pande, Jyotsna and S.M. Das (1983) : Experimental bioassay of toxic effect of lake water on the fish, *Puntius conchenius*. Proc. Natl. Acad. Sc. India Sect. B. (Biol. Sci.) 53(3) : 169-173.
- Pande, R. and D.N. Saksena (1992) : The toxicity of copper sulphate to the fingerlings of *Labeo rohita* (Ham.). Bull. Env. Sci. Vol. X. P. 23-27.
- Pandey, A.K., S.Z. Siddiqui and K.V. Rama Rao (1993): Physico-chemical and biological characteristics of Hussain sagar, an industrially polluted lake, Hyderabad. Proc. Acad. Environ. Biol. 2(2); 161-167.
- Panigrahi, A.K. and B.N. Misra (1980) : Toxicological effects of a sublethal concentration of inorganic mercury on the freshwater fish. *Tilapia mossambica*. Arch. Toxicol. 44(4) : 269-278.
- Patel, C.B. and G.S. Pandey (1985) : Alumina plant effluents ; appraisal of pollution load. Indian J. Environ. prot, 5(4): 268-270.

- Patil, G.P. and R.R. Dhande (2000): Effect of some heavy metal compounds on testicular recrudescence in the freshwater teleost, *Channa punctatus* (Bloch). J. Aquat. Biol. Vol.15(1&2): 77-80.
- Patil, S.G., D.F. Singh and D.K. Harshey (1986) : Impact of gelatine factory effluent on the water quality and biota of a stream near Jabalpur, M.P. India. J. Environ. Biol., 7(1): 61-65.
- Patra, A.K., S. Patra and S.P. Misra (1990) : Effect of paper mill effluents on hatching and survival of spawns of a major carp *Labeo rohita* (Ham.). J. Freshwater Biol. 2 (4); 323-327.
- Pawar, C.D. (1988) : Studies on fish and fisheries of river Panchganga. M.Phil. dissertation, Shivaji University, Kolhapur (India).
- Perez, C.S., J. Herkovits and A. Salibian (1986): Tetrogenic effects of Cd, on *Bufo arenaratum* during gastrulation. Experimenta. 42:1174-1176.
- Petersen, C. James, S.L. Berks and R.J. Miller (1986) : Sublethal effects of biologically treated petroleum refinery wastewaters on agonistic of male orangespotted sunfish, *Lepomis humills*. Environ. Toxicol. Chem. 5(5) : 463-472.
- Peterson, R.H., P.G. Daye and J.L. Metcalfe (1980): Inhibition of Atlantic Salmon (*Salmo Salar*) hatching at low pH. Can. J. Fish. Aquat. Sci. 37(5): 770-774.

- Prahallad, A. K. and G. Seenaya (1986) : In situ compartmentation and biomagnification of copper and cadmium in industrially polluted Hussain sagar Lake, Hyderabad, India. Arch. Environ. Contam. Toxicol. 15(4) : 417-426.
- Prakasan, V.R., P.S. Johnikutty, P.G. Sanjeev and P. M. George (1989): Combined effect of Hg and Zn in modifying the rate of ciliary activity of *Lamellidens marginalis* J. Environ. Biol. Vol. 10 (2 supp.): 211-215.
- Prakash, D.J. (1996): Effect of detergent, soap and fertilizer on teleost fishes. J. Ecobiol. 8(4): 271-274.
- Pramod, C., Mishra and M. C. Das (1989): Effect of municipal waste on water quality and ecology of Burla lake near Hirakud Dam Reservior. Poll. Res. No.8(3):pp 145-152.
- Radhakrishnaiah, K. (1988) : Accumulation of copper in the organs of freshwater fish. *Labeo rohita* (Ham.) on exposure to lethal and sublethal concentrations of copper. J. Environ. Biol. 9(3) Suppl. 319-326.
- Radhakrishnaiah, K., P. Venkataramana, A. Suresh and B.S Sivaramakrishna (1992): Effect of lethal and sublethal concentration of copper on glycolysis in liver and muscle of the freshwater teleost, *Labeo rohita* (Ham.). J. Environ. Biol. 13(1): 063-068.
- Rajan, M.R. (1990): Sublethal effects of textile mill effluent on protein carbohydrate and lipid content of different tissues of the fish *Cyprinus carpio*. Env. Eco. 8(1): 54-58.



- Rajkumar, R., M. Das and S. Sam (1991): Effect of copper intoxication on muscle glycogen level in *Oreochromis mossambicus*. *Env. Eco.* 9(1); 1-3.
- Ram, R.N. and A.G. Sathyanesan (1984) : Mercuric chloride induced changes in the protein, lipid and cholesterol levels of the liver and ovary of the fish *Channa punctatus*. *Environ. Ecol.* 2, 113-117.
- Ramaswamy, S.N., R.K. Somasekhar and G.D. Artal (1982): Ecological studies on algae in waste from rubber tyre factory, *Ind. J. Environ. Hlth.*, 24(1); 1-7.
- Ranjithakani (1988); A Study on tannery effluent at Erode. National Academy of Sciences, India, 58<sup>th</sup> annual session Jammu Nov. 5-7 (Abstract).
- Rao, K.S., S. Srivastava and S. Dhanekar (1988) : Acute toxicity of reactive textile dyes to eggs and early life history stages of *Cyprinus carpio*. *Geobios.* 15 (2 and 3), 111-113.
- Rao, K.S.P., KRSS Rao, I.K.A. Sahib and KVR Rao (1985); Combined action of carbaryl and phenthoate on tissue lipid derivative of muscle of *Channa punctatus* (Bloch.). *Ecotoxicol. Environ. Saf.* 9(1): 107-111.
- Rathore, V.S., R. Saxena, P.J. John, Rekha (1989) : Acute toxicity of textile dyes to *Channa punctatus*. *J. Nature Conserv.* 1(2), 155-157.
- Ray, P.K. (1986); Environmental Protection and Cancer. *J. Sci. & Industrial Research* 45: 370-371.

- Reusink, R.G. and L.L. Smith (1975): Relationship of 96 h LC<sub>50</sub> to lethal threshold concentration of hexavalent chromium, phenol and sodium pentachlorophenate for fat head minnow, *Pimephales promelas* (Raf.). Trans. Amer. Fish. Soc. Vol.104:567.
- Robbert, G.S., J. Grindley and E.M. William (1940): Chemical methods for the study of river pollution, London. H.M.S.O. Fish Invert. Ser. I 4(2).
- Roe, J.R. and KVR Rao (1981): Lipid derivatives in the tissue of freshwater teleost *Sarotherodon mossambicus* (Peters): Effects of methyl parathion. Proc. Ind. Natl. Sci. Acad. 47:53-57.
- Rosenthal, H. and K.R. Sperling (1974): Effect of cadmium on development and survival of herring eggs. In: The early life history of fish (Ed. J.H.S. Blexter) Berlin Springer pp. 383-396.
- Sakthivel, M. (1989) : Toxic effect of tannery and textile mill effluents on the fishes *Cyprinus carpio* and *Oreochromis mossambicus*. Env. Eco. 7(8), 685-689.
- Sampath, K. and Sakthivel (1989) : Effects of textile dye stuff effluent on food utilization of *Cyprinus carpio*. Indian J. Exptl. Biol. 27(12), 1032-34.
- Sarkar, S.K. (1991b): Effects of temperature on eggs, fry and fingerlings of *Labeo rohita* exposed to urea Prog. Fish. Cult. 53; 242-245.

- Sarkar, S.K. (1995); Influence of temperature on the toxicity of nitrogen and phosphorous fertilizer of fish and aquatic organisms. Proc. Acad. Environ. Biol. 4(1): 63-75.
- Sarkar, S.K. (1997) : Influence of temperature on the feeding rate of fish exposed to some agricultural chemicals. J. Environ. Biol 18(3); 283-286.
- Sastry, C.A. and R. Kamatchiammal (1988) : Effect of paper mill effluents on fish life. Indian J. Environ. Prot. 8(1): 31-38.
- Sastry, K.V. and K. Sunitha (1984) : Chronic toxic effect of chromium in *Channa punctatus* : Biochemical studies. J. Environ. Biol. Vol. 5 (1) : 53-56.
- Sastry, K.V. and V. Shukla (1993): Effect of cadmium on the rate of oxygen uptake by the fish *Channa paunctatus*. J. Ecobiol. 5(4): 295-298.
- Sastry, K.V. and V. Shukla (1994) : Acute and chronic effects of cadmium on some parameters in the freshwater teleost fish *Channa punctatus*. Acta. Hydrochim. Hydrobiol. 22(4), 171-176.
- Saxena, M.M. (1998): Environmental Analysis Water, Soil and Air. Published by Agro Botanica Bikaner.
- Shaffi, S.A. (1978) : Cadmium intoxication on tissue glycogen content in three freshwater teleost. Curr. Sci. 47, 868-870.
- Shaffi, S.A. (1980) : The acute industrial effluent toxicity of freshwater fish. Toxicol. Lett. (AMST) 5 (3/4) : 183-190.

- Sharma, A. and M. Sharma (1994) : Toxic effect of zinc smelter effluent to some developmental stages of freshwater fish, *Cyprinus carpio* (Linn.). J. Environ. Biol. 15(3), 221-229.
- Sharma A. and M.S. Sharma (1995) : Acute toxicity of zinc to certain developmental stages of *Cirrhinus mrigala* (Ham.). J. Environ. Biol. 16(2), 157-162.
- Sharma, K.C. (1984) : Effect of mercury pollution on the general biology and carbohydrate metabolism of a freshwater murrel *Channa punctatus*. Geobios, 11(3) : 122-127.
- Shivaraj, K.M. and H.S. Patil (1988) : Toxicity of cadmium and copper to a freshwater fish. Environ. Ecol. 6, 5-8.
- Shivraj, K.M. and H.S. Patil (1985) : Acute toxicity of cadmium and cobalt to a freshwater fish *Lepidocephalichthys guntea*. Indian J. Comp. Anim. Physiol. 3(2): 24-28.
- Shrivastava, A.K. (1989) : Pollution of the Faren stream by the effluent of sugar factory and distillery. Adv. Plant Sci. 2(2): 308-310.
- Shrivastava, V.S. (1996) : Heavy metals in industrial wastewater and sludge by ICP-atomic emission spectrophotometer. Indian J. Environ. Prot. 15(5) : 328-329.
- Shumway, D.F. and J.R. Palensky (1973) : Impairment of flavour of fish by water pollutants. U.S. Environ. Prot. Agency, EPA R3, 73-10.
- Sijm, D.T.H.M., P. Peart and A. Opperhuizen (1993) : The influence of temperature on the uptake rate constant of

hydrophobic compounds determined by the isolated perfused gills of rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.* 25(1-2) : 1-14.

Singh, C., H.K. Parwana, S.S. Marwaha, R. Garg and Gajendra. Singh (1996): Toxicity of electroplating effluents. *Journal of Industrial pollution control* 12(1): 15-19.

Singh, R.K. and S.P. Singh (1996) : Effect of industrial effluent on the food fishes of river Sone. *Proceeding of 83<sup>rd</sup> session of the Indian Science Congress, Patiala.*

Singhal, K.C. (1994): Biochemical and enzymatic alterations due to chronic lead exposure in the fresh water catfish, *Heteropneustes fossilis*. *J. Environ. Biol.* 15(3): 185-191.

Skidmor, J.F. (1964): Toxicity of zinc compound to aquatic animals with special reference to fish. *Q. Rev. Biol.* 39: 224-248.

Skidmor, T.F. and P.W.A. Tovell (1972): Toxic effects of zinc sulphate on the gills of Rainbow trout-*Wat Res.* 6: 217-230.

Smith, R. Leverett, T.M. Holsen, C.I.B. Noel, R.M. Block and A.B. DeLeon (1985) : Studies on acute toxicity of fluoride ions to stickle back (*Gasterosteus aculeatus*), fat head minnow (*Pimephales promelas*) and rainbow trout (*Salmo gairdneri*). *Chemosphere.* 14(9) : 1383-1390.

Somanath, B. (1991): Effect of acute sublethal concentration of tannic acid on the protein, carbohydrate and lipid levels in the tissue of the fish *Labeo rohita*. *J. Environ. Biol.* 12(2): 107-112.

- Somasundara, B., P.E. King and S.E. Shackley (1984): Some morphological effects of zinc upon the yolk-sac larvae of *Clupea harengus*. J. Fish. Biol. 25(3): 333-344.
- Spear, P.A. and R.C. Pierce (1979): Copper in the aquatic environment: Chemistry, distribution and toxicology. Natl. Res. Counc. Can. Assoc. Comm. Sci. Criter Environ. Qual. Publ. A (16454): 13-184.
- Spear, P.A.(1981): Zinc in aquatic environment: Chemistry distribution and toxicology and Environ. Quant. Publ. 0(175769); 1-145.
- Sprague, J.B. and D.W. Melease (1968) : Different toxic mechanism in kraft pulp mill effluent for two aquatic animals. Water Res. 2, 761-765.
- Srinivas, S.V. and M.B.P. Rao (2000): Impact of Hexavalent chromium on survival of the freshwater fish *Sarotherodon mossambicus*. J. Aquat. Biol. Vol. 15(1&2): 71-73.
- Srinivas, V. and M.B.P. Rao (1996) : Acute toxicity of chromium to the freshwater teleost fish *Labeo rohita* (Ham.). Indian J. Comp. Anim. Physiol. 14-2, 30-32.
- Srinivas, V. and M.B.P. Rao (1998) : Chromium induced histological alteration in the gill of the freshwater teleost fish, *Labeo rohita* (Ham.). Indian J. Comp. Anim. Physiol. Vol. 16(1), 54-62.
- Srivastava, D.K. (1982) : Comparative effects of copper, cadmium and mercury on tissue glycogen of catfish. *Heteropneustes fossilis*. Toxicol. Lett. 11, 135-140.



- Stephen, M., R. S. Antony, A.G. Murugesan and M.A. Hanifa (1987): Toxicity of industrial effluents to fresh water cat fish *Mystus kelitius* Curr. Sci. 56, 733-734.
- Strain, R. (1984) : New directions of future water quality legislation, speciation of heavy metals. National Symp. On Assessment of Env. Polln. At Aurangabad.
- Subramanian, M.A. and G. Varadaraj (1993) ; Impact of Tannery effluent on the biochemical constituents in the haemolymph of the freshwater prawn, *Macrobrachium idella* Heller. J. Environ. Biol. 14(3); 255-259.
- Subramanian, M.A., S. Vishwanathan, G. Varadaraj (1988): A report on the physico-chemical parameters of the sewage and industrial effluents in and around Erode, Tamil Nadu. The Indian Zoologist. 12(1 & 2) ; 185-187.
- Sunila, I. (1981) : Toxicology of copper and cadmium to *Mytilus edulis* L. (Bivalvia) in brackish water. Ann. Zool. Fennici, 18, 213-223.
- Surendrakumar, A. Prasad and J. Prasad (1996): Studies on biochemical composition in relation to body weight in freshwater major carp, *Labeo rohita* (hom.). Proceeding of the 83<sup>rd</sup> session of the Indian science congress, Patiala.
- Suresh, I.V. and A. Wanganeo (1996): Municipal sewage sludge- Heavy metals. Proc. Acad. Environ. Biol. 5(1): 65-70.
- Tam, W.H., L. Birkett, R. Makaran, P.P. Payson, D.K. Whitney, CK Yu (1987) : Modification of carbohydrate metabolism and liver vitellogenic function in brook trout (*Salvelinus*

- fontanalis*) by exposure to low pH. Can. J. Fish. Aquat. Sci. 44(3) : 630-635.
- Tana, J. and E. Nikunen. (1986) : Physiological responses of rainbow trout (*Salmo gairdneri*) in a pulp and paper mill recipient during four seasons. Ecotoxcol. Environ. Saf. 12(1); 22-34.
- Tariq, J., M. Ashraf, M. Jaffar and M. Afjal (1996): Pollution status of the Indus river, Pakistan through heavy metal and macronutrient content of fish, sediment and water. Water Research, 30(6):1337-1344.
- Taylor, D., B.G. Maddock and G. Mance (1985) : The acute toxicity of nine 'gray list' metals (Ar, Bo, Cr, Cu, Pb, Ni, Sb, Va and Zn) to two marine fishes : DAB (*Limanda limanda*) and gray mullet (*Chelon labrosus*). Aquat. Toxicol. (AMST) 7(3) : 135-144.
- Thakre, R., D. Ramteke and V.P. Thergaonkar (1986) : Impact of thermal power plant emissions and effluents on water quality. J. Indian Water Assoc. 18 (1): 39-47.
- Thingran, V.G. (1974) : Fisheries in India in the context of aquatic pollution. Int. J. Ecol. Environ. Sci. 1, 15-18.
- Thorat, S.R. and S.B. Wagh (1998); Study on physico-chemical parameters of Tannery effluent at Aurangabad. National Symposium on Environmental issues and strategies for sustainable development. No. 2.36. Jan. 23-25 Aurangabad.



- Tort, L., R. Flos and J. Balasch (1984) : Metabolic levels in dogfish (*Scyliorhinus canicula*) gill tissue after zinc treatment. Rev. Esp. Fisiol. 40(2). 149-152.
- Tort, Lluís, P. Torres and R. Flos (1987) : Effects on dogfish hematology and liver composition after acute copper exposure. Comp. Biochem. Physiol. C. Comp. Pharmacol. Toxicol. 87 (2):349-354.
- Trivedy, R.K., S.D. Khatavkar and P.K. Goel (1986); Characterization, treatment and disposal of water in a textile industry. J. Ind. Pol. Cont. 2(1): 1-2.
- Trivedy, R.M., P.K. Goel and C.L. Trisal (1987): Practical Methods In Ecology And Environmental Science. Published by Enviromedia, Karad.
- Trotman, S.R. and E.R. Trotman (1984): The Bleaching, Dyeing chemical Technology of textile Fibres. Charles Griffin and Comp. Ltd., London.
- Valley, B.L. and D.D. Ulmer (1972): Biochemical effect of Hg, Cd and Pb. Annual Res. Biochem. 41: 91-128.
- Varadaraj, G., M.A. Subramanian and B. Nagarajan (1993) : The effect of sublethal concentration of paper and pulp mill effluent on the haematological parameters of *Oreochromis mossambicus* (Peters). J. Environ. Biol. 14(4), 321-325.
- Varadraj, G., S. Jayasuria and M.A. Subramanian (1994): Toxic effect of tannery effluent on the biochemical constituents in different tissues of *Pila globosa*. Env. Eco. 12(2); 303-307.

- Venkataramana, P. and K. Radhakrishnaiah (1987) : Lethal and sublethal effects of copper on the protein metabolism of the freshwater teleost *Labeo rohita* (Ham.). Trends Life Sci. 2(2) : 82-86.
- Verma, S.R. and R.C. Dalela (1975) : Studies on the pollution of Kalinadi by industrial waste near Mansurpur part. Biological index of pollution and biological characteristics of river. Acta Hydrochim. Hydrobiol. 3, 259-274.
- Verma, S.R. and R.P. Mathur (1974) : Studies on toxicity of Industrial waste to *Macrobrachium dyanum*. Indian J. Environ. Hlth. 16, 1-11.
- Verma, S.R., P. Sharma,, A. Tyagi,, S. Rani,, A.K. Gupta and R.C. Dalela (1984): Pollution and saprobic status of Eastern Kalinadi Linnologica (Berlin) 15(1): 69-133.
- Victor, B., S. Mahalingam and R. Sarojini (1986) : Toxicity of mercury and cadmium on oocyte differentiation and vitellogenesis of the teleost, *Lepidocephalichthys thermalis* (Bleeker). J. Environ. Biol. 7 (4) : 209-214.
- Vijaya Lakshmi, G. Umadevi, M. Prasad, M.C. Venkatasubbaiah and S. Govindappa (1990): Effect of environmental acidity on tissue metabolism of fresh water fish, *Cyprinus carpio* (Linn.) I. Muscle metabolism. J. Environ. Biol. 11(3): 269-274.
- Virk, S. and R.C. Sharma (1995) : Effect of nickel and chromium on various life stages of *Cyprinus carpio* (Linn.). Indian J. Eco., 22(2), 77-81.

- Viswaranjan, S. and J. Muthukrishnan (1982) : Toxicity of tannery effluent to mosquito *Culex pipiens quinquefasciatus*. Proceedings of III Oriental Entomology Symposium, University of Kerala Trivendrum, 3, 29-34.
- Viswaranjan, S. and J. Muthukrishnan (1982) : Toxicity of tannery effluent to mosquito *Culex pipiens quinque fasciatus*. Proceedings of III Oriental Entomology Symposium University of Kerala Trivendrum, 3, 29-34.
- Von Hoof, F., J.P. Nauwelaers and N.V. M. Seslenmeg (1984): Distribution of nickel in the Roach, *Rutilus rutilus* Chemosphere 13(9): 1053-1059.
- Von, Westernhagon, H. and V. Dethlefsen (1975) : Combined effect of cadmium and salinity on development and survival of flounder eggs. J. Mar. Biol. Ass. U.K. 55, 945-957.
- Wagh, S.B. (1986) : Seasonal variation in the biochemical composition of freshwater Cyprinid fish *Chela clupeoides* : Part I. Muscle. Indian J. Comp. Animal Physiol. 4(1) : 49-54.
- Wagh, S.B., K. Sharif and S. Shaikh (1985) : Acute toxicity of cadmium sulphate, zinc sulphate and copper sulphate on *Barbus ticto*, oxygen consumption and gill histology. J. Environ. Biol., 6(4) : 287-293.
- Waiwood, K.G. and F.W. Beamish (1978): Effects of copper, pH and hardness on the critical swimming performance of rainbow trout *Salmo gardneri* Wat. Res. Vol. 67 No.B 12, 611-620.

- Walsh, D.F., L. Bernard, Berger and R. Jerry (1977); Mercury, arsenic, lead, cadmium and selenium residue in fish. *Pestic. Monit. J.*, 11(1): 5-34.
- WHO (1977): *Environ: Health Criteria – 3 lead* WHO Geneva.
- Wilson, D., B. Finlayson and N. Morgan (1981): Copper, zinc and cadmium concentrations of resident trout related to acid mine wastes. *Calif Fish. Game.* 67(3): 176-186.
- Wong, M.H., K.C. Luk and K.Y. Choi (1977): Effect of zinc and copper salts on *Cyprinus Carpio* and *Ctenopharyngodon idellus*. *Acta. Anat.*, 99(4): 450-454.
- Woo, P.T.K. , Y.M. sin and M.K. Wong (1993) : The effect of short term acute cadmium exposure on blue tilapia *Oreochromis aureus*. *Environ. Biol. Fish.* 37(1) : 67-74.
- Yamagata, Y.M. and Niwa (1982) : Acute and chronic toxicity of ammonia to eel *Anguilla japonica*. *Bull. Jap. Soc. Sci. fish* 49(2) : 171-176.
- Yamamoto, Y,T., Ishii and S. Ikeda (1978): Studies on copper metabolism in fishes-III. Existence of metallothionein like protein in carp hepatopancreas. *Bull. Jap. Soc. Sci. Fisheries* 44: 149-153.