

## REFERENCES

=====

- 1) L. Benson, 1962, Plant Classification, 'Family Leguminosae - Pea Family', P. 244.
- 2) T. R. Narayan and P. M. Dabaghao, 1972, Forage Crops of India, Medics (*Medicago* species), pp. 54-61.
- 3) M. S. Tacker (Ed) and Associating Editors, 1962. 'Medicago Sativa', The Wealth of India : A Dictionary of Indian Raw Materials & Industrial Products, (CSIR), pp 316-317.
- 4) E. Bottini, 1949, Vitamins as Factors of Quality in Forage Plants, Ann. Sper. Agar. (Rome) [N.S.] 3, pp 1167-79.
- 5) R. L. Horst, T. A. Reinhardt, J. R. Russell and Napoli, 1984, The Isolation & Identification of vitamin D<sub>2</sub> and Vitamin D<sub>3</sub> from Medicago Sativa, 'Arch. Biochem. Biophys', 231 (1), 67-71.
- 6) M. A. Sahadev, P. S. Rao, S. Deolakar and K. V. Giri, 1949, Vitamin C Content of Lucerne & Factors Controlling its Stability.
- 7) E. M. Bickoff, A. I. Livingston and N. S. Snell, 1950, 'Occurance of Vitamin B<sub>12</sub> and other Growth in Alfalfa', Arch. Biochem., 28, pp 242-52.

- 8) A. Atom, and A. Monzini, 1952, 'The Vitamin B<sub>2</sub> Content of Vegetable fruits, Fodders & Feedstuff', Ann. Sper. Agar (Rome), 6, pp 1021-26.
- 9) M. S. Tacker (Ed) and Associating Editors, 1962. 'Medicago Sativa', The Wealth of India : A Dictionary of Indian Raw Materials & Industrial Products (CSIR), p318.
- 10) R. Douillard, 1986, 'Ribulose Biphosphate Carboxylase [EC 4.1.1.39] from Alfalfa Leaves', Sci. Aliments., 6(1), pp 81-89.
- 11) H. Cynthia, A. Stanley, H. Duke and M. Collins, 1986, 'NADP<sup>+</sup> -isocitrate Dehydrogenase [E.C.1.1.1.42] Purified from the Host Plant Cytosol of Medicago L. av.', Physiol. Plant, 67(4), pp 538-564. sativa
- 12) A. E. Timbekova, 1986, 'Two Triterpene Glucosides Isolated from Roots of Medicago Sativa, Caulosaponin B and a New Glycoside Medicoside C.', Khim. Prir. Soedin (Tashk), 0(6), pp. 805 -8.
- 13) M. Paul, M. Dewick, and M. Martin, 'Biosynthesis of pterocarpan & Isoflavan Phytolexin in Medicago Sativa : The Biochemical Interconversion of Pterocarpans and 2-Hydroxyisoflavans', Phytochemistry, 18, pp 591-96.
- 14) D. A. Bender, 1975, 'Amino Acid Metabolism : Amino Acids Synthesised from Glutamate : Proline, Ornithine & Arginine', pp 80-97.

- 15) ibid.
- 16) D. R. Morris, and A. B. Pardee, 1966. 'Multiple Pathways of Putrescine Biosynthesis in E. Coli', J. Biol. Chem., 241, pp 3129-35.
- 17) T. A. Smith, 1969, 'Biosynthesis of Putrescine in Maize'.
- 18) A. Maretzki, M. Thom and L. G. Mickell, 1969, The Biosynthesis of Polyamines in Sugarcane, Phytochemistry, 8, pp 811-972.
- 19) H. Tabor and C. W. Tabor, 1972, 'Biosynthesis and Metabolism of 1,4-diaminobutane, Spermidine, Spermine & Related Amines,' Adv. Enzymol., 36, pp 203-69.
- 20) B. N. Ames and D. T. Dubin, 1960, 'The Role of Polyamines in the function of Bacteriophage DNA', J. Biol. Chem., 235, pp 769-775.
- 21) J. Miersch and H. Reinbothe, 1966, 'Chromatographic Separation of Amino Acids and Guanidine Compounds from Fruit Bodies of Higher Fungi', Flora, 156, p. 543.
- 22) E.F. Gale, 1971, 'Metabolism of Fruit Bodies of Fungus Panus Tigrinus', Phytochemistry, 10, p. 735.
- 23) A. Meuberger, 1981, Special Problem in Nitrogen Metabolism in Plants, Comprehensive Biochemistry, 19A : Amino Acid Metabolism & Sulphur Metabolism, pp. 135-140.

- 24) D. J. Durzan, 1968, 'Nitrogen Metabolism of Picea glauca; Diurnal changes of free Amino Acids, Amides & Guanidino Compounds in Roots, Buds and Leaves during the Onset of Dormancy of White Spruce Saplings', Can. J. Biochem., 47, p. 921.
- 25) A. Uhlemann, H. Reinbothe, 1977, 'Characterization of L-Arginine Carboxylase of Panus tigrinus', Biochem. Physiol., Pflanzen, 171, p. 85.
- 26) J. Durzan, 1968, 'Nitrogen Metabolism of Picea glauca: Seasonal Changes of Free Amino Acids in Buds, Shoot apices and Leaves & the Metabolism of Uniformly labelled L-Arginine  $^{14}\text{C}$  by buds during the Onset of Dormancy', Can. J. Bot., 46, p. 909.
- 27) Thoai, V. Mauyen and J. Roche, 1960, 'Biological Guanidine Derivatives', Fortschr. Chem. Organ. Naturstoffe, 18, p. 83.
- 28) E. Muller, and K. Armbrust, 1940, 'Nitrogenous Component of Soybeans', 2, Physiol. Chem., 263, p. 41
- 29) Shingo, Mitsui and K. Kumazawa, 1961, 'Nutrient Uptake by crop plants, Amino acids and other Nitrogenous Compounds in rice root under Different Nutritional Conditions' J. Sci. Soil Manure (Tokyo), 32, p. 57.
- 30) J. B. Walker, P. A. Hoffee and M. E. Jones (Ed), 1975, 'Methods in Enzymology', Vol. 43, p. 429.

- 31) A. Neuberger, 1981, 'Special Problem of Nitrogen Metabolism in Plants', Comprehensive Biochem., 19A : Amino Acid and Sulphur Metabolism. pp. 135-140.
- 32) G. Reuter, A. Barthel, 1967, 'Creatine as a Plant Product', Pharmazie, 22, p. 261.
- 33) J. A. Lippincott and B. B. Lippincott, 1975, 'Induction of Responsiveness to Octopine and Lysopine in Crown gall Tissues', Ann. Rev. Microbiol. 29, 377
- 34) A. Menage and G. Moret, 1965, 'Presence of Amino acids in Crown gall tissues', Acad. Sci., 261(1965), p. 2001
- 35) M. F. Wendt, Gallitelli and Dobrigkeit, 1973, 'Octopine in Pea Seedlings', 2, Naturforch, 28C, p. 768
- 36) Petit, Annick and G. Morel, 1966, 'Metabolism of Homoarginine by crown gall Tissues', C.R. Soc. Biol. 160, p. 806.
- 37) G. S. Cristie, N. P. Madsen, M. p. Hegarty, 1970, 'Acute Biochemical Changes in rat liver induced by the naturally occurring amino acid Indospicine', Biochem. Pharmacol, 19, p. 853.
- 38) G. Reuter, A. Barthel, 1967, 'Synthesis of radioactive labelled compounds of Guanidine acetic acid- Amidine  $^{14}\text{C}$  and Guanidine acetic acid - 2- $^{14}\text{C}$ - Amidine  $^{14}\text{C}$ ', Pharmazia, 22, p. 261.

- 39) G. Reuer, A. Barthel, 1967, 'Biochemistry and Physiology of Isoprenoid Guanidine especially (L-Hydroxy - 3 - methyl -2-buten(yl), guanidine in Galega officinalis', Pharmazie, 23, p. 261.
- 40) A. petit, J. Tempe, G. Morel, 1968, 'Presence of Conavanine Metabolite in Canavalia ensiformis Tumor Tissue', C. R. Soc. Biol. 162, p. 632.
- 41) I. Krone and G. Reuter, 1968, 'Metabolism of Spherophysine (4-(3-methyl-2-butenyl amino butyl)guanidine) in Sphaerophysa salsula', Dtsch. Pharm. Ges., 301, p. 64.
- 42) Y. Fugita, 1959, 'Hydroxy arginine, guanidino compound from a sea cucumber isolation and identification', Bull. Chem. Soc. Japan, 32, pp. 439 - 442.
- 43) G. E. Wolstenholme and M. P. Cameron, 1965, Comparative Biochem. of Arginine and Derivatives, London, p. 20..
- 44) J. Cabello, V. Prajoux and M. Plaza, 1966, 'Separation of two enzymes in purified preparation of human liver and erythrocyte arginase by zone electrophoresis mobility of the fast component in both preparation', Arch. Med. Exptl. 3(1), pp. 7 - 13.

- 45) G. Brabdner, and A. I. Virtanen, 1964. ' $\alpha$ - oxo - $\delta$ - guanidinovaleric acid and  $\gamma$ - hydroxy- $\alpha$ - oxoglutaric acid in Green Parts & Seeds of Phloex Plants, *Acta. Chem. Scand.*, 18, p. 574.
- 46) A. E. Bender, H. A. Krebs, 1950, 'The Oxidation of Various Synthetic  $\alpha$  - Amino Acids by Mammalian D-amino acid oxidase, L-Amino Acid Oxidase of Cobra Venom & the L & D-Amino Acid Oxidases of Neurospora Crassa', *The Biochemical Journal*, vol. 46, p. 210-19.
- 47) K. Burton, 1952, ' The l-Amino Acid Oxidase of Neurospora ', *Biochemical Journal*, 50, p. 258.
- 48) L. K. Ramachandran, K. Rao, 1963, 'Preparation of & Properties of Lathyrine', *Biochem. Biophys. Res., Commun.*, 13, p. 49.
- 49) ibid, 2, p. 298.
- 50) E. A. Bell, G. Fooster, 1962, 'Structure of Lathyrin, *Nature*, 91(2), 1962.
- 51) G. A. Rosenthal, 1982, 'Plant Nonprotein Amino & Imino Acids, Biological, Biocem. & Toxicological Properties, Nomenclature & Certain Physicochemical Properties, pp. 1-2.
- 52) P. M. Dunill and L. Fowden, 1965, 'The Amino Acids of Seeds of the Cucurbitaceae, *Phytochemistry*, 4, pp. 933-44.

- 53) D. O. Gray & L. Fowden, 1961, 'N-ethyl-l-asparagine : A New Amino Acid Amide from Ecballium', Nature, 189, pp. 401-401.
- 54) L. Fowden, 1961, 'A New Asparagine Derivative N<sup>4</sup>-(2-hydroxyethyl)-l-asparagine from bryony (Bryonia dioica), Biochem. J. 81, pp. 154-156.
- 55) A. I. Virtanen, 'Studies on Organic Sulfur Compounds & Other Labile Substances in Plants, Phytochemistry, 4, pp. 207-28.
- 56) L. Fowden, 1955, 'Azetidine-2-Carboxylic Acid : A New Constituent of Plants', Nature, 176, pp. 347-8.
- 58) L. Fowden, et al., 1969, 'Amino Acids', Phytochemistry, (Ed. Lawrence P. Miller), p. 5.
- 57) R. Gmelin and P. O. Larsen, 1967, L-γ-Ethylidene glutamic acid & L-γ-Ethylene glutamic acid in seeds of Tetrapleura tetraptera, Bichem. Biophys. Acts., 136, pp. 572-3.
- 59) A. I. Virtanen and P. K. Hietala, 1955, -Hydroxy Glutamic Acid in Green Plants, Acta. Chem. Scand., 9, pp. 175-6.
- 60) J. Done, and L. Fowden, 1952, 'A New Amino Acid, Amide in the Ground Plants (Arachis hypogaea) : 'Evidence of the occurrence of γ-methylene glutamine and γ-methylene glutamic acid', Biochem. J., 51, pp. 451 - 458.

- 61) A. I. Virtanen and T. Ettala, 1957, 'Dihydroxy Glutamic Acid in Plants', Acta. Chem. Scand., 11, pp. 182 - 4.
- 62) L. Fouwden, 1966, 'The Acidic Amino Acids of Tulip: Isolation of  $\gamma$ - ethyldene -Glutamic Acid', Biochem. J., p. 98
- 63) R. Gnelin and P. O. Larsen, 1967, 'L- - ethyldone Glutamic Acid and L -  $\gamma$  - Methylene Glutamic Acid in seeds of Tetrapleura tetraptera', Biochem. Biophys. Acta., 136, pp. 572 - 3.
- 64) E. A. Dell, 1964c, 'Relevance of biochemical taxonomy to the problem of Lathyrism', Nature, 203, pp. 378-80
- 65) M. P. Hegarty and A. W. Pound, 1968, Indospicine, a New Hepato toxic Amino Acid from Indigofera spicata', Nature, 217, pp. 354 - 5.
- 66) E. Bell and A. S. L. Tirimanna, 1965, 'Association of Amino Acids and Related compounds in seeds of forty seven species of Vicia: their taxonomic and nutritional significance', Biochem. J. 97, pp. 104 11
- 67) T. Takemoto, K. Daigo and N. Takagi, 1964, 'Hypotensive Constituents of marine algae ; A new basic Amino Acid Laminine', Yakugaku Zasshi, 84, pp. 1176 - 9.



- 68) K. Biemann, C. Lionet, J. Asselineau, E. Lederer, and J. Polomsky, 1960, 'On the Structure of Lysopine, A New Amino Acid Isolated from Crown-gall Tissue, Biochem. Biophys. Acta., 40, pp. 369-70.
- 69) B. L. Turner and J. B. Harborne, 1967, 'Distribution of Canavanine in the Plant Kingdom', Phytochemistry, 6, pp. 836-6.
- 70) E. A. Bell, and R. G. Foster, 1962, 'Structure of Lathyrine', Nature, 194, pp. 91-2.
- 71) L. F. Burroughs, 1957, '2-aminocyclopropane-1-Carboxylic Acid : A New Amino Acid in Perry Pears & Cider Apples', Nature, 179, pp. 360-1.
- 72) L. Fowden and A. Smith, 1968, 'Newly Characterised Amino Acids from Aesculus Californica', Phytochemistry, 7, pp. 809-19.
- 73) L. Fowden, 1964, 'The Chemistry and Metabolism of Recently Isolated Amino Acids', Ann. Rev. Biochem., 33, pp. 451-5.
- 74) M. Takagi, K. Oishi, A. Okumura, 1967, 'Free Amino Acid Composition of Some Species of Marine Algae', Bull. Japan Soc. Sci. Fish., 33, pp. 669-73.
- 75) J. F. Fisher, M. F. Mallette, 1961, 'The Natural Occurance of Ethionine in Bacteria', J. Gen. Physiol., 45, pp. 1-13.

- 76) L. Fowden, 1955, Azetidine-2-Carboxylic Acid : A New Constituent of Plants', Nature, 176, pp. 347-8.
- 77) L. Fowden and F. C. Steward, 1957, 'Nitrogenous Compounds & Nitrogen Metabolism in the Liliaceae', Ann. Bot. N. S., 21, pp. 53-67.
- 78) H. Tomita, S. Mizusaki and E. Tamaki, 1964, Chemical Studies on Ninhydrin Positive Compounds in Cured Tobacco Leaves', Agric and Biol. Chem., 28, pp. 451-5.
- 79) A. N. Radhakrishnan and K. V. Giri, 1954, 'The Isolation of Allohydroxy-L-Proline from Sandal (Santalum album L.) Biochem. J., 58, pp. 57-61.
- 80) M. L. Sung and L. Fowden, 1968, 'Trans-3-hydroxy-L-proline : A Constituent of Delonix regia., Phytochemistry, 7, 2061-3, p. 889.
- 81) M. Guggenheim, 1958, 'Agmatine Metabolism, Handbook of Plant Physiology, Vol. III
- 82) J. W. Clark Lewis and P. I. Mortimer, 1961, The 4-hydroxypipeolic Acid from Acacia Species & its Stereoisomers', J. Chem. Soc., pp. 189-96.
- 83) M. D. Wilding and H. W. Schnabel, 1962, 'Hydroxylysine from Alfalfa Roots', Phytochemistry, 1, 263-5.
- 84) J. Przebylska and J. Pawelkiewicz, 1965, 'O-oxallyl-homoscrine, A New Homoserine Derivative in Young pools of Lathyrus sativa', Bull. Acad. Pol. Sci. Ser. Sci. Bio., 13 pp. 327-29.

- 85) J. K. Pollard, E. Sondheimer and F. C. Steward, 1958,  
'New Hydroxyamino Acids in Plants & Their  
Identification', Nature, 182, pp. 1356-59.
- 86) L. Fowden, 1973, 'Amino Acids', Phytochemistry, p. 15.  
L. Fowden, and A. Simth, Op. cit.
- 88) R. R. Doyle and B. Levenberg, 1967, 'Structure of  
Several Unusual  $\alpha$ - Amino Acids from Basidiomycetes',  
Fed. Proc., 26, p. 45.
- 89) V. V. S. Murti, T. R. Seshadri and Venkitasubramanian,  
1964, 'Physiological Importance of Nonprotein Amino  
Acids', Photochemistry, 3, p. 73.
- 90) M. P. Hegarty, P. G. Schinckel and R. D. Court, 1964,  
'Reaction of Sheep to the Consumption of Leucaena  
gluca, Benth. and Its Toxic Principle Mimosine, Aust.  
J. Agric. Res., 15, pp. 153-67.
- 91) L. Fowden, 1963, 'Amino acid Analogues and the Growth  
of Seedlings', J. Exptl. Bot., 14, pp. 387 - 98.
- 92) L. Fowden, 1967, 'Aspects of Amino Acid Metabolism in  
Plants', Ann. Rev. Plant Physiol., 18, pp. 85 -106.
- 93) E. Bege, 1978, 'Plant Ecology', Plant Res. Develop,  
7, p. 14.
- 94) C. S. Levenworth, A.J. Wakeman and T. B. Osborne,  
1924, J. Biol. Chem. 24, viii, p. 209, 'The  
nitrogenous constituent of the juice of the Alfalfa  
Plant'.

- 95) R. L. M. Synge, 1951, 'Non protein Nitrogenous constituent of Rye Grass': Biochem. J. 49, p. 642
- 96) G. Gomori, 'Preparation of Buffers for use in Enzyme Studies', Methods in Enzymology, Vol. I., pp. 138-143
- 97) S. Moore and W. Stein, 1954, 'Determination of Amino Acid by Ninhydrin Method', J. Biol. Chem. 211, p. 893
- 99) A. Harden and D. Morris, 1911, 'The Diacetyl C - naphthol reaction of Arginine', J. Physiol., 42, p. 332.
- 100) T. Shinzo, O. Takeshi, S. Takeichi, 1975, 'New Colurimetric method for guanidines and mono substituted Guanidines with 9, 10 -Phenanthraquinoline and 3,5-dihydrxy benzoic acid', 23, (8), pp. 1657 63.
- 101) F. F. Ventimigli, D. J. Durzan, 1986, 'Determination of Mono Substituted Guanidines using a dedicated amino acid analyser', LC -GC, 4(11), pp. 112 - 24.
- 98) P. N. Warness and J. F. A. McManus, 1951, 'Sakaguchi Oxine Test for Arginine applied to the Tissue section Exptl. Cell Res., 2, pp. 703-7.
- 102) R. Yomasaki, S. Bryan, A. Dorothy, R. E. Feeney, 1981, 'Colourimetric determination of Arginine in Residues in Proteins by Nitro phenyl Glyoxal.
- 103) V. V. Pedro, 1980, 'Enzymatic Immobilization for Determining Arginine Level in Biological Products', Diss. Abstr. Int. B., 41(5), p. 1706.

- 104) M. Peterkova, O. Matousova, B. Kakae, 1980, 'Use of Guanidine Group for the Determination of the Drug 2, 2, 5, 7, 8-pentamethyl-6(2-guanidino ethoxy chroman', VYZK. Ustav Farm, Biochem. Prague(Zesk) Farm, 29 (5), pp. 125 - 7.
- 105) J. Jayaraman, 1981, 'Carboxyhydrates', Laboratory Manuel in Biochemistry, p. 78.
- 106) W. W. Pigman, M. L. Wolfrom (Ed), 'Amino Sugar', Advances in carbohydrate Chemistry, Vol.2, 1946, p. 170.
- 107) O. H. Lowry, N. J. Rosebrough, A. G. Farv and R. J. Randall, 1951, 'Protein Estimation by Folin Phenol', J. Biol. Chem. 193, p. 265.
- 108) Hawk, 'Starch', Hawk's Physiological Chem. pp. 94-96.
- 109) T. J. Schoch, W. W. Pigman, M. L. Wolfrom (Ed), 'The Fractionation of Starch', Advances in Carbohydrate Chemistry, Vol. I, p. 247 - 277.
- 110) C. H. W. Hirs, S. Moore and W. H. Stein, 1951, 'Isolation of Amino acids by chromatography on Ion-exchange Column, Use of Volatile Buffers, J. Biol. Chem. 14, pp. 669-84.
- 111) S. Moore and W. H. Stein, 1954, 'Procedure for chromatographic determination of Amino Acids in four Percent Cross Linked Sulfonated Polystyrene Resins', J. Biol. Chem., pp. 893 - 97.

- 112) R. Consden, A. H. Gordon and A. J. P. Martin, 1944,  
'Partition Chromatography', Biochem. J. 38, p. 224.  
ibid.
- 114) J. Bascur, J. Cabello, M. Veliz, A. Gonzalez, 1966,  
'Molecular Forms of Human Liver Arginase', Biochim.  
Biophys. Acta. 128, pp. 149 - 54.
- 115) J. Beruter, J. P. Colombo, C. Bachmann, 1978,  
'Purification and Properties of Arginase from Human  
Liver and Erythrocytes', Biochem. J. 175, pp. 449-54.
- 116) J. Cobello, C. Basilio, V. Prajoux, 1961, Kinetic  
properties of Erythrocyte and Liver Arginase',  
Biochim. Biophys. Acta., 48, pp. 148 - 152.
- 117) J. L. Rosenfield, S. P. Dutta, G. B. Chdeda, G. L.  
Tritsch, 1975, 'Purine and Pyrimidine inhibitors of  
Arginase', Biochim. Biophys. Acta., 4107, pp. 164  
-66.
- 118) V. L. Villanueva, M. Giret, 1980, 'Human Platelet  
Arginase', Mol. Cell Biochem., 33 (1-2), pp. 97-100.
- 119) W. H. Stein and S. Moore, 1950, Chromatographic  
determination of Amino Acids Composition of Protein',  
Cold Spring Harbor, Symp. Quart. Biol. 14, p. 179.
- 120) G. H. Rabinson, 1978, 'Factors affecting the  
seperation of Amino Acids on Ion-Exchange Columns',  
Amino Acid Determination, p. 47.

- 121) H. B. Vickery, 1925, 'Some Nitrogenous Constituents  
of the Plant Juice of Alfalfa', J. Biol. Chem. 65,  
pp. 657 - 64.
- 122) J. F. Thompson
- 123) B. S. Somvanshi, 1982, A Ph.D. Thesis, Marathawada  
University, Auranganbad.