
BIBLIOGRAPHY

- ✓ *Aalders, A.J.G. and Pieters,R. (1987). Resistance in Vicia faba to Orobanche crenata true resistance versus hidden susceptibility. Euphytica, 36(1) : 227-236.
- ✓ Aber,M., Fer,A. and Salle,G. (1983). Transfer of organic substances from the host plant Vicia faba to the parasite Orobanche crenata Forsk. International J.Plant Physiol., 112(1) : 297-308.
- *Addy,S.K. (1956). Preliminary experiments on the control of Orobanche on brinjal by 'Crag' Herbicide 1. Sci.Cult., 22 : 231-232.
- ✓ Agarwal,M.L., Kumar,S., Goel,A.K. and Tayal,M.S. (1982). Biochemical analysis in leaf spot disease of turmeric : Some hydrolysing and oxidative enzymes and related chemical metabolites. Indian Phytopath., 35(3) : 438-441.
- ✓ *Agrawal,K.C. (1982). Mineral status of Cistanche tubulosa and the host roots. Acta.Bot.Indica, 11(2) : 7-12.
- Ahmad,I., Owera,S.A.P., Farrar,J.F. and Whitbrand,R. (1982). The distribution of five major nutrients in barley plants infected with brown rust. Physiol.Plant Pathol., 21(3) : 335-346.
- ✓ *Alabi,R.O. and Naqvi,S.H.Z. (1977). Effect of infection by Cercospora arachidicola on the food contents of groundnut leaf. Trans.Br.Mycol.Soc., 68 : 295-307.

- *Aleksiev,A. (1967). Trench ploughing as a control measure against broomrape on tobacco. Bulg.Tyutym, 12(3) : 13-16.
- Allen,P.J. (1954). Physiological aspects of fungus diseases of plants. Ann.Rev.Plant Physiol., 5 : 225-248.
- ✓ Arnon,D.I. (1949). Copper enzymes in isolated chloroplasts : Polyphenol oxidase in Beta vulgaris. Plant Physiol., 24:1-15.
- Bains,S.S. and Jhooty,J.S. (1978). Relationship between mineral nutrition of muskmelon and development of downy mildew caused by Pseudoperanospora cubensis. Plant and Soil, 49 : 85-90.
- ✓ Balasubramanian,K.A. (1975). Pattern of distribution of certain cations in downy mildew affected sorghum and soil. Plant and Soil, 43(3) : 621-626.
- Balsubramanian,K.A. (1981). Chlorophyll content and mineral composition of downy mildew affected chlorotic leaves of sorghum. Indian Phytopath., 34(4) : 500-501.
- *Bamber,C.J. (1916). Plants of Panjab. Superintendent, Government Printing Press, Panjab.
- Barna,B. and Janos,E. (1981). The role of protein metabolism of wheat in amino acid induced resistance to rust. Acta Phytopathologica Academiae Scientiarum Hungaricae, 16(1/2) : 49-57.
- *Beck-Mannogetta,G. (1930). Orobanchaceas. Pflanzenreich, IV, 261, pp.348.

- K.
- Benoit,M.A., Thiagalingam and K.L.Khew (1978). Mineral nutrition and plant disease relationship. Proc.Pl.Prot.Conf., Kualalumpur, pp.67-71.
- ✓ *Betz,H., Schlosser,E. and Landbouwwet, Meded, Fac. (1980). Plant disease product quality. 1. Effect of club root (Plasmodiophora brassicae. Woronin.) on the quality of three kinds of cabbage. Rijksuniv. Gent., 45(2) : 141-145.
- ✓ Bhargava,L.P., Handa,D.K. and Mathur,B.N. (1976). Occurrence of Orobanche indica on Trigonella foenum-graecum and Phasalis minima. Plant Dis.Res., 60(1C) : 871-872.
- ✓ Bishop,N.J. (1966). Partial reactions of photosynthesis and photoreduction. Ann.Rev.Plant Physiol., 17 : 185-208.
- ✓ Boardman,N.J. (1975). Trace elements in photosynthesis, p.199-212. In : Trace Elements in Soil Plant Animal Systems. Nicholas, Ed. Egan,D.J.D. and Egan,A.R. Academic Press.
- ✓ *Boskovic,M. (1962). Crobanche on sunflower. Sborn.Mat.Spiske Prirodne Nauk, 22 : 44-65.
- ✓ *Bottrill,D.E. and Pollassingham (1969). Biochim.Biophys.Acta., 189 : 80.
- ✓ Brown,R. (1965). The germination of angiospermous parasite seeds. - In : Encyclopaedia of Plant Physiology, 15(2) : 925-932. Springer.

- ✓ Brownell,P.F. and Crossland,C.J. (1972). The requirement for sodium as a micromutrient by species having the C₄ dicarboxylic photosynthetic pathway. Plant Physiol., 49 : 794-797.
- ✓ Burstrom,H.G. (1968). Calcium and plant growth. Biol.Rev., 43 : 287-316.
- *Central Tobacco Research Institute, India (1966). Selected reference on Orobanche (1950-1964). DOC List.1.
- ✓ Chahal,A.S. and Kang,M.S. (1979). Changes in free amino acids and sugars in the leaves of brown sarson (rape) infected by Alternaria brassicae. J.Res. (Panjab Agri.Univ.), 16(1) : 57-59.
- ✓ Chaudhary,S.K. and Prasad,M. (1974). Variations in sugar contents of healthy and Fusarium oxysporum, F.udum infected plants of Cajanus cajan. Phytopath.Z., 80(4) : 303-305.
- ✓ Chen,C.T. and Chen,M.J. (1974). Pathological effects of the sugarcane white leaf agent on chlorophyll content and chloroplast ultrastructure. Proc.ISSCT., 15 : 343-347.
- ✓ Choudhari,R.S. and Garg,O.K. (1955). Effect of little leaf disease on the growth and physiology of Brinjal (Solanum melongena L.). Indian J.Hort., 12 : 180-187.
- ✓ Cruickshank,I.A.M. and Perrin,D.R. (1964). Pathological function of phenolic compounds in plants. - In : Biochemistry of phenolic compounds (ed. J.B.Harborne). pp.511-544. Academic Press, London and New York.

- ✓ *Dakshinamurty,V., Govindarao,P. and Reddy,G.S. (1964). A survey of torko (Orobanche cernua Loefl. var. desertorum Beck) incidence on tobacco in coastal Andhra districts. Andhra Agric.J., 11 : 152-156.
- Davis,E.L. and Rich,J.R.(1987). Nicotine content of tobacco roots and toxicity to Meloidogyne incognita. J.Nematol., 19(1) : 23-29.
- ✓ Dawson,R.F. (1942). Accumulation of nicotine in reciprocal grafts of tomato and tobacco. Am.J.Bot., 29 : 66-71.
- ✓ Dekock,P.C. (1964). The physiological significance of the potassium-calcium relationship in plant growth. Outlook on Agriculture, 4(2) : 93-98.
- ✓ Deshmukh,S.S. and Mayee,C.D. (1978). Pearl millet mildew development as influenced by some nutrients. J.OF MAU, 3(1) : 79-80.
- ✓ Dhingra,R.K., Chauhan,N. and Chauhan,S.V.S. (1982). Biochemical changes in the floral part of Brassica campestris infected by Albugo candida. Indian Phytopath., 35(1) : 177-179.
- ✓ Dixit,R.B. and Gupta,J.S. (1983). Post infection alterations in the composition of organic acids and phenolic substances in leaf leachates of barley infected with Alternaria alternata (FR.) KEISSEER. Geobios, 10 : 122-123.
- ✓ Dorr,I. and Kollmann,R. (1975). Structural features of parasitism of Orobanche. II. The differentiation of assimilate conducting elements within the haustorium. Protoplasma, 83(3) : 185-200.

- ✓ Dorr,I. and Kollmann,R. (1975). Structural features of parasitism of Orobanche. III. The differentiation of xylem connection of O.crenata. Protoplasma, 89(3/4) : 235-249.
- ✓ Dubey,L.N. and Joshi,R.D. (1985). Influence of cucumber mosaic virus on total nitrogen and reducing sugar content of Chilli (Capsicum annum). Indian Journal of Mycology and Plant Pathology, 15(2) : 221-223.
- Duggal,S., Singh,R.A. and Lakhanpol,T.N. (1981). Physiological studies on potato virus X infected Capsicum penicillum Plants-I. Indian J.Mycol.and Pl.Pathol., 11(1) : 84-88.
- ✓ Eames,A.J. and Laurence,H.M. (1972). An Introduction to Plant Anatomy. Publ. McGraw-Hill Company Ltd., Bombay, New Delhi.
- *Ekaterina,S. (1959). Invertase and peroxidase of sunflower in relation to resistance to broomrape (Orobanche cernua). Invest.Inst.Rast.Bulgar.Akad.Nauk., 7 : 151-152.
- Epstein,E. (1972). Mineral Nutrition of Plant :Principles and Perspectives. John Wiley and Sons, Inc., New York, London, Sydney, Toronto.
- ✓ Farkas,G.L. and Kiraly,Z. (1962). Role of phenolic compounds in the physiology of plant diseases and disease resistance. Phytopath.Z., 44 : 105-150.
- ✓ Folin,O. and Denis,W. (1915). A colorimetric estimation of phenols (Phenol-derivatives) in Urine. J.Biol.Chem., 22 : 305-308.

- Gangopadhyay,S. and Chattopadhyay,S.B. (1977). Biochemical changes of rice plants at different levels of nitrogen in relation to brown spot disease development. I. Nitrogen. Indian Phytopath. , 28(3) : 326-329.
- Garg,I.D. and Mandahar,C.I. (1976). Physiology of powdery mildew infected leaves of Abetmoschus esculentus lll. Carbohydrates. Indian Phytopath., 29 : 173-177.
- Garg,I.D. and Mandahar,C.I. (1977 a). Effect of net blotch infection on the physiology of barley : Carbohydrate and nitrogen metabolism. Indian J.Exp.Biol., 14 : 66-68.
- Garg,I.D. and Mandahar,C.I. (1977 b). Physiology of Phaseolus mungo leaves infected with bean yellow mosaic virus. Indian Phytopath., 30 : 123-125.
- *Gharib,M.S. (1973). Biological and economical aspects of broomrape (Orobanche spp.) in nothern Iraq. Proc.Eur.Weed.Res.Counc.Symp.Parasit.Weeds, pp.44-47.
- Ghorpade,L.N. and Joshi,G.V. (1981). Studies in sugarcane mosaic virus in sugarcane var.CO 740. Indian J.Plant Physiol., 24(1) : 1-7.
- ✓ *Gill,L.S. and Hawksworth,F.G. (1961). The mistletoe : A literature review. U.S.Dep.Agric., Tech.Full., No.1242.
- ✓ Goel,A.K., Kumar,S. and Tayal,M.S. (1983). Biochemical alteration induced in Coriandrum sativum L. by Protomyces macrosporus Unger. Indian Bot.Reptr., 2(1) : 62-64.

- ✓Gonzoles-Torres,R., Jimenez-Diaz,R.M. and Melero-Vara,J.M. (1982). Distribution and virulence of Orobanche cernua in sunflower crops in Spain. Phytopathol.Z., 104(1) : 78-89.
- ✓Goswami,B.K., Singh,S. and Verma,V.S. (1976). Uptake and translocation of calcium and magnesium in tomato plants as influenced by infection with root-knot nematode Meloidogyne incognita and tobacco mosaic virus. Nematologica, 22 : 116-117.
- ✓Greenham,C.G. and Leonard,O.A. (1965). The amino acids of some mistletoes and their hosts. Am.J.Bot., 52(1) : 41-47.
- ✓Gujar,V.R. and B.A.Hegde (1983). Effect of TMV on photosynthesis and nicotine content of tobacco. Tobacco Res., 9(1) : 25-28.
- ✓Gupta,R.N. (1975). Mineral matter content in coriander leaves and fruit as influenced by stem gall disease. Indian Phytopath., 28 : 136-137.
- ✓Gupta,J.P. and Chatrath,M.S. (1979). Physiology of resistance to anthracnose in guava. Indian Phytopath., 32(1) : 64-67.
- Hall,S.M. and Baker,D.A. (1972). The O' Chemical composition of Ricinus phloem exudate. Planta, 106 : 131-140.
- Hare,R.C. (1966). Physiology of resistance to fungal diseases in plants. Bot.Rev., 32 : 95-137.
- Harms,H. and Terbea,M. (1984). Metabolism of phenolic compounds in healthy and brown-rust infected barley and wheat varieties. Phytopathol.Z., 111(3/4) : 283-296.

- *Hassan, A.A. and Khalid, E.A. (1986). Assessment of broomrape tolerance in the genus Lycopersicon. Egypt J.Hortic., 13(2) : 153-158.
- Hawk, P.B., Oser, B.L. and Summerson, W.H. (1948). 'Practical Physiological Chemistry' Publ. The Blakiston Company, U.S.A.
- Hegde, R.K. (1967). Studies on bean anthracnose organism Colletotrichum lindemuthianum (Sacc. and Magn.) Bri. and Cav. in India. Ph.D. Thesis, I.A.R.I., Delhi.
- ✓ Hegde, B.A. and Karande, S.M. (1978). Effect of presowing treatment of sodium chloride on the incidence of green ear disease of Pennisetum typhoides (Burm) Stapf. and Hubb var. HB₃. Plant and Soil, 49(3) : 551-559.
- Heitefuss, R. and Williams, P.H. (1976). Physiological Plant Pathology. Encyclopedia of Plant Physiology, Vol. 4. eds. A. Pirson, M.H.Z. Immermann. Berlin. Springer-Verlag. pp. 890.
- ✓ Hodosy, S. (1980). Biological control of Orobanche ramosa, a parasite of tomato. 1. Hyperparasitic Fusarium species in the control of broomrape in Hungary. 14 : 21-29.
- Hopkins, D.L. and Hampton, R.E. (1969). Effect of tobacco etch virus infection upon the dark reactions of photosynthesis in tobacco leaf tissue. Phytopath. , 59 : 1136-1140.
- ✓ Horak, O. (1974). Comparative analysis on the mineral ion content of some Loranthaceae and their hosts. Z.Pflanzenphysiol., 73(5) : 461-466.

- ✓ Horowitz, W. (1965). Silicotungstic Acid Method (S_1). In : "Official methods of Analysis". A.O.A.C. pp.48.
- ✓ Horsfall, J.G. and Dimond, A.E. (1957). Interaction of tissue sugar, growth substances and disease susceptibility. J.Pflanzenkrnkh. 64 : 415-421.
- ✓ Huber, D.M. (1978). Disturbed mineral nutrition. In : Plant disease an advanced treatise. Vol.III. eds. Horsfall, J.G. and Cowling, E.B. Academic Press Inc., New York, pp.163-200.
- ✓ Hull, R.J. and Leonard, O.A. (1964). Physiological aspects of parasitism in mistletoe (Arceuthobium and Phoradendron) I. The carbohydrate nutrition in mistletoes. II. The Photosynthetic capacity of mistletoe. Plant Physiol., 39 : 996-1017.
- ✓ Hwang, B.K., Ibenithal, W.D. and Heitefuss, R. (1983). Age, rate of growth, carbohydrate and amino acid contents of Spring barley plants in relation to their resistance to powdery mildew (Erysiphe graminis & Sp. hardei). Physiol. Plant Pathol., 22(1) : 1-14.
- ✓ Indian Central Tobacco Committee (1960). Indian Tobacco - A monograph.
- * Iskender, G. (1975). Quantitative estimation of the alkaloids of mildew resistant turkish tobacco. Istanbul Univ. Eczacilik Fak. Mecm., 11(1) : 27-37.
- ✓ Ismail, A.M.A. and Obeid, M. (1976). A study of assimilation and translocation in Cuscuta hyalina Heyne ex. Roth. Orobanche ramosa L. and Striga hermonthica Benth. Weed Res., 16(2) : 87-92.

- ✓ *Izard,C. (1959). L' Orobanche du tabac. Ann.Inst.Expl.Tabac.
Bergerac, 3 : 299-316.
- ✓ Jaworski,E.G. (1971). Nitrate reductase assay in intact plant tissue. Biochem.Biophys.Res.Commun., 43 : 1274-1279.
- ✓ Johansen,D.A. (1940). Plant Microtechnique. McGraw-Hill Co., New York. pp.523.
- Johri,J.K. (1975). Post infection changes in nitrogen metabolism in virus infected Carica papaya L. Indian J.Exp.Biol., 13 : 220-221.
- ✓ Johri,J.K. and Padhi,B.(1981). Biochemical changes induced by Cercospora arachidicola in groundnut leaves. Third International Symposium on Plant Pathology, New Delhi (Abstr.) pp.189.
- ✓ *Kadry,A.E.R. and Tewfic,H. (1956). Seed germination in Orobanche cernua Forsk. Svensk.Bot.Tidsk., 50 : 270-286.
- ✓ Kapoor,I.J. and Tandon,R.N. (1969). Post infection changes in organic acids of tomato under pathogenesis. Naturwissenschaften, 56 : 40-41.
- ✓ Kapur,S., Kapur,S.P. and Rehalia,A.S. (1978). Studies on some of the biochemical changes occurring in date palm leaves infected by smut. Indian Phytopath., 31(3) : 394-395.
- ✓ Kasasian,L.(1973). Miscellaneous observations on the biology of Orobanche crenata and O.aegyptiaca. Proc.Eur.Weed.Res.Counc.Symp.Parasit.Weeds, pp.68-75.

- Kauraw, L.P. (1979). Effect of different soil types, pH and phosphorus levels on the wheat seedling and the incidence of Phythium graminocolum. Plant and Soil, 53(4) : 551-570.
- ✓ Khanna, S.K., Viswanathan, P.N., Tewari, C.P., Krishnan, P.S. and Sanwal, G.G. (1968). Biochemical aspects of parasitism by angiosperm parasites : Phenolics in parasites and hosts. Physiol. Plant., 21 : 949-959.
- ✓ *Knutson, D.M. and Toevs, W.J. (1973). Dwarf mistletoe reduces root growth of Ponderosa pine seedlings. For. Sci., 18(4) : 323-324.
- ✓ *Kolhoff, I.M. and Noponen, G.E. (1933). J. Am. Chem. Soc., 55 : 1148-1453.
- ✓ Krishnamurty, G.V.C., Nagarajan, K. and Lal, R. (1977). Some studies on Orobanche cernua Loefl., a Parasitic Weed on Tobacco in India. Indian J. Weed. Sci., 9(2) : 95-106.
- ✓ Krishnarao, N.S.R. and Nayadu, M.V. (1979). Changes in organic constituents of rice leaves infected by Xanthomonas oryzae. Phytopathol. Z., 94(4) : 357-366.
- Kuijt, J. (1955). Dwarf mistletoes. Bot. Rev., 21 : 569-626.
- ✓ *Kuijt, J. (1969). The Biology of parasitic flowering plants. Univ. Calif. Press, Berkeley.
- Kulkarni, D.K. and Kulkarni, U.K. (1978). Physiology of mango leaves infected by Capnodium ramosum Cooke : II. Mineral contents. Biovigyanum, 4 : 173-174.

- ✓ Kumar, N.C. and Mukherjee, K.L. (1973). Host-parasite relation in Dendrophthoe falcata infection. Indian Phytopath., 26 : 148-151.
- ✓ *Lee, A. and Goseco, F. (1932). Studies of the sugarcane root parasite Aeginetia indica. Proc. Int. Soc. Sugarcane Technol. Congr., San Juan, 4 : 1-12.
- ✓ Livne, A. (1964). Photosynthesis in healthy and rust-affected plants. Plant Physiol., 39 : 614-621.
- ✓ Lolas, P.C. (1986). Control of broomrape (Orobanche ramosa) in tobacco (Nicotiana tabacum). Weed Sci., 34(3) : 427-430.
- ✓ Loneragan, J.F., Grove, T.S., Robson, A.D. and Sunowball, K. (1979). Phosphorus toxicity as a factor in zinc-phosphorus interactions in plants. Soil Sci. Soc. Am. J., 43 : 966-972.
- ✓ *Loomis, L.C. and Wilson, W.E. (1953). Botany Holt., Rinehart and Winston, New York.
- ✓ Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with the folin phenol reagent. J. Biol. Chem., 193 : 262-275.
- ✓ MacDonald, P.W. and Strobel, G.A. (1969). Starch accumulation in wheat plants infected with stripe rust. Phytopathol., 59 : 1139 (Abstr.)
- *MacLeod, D.G. (1962). Some anatomical and physiological observations on two species of Cuscuta. Trans. Bot. Soc. Edinb., 39 : 302-315.

- ✓ MacLeod,D.G. (1963). The parasitism of Cuscuta. New Phytol., 62 : 257-263.
- ✓ Magyarosy,A.C., Schurmann,P. and Buchanan,B.B. (1976). Effect of powdery mildew infection on photosynthesis by leaves and chloroplasts of sugarbeets. Plant Physiol., 57 : 486-489.
- Mahadevan,A. (1966). Biochemistry of infection and resistance. Phytopath.Z., 57 : 96-99.
- Majumdar,M. and Raychaudhuri,S.P. (1978). Changes in water, carbohydrates and nitrogen contents of pea plants during the pathogenesis of Fusarium solani infection. Ann.Physio-pathol.Soc.Jon., 48 : 199-201.
- ✓ Malik,C.P. and Komal,A. (Unpublished). Cf.Malik,C.P. and Singh, M.B. (1979). Physiological and biochemical aspects of parasitism in Cuscuta. A review. Ann.Rev.Plant Sciences, (C.P.Malik ed.) Kalyani Publishers, New Delhi, pp.67-113.
- ✓ Mandokhot,A.M., Singh,D.F., Chaudhary,K.C.E. and Singh,J.S. (1979). Chemical changes in maize leaves in response to leaf spot pathogen. Indian Phytopath., 32(4) : 658-660.
- ✓ Marimuthu,T. and Kandaswamy,T.K. (1981). Role of soluble sugars in the resistance or susceptibility of green gram varieties to bacterial leaf blight caused by Xanthomonas phaseoli. Curr.Sci., 50(14) : 643-645.
- ✓ Marudarajan,D. (1950). Note on Orobanche cernua Loefl. Curr. Sci., 19 : 64-65.

- ✓ Mattoo, R.L., Viswanathan, P.N. and Krishnan, P.S. (1969). Biochemical aspects of parasitism by the angiosperm parasites : Host-parasite interrelationship in phosphatase activity. Physiol. Plant., 22 : 638-647.
- ✓ *Mc Dowell, L.L. (1964). Physiological relationships between dwarf mistletoe and Ponderosa pine. Ph.D.Thesis. Oregon State Univ. Corvallis, Diss. Abstr., 25(1) : 53.
- *Mc Murtrey, J.E. Jr. (1941). Plant nutrient deficiency in tobacco. In "Hunger Signs in Crops", 15-54. American Soc. Agron. and American Fertilizer Assoc. Washington, D.C.
- Mehta, P.K., Vyay, M. and Saksena, S.B. (1975). Metabolic changes during pathogenesis of fruit rot disease of tomato. Indian Phytopath., 28 : 253-255.
- Mengel, E. and Kirkby, E.A. (1982). Principles of Plant Nutrition. International Potash Institute. Worblaufen-Bern/Switzerland.
- ✓ Meyer, B.S., Anderson, D.B. and Bohning, R.H. (1960). Introduction to plant physiology. Van Nostrand Company Inc., New York, pp.185.
- ✓ *Mijatovic, K. and Stajanovic, D. (1973). Distribution of Orobanche spp. on the agricultural crops in Yugoslavia. Proc. Eur. Weed Res. Counc. Symp. Parasit. Weeds., pp. 28-34.
- ✓ Mishra, U.S., Saxena, S.K. and Pramod Kumar (1989). Amino acid composition in seedlings of Sesamum indicum L. infected with Aspergillus niger. In Proceedings of the Seventy Six Session of the Indian Science Congress, Madurai, 83 : 47-48.

- Misra,A. and Padhi,B.(1981). Physiological alterations due to white rust infection in Ipomoea reoptans. IIIrd Intern.Sym. on Plant Pathol., (Abstr.) pp.57.
- Misra,A.K., Rao,N.N. and Ganapati,M.M. (1981). Zinc deficiency in Coorg mandarin associated with Phytopathora population in soil. IIIrd Intern.Symp.on Plant Pathology, New Delhi, (Abstr.) pp.38.
- Misra,P.C., Setty,P.N., Singh,B.V., Lal,R.K., Viswanathan,P.N., Saxena,Y.R. and Krishnan,P.S. (1970). Heterogeneity in composition along the length of Cuscuta filaments. Physiol. Plant., 23 : 1025-1032.
- Mitra,D.K. and Majumdar,M. (1980). Absorption of ^{32}P by brinjal and tobacco leaf tissues infected with little leaf and tobacco mosaic virus respectively. Indian Phytopath., 33(4): 619-620.
- Mitra,D.K. and Sengupta,U.K. (1980). Influence of little leaf disease on photosynthesis and chlorophyll content in brinjal (Solanum melongena Linn.) leaf tissues. Ann.Agric.Res., 1(1) : 108-111.
- Mogle,T.R. and Mayee,C.D. (1981). Changes in reducing sugars, starch and free amino acid accompanying downy mildew infection in resistant, moderately resistant and susceptible lines of pearl-millet. Indian J.Mycol.Plant Pathol., 11(1) : 61-65.

- ✓ Mohan,R., Murugappan,V., Doraiswamy,S. and Marimuthu,T. (1978). Changes in mineral content in chilli leaves to graded doses of potash and its effect on Xanthomonas infection. Indian Potash J., 3 : 2-4.
- Mukherjee,M., Srivastava,H.S. and Bhargava,K.S. (1979). Effect of cucumber mosaic virus infection on some nitrogenous components in tobacco. (Nicotiana tabacum L.) var. White Burley. Indian J.Exp.Biol., 17(1) : 116-117.
- ✓ Musselman,L.J. and Nixon,K.C. (1981). Branched broomrape (Orobanche ramosa) in Texas, USA. Plant Dis., 65(9) : 752-753.
- ✓ Musselmann,L.J. and Parker,C. (1982). Preliminary host ranges of some strains of economically important broomrapes (Orobanche). Econ.Bot., 36(3) : 270-273.
- ✓ Nassib,A.M., Ibrahim,A.A. and Saber,H.A. (1978). Broomrape (Orobanche crenata) resistance in broad beans : Breeding work in Egypt : A review. In 'Food, Legume Improvement and Development' Ed. Geoffrey C.Hawtin and George J. Chancellor. pp.133-135.
- ✓ Nelson,N. (1944). A photometric adaptation of the Somogyi method for the determination of glucose : J.Riol.Chem., 153:375-380.
- ✓ Nicholas,D.J.D. (1961). Minor mineral nutrients. Ann.Rev.Plant Physiol., 12 : 63-90.
- ✓ Nicoloff,T.H. (1923). Contribution a la physiologie de la nutrition des parasites vegetaux superieurs. Rev.Gen.Bot., 25 : 545-554, 593-601.

- ✓ Nour-Eldin, F. (1955). The effect of organic acids on TMV - multiplication. Phytopath., 45 : 291.
- ✓ Okasha, K.A., Ryugo, K. and Bringhurst, R.S. (1968). Relationship of tannins, polyphenols and reducing sugar to Verticillium wilt resistance of strawberry cultivars. Phytopath., 58 : 1118-1122.
- ✓ Orabi, A.A., Abdallah, A., Mashadi, H. and Barakat, A.R. (1981). Zinc-phosphorus relationship in the nutrition of corn plants (Zea mays L.) grown on some calcareous soils. Plant and Soil, 59 : 51-59.
- Patil, B.D. and Kulkarni, U.K. (1977 a). Physiological changes in sunflower leaves due to infection of Puccinia helianthi Schw. Biovigyanum, 3 : 11-15.
- ✓ Patil, B.D. and Kulkarni, U.K. (1977 b). Physiological changes in the leaves of sunflower due to infection of Puccinia helianthi Schw. Indian Phytopath., 30(4) : 560-561.
- ✓ Patil, L.K., Wani, P.V. and More, B.B. (1979). Chromatographic studies on the amino acid metabolism of healthy and diseased leaves of durum wheat (Triticum durum) infected by Alternaria triticola. J. Maharashtra Agric. Univ., 4(3) : 268-270.
- ✓ Pennypacker, B.W., Nelson, P.E. and Wilhelm, S. (1979). Anatomic changes resulting from parasitism of tomato by Orobanche ramosa. Phytopath., 69(7) : 741-748.
- Perkin-Elmer Corp. (1971). Analytical methods for atomic absorption spectrophotometry. Norwalk, Connecticut, U.S.A.

- ✓ Pero,R.W., Main,C.E. (1970). Chlorosis of tobacco induced by alternariol manomethyl ether produced by Alternaria tennise. Phytopath., 66 : 1570-1573.
- ✓ Prasad,B.K. (1983). Free sugar and amino acid contents of coriander infected with stem gall disease. Indian Phytopath., 36(1) : 150-151.
- ✓ Prasad,N. (1954). Control of Orobanche in tobacco by Crag herbicide 1. Indian Tobacco, 4(3) : 139.
- ✓ *Prasad,M. and Chaudhary,S.K. (1974). Amino acids and fusaric acid production in tissue of Cajanus cajan infected with Fusarium oxysporum, F.udum. Phytopathol.Z., 81(4) : 339-345.
- ✓ Pridham,J.B. (1965). Low molecular weight phenols in higher plants. Ann.Rev.Plant Physiol., 16 : 13-36.
- ✓ Purohit,S.D., Ramawat,K.G. and Arya,H.C. (1979). Polyphenols and some oxidative enzymes in rust infected leaves of Cyperus rotundus. Indian Phytopath., 32 : 255-259.
- ✓ Raggi,V. (1974). Synthesis of free amino acids in rusted bean (infected and non-infected) leaves. Phytopathol.Z., 81(3) : 262-272.
- *Rakhimov,Yu. Kh. (1967). Comparative analysis of the susceptibility of host plants to certain species of Orobanche L. Ispol.Rastit.Resursov-Povysh.Prod.Kult.Rast., 77-81.
- ✓ *Rangaswami,N.S.(1963). Studies on culturing seeds of O.aegyptiaca. - In plant tissue and organ culture. Symp.Int.Soc.Plant Morphologists, Delhi (Eds. Maheshwari,P. and Rangaswami,N.S.).
- ✓ Rao,D. (1953). New hosts of Orobanche. Curr.Sci., 22 : 311.

- ✓ Rao,R.S.N., Joshi,B.G. and Sharma,C.B. (1982). Maintenance of Orobanche cernua on brinjal plants, (Solanum melongena). Indian J.Agric.Sci., 52(11) : 764-767.
- ✓ Rao,S.C., Sreenivasa Rao,B. and Naidue,V.D.(1978). Effect of nitrogen on the biochemistry of rice leaves in relation to bacterial leaf streak disease. Indian Phytopath., 31 : 547-548.
- ✓ Rawat,V.S. and Ambasht,R.S. (1958). Root relations of Orobanche and its hosts. Curr.Sci., 27 : 445-446.
- ✓ Raychaudhuri,S.P. and Varma,A. (1980). Sandle spike. Rev.Plant Pathol., 59(3) : 99-107.
- ✓ Reddy,M.N. and Ramagopal,G. (1982). Changes in organic acids and sugars in the rust infected groundnut leaves. Indian Phytopath., 35(4) : 728-730.
- ✓ *Rensing,L. and Corenlius,G. (1980). Biological membranes as components of oscillating systems. Biol.Rdsch., 18:197-209.
- ✓ Roberts,B.R. and Jensen,K.F. (1970). The influence of dutch elm disease and plant water stress on the foliar nutrient content of American and Siberian elm. Phytopathol., 60 : 1831-1833.
- ✓ Roberts,P.L. and Wood,K.R. (1982). Effects of a severe (P₆) and a mild (W) strain of cucumber mosaic virus on tobacco leaf chlorophyll, starch and cell ultrastructure. Physiol. Plant Pathol., 21(1) : 31-37.

- Sadik,A., Rey,L. and Renqudin,S. (1986). The endophytic system of Arceuthobium oxycedri. II. Ultrastructural aspects of contact zones, between tissues of host and parasite. Can.J. Bot., 64(11) : 2778-2784.
- ✓ Saghir,A.R., Dastgheib,F. (1978). The biology and control of Orobanche : A Review, in : 'Food Legume Improvement and Development'. Ed.Geoffey,C. Hawtin and George J.Chancellor. pp. 126-132.
- ✓ Saghir,A.R., Foy,C.L., Hameed,K.M., Drake,C.R. and Tolin,S.A. (1973). Studies on biology and control of Orobanche ramosa L. Proc.Eur.Weed Res.Counc.Symp.Parasit.Weeds, pp.106-116.
- ✓ Saini,S.S. and Suppal,H.S. (1982). Changes in sugar and amino acid content in healthy and diseased leaves of chicory. Geobios., 5(3) : 121-124.
- Sankpal,S.D. (1981). Physiological studies in sugarcane. (Post infection physiological changes in smut infected sugarcane Var. Co 740). Ph.D. Thesis, Shivaji University, Kolhapur, India.
- ✓ Sankpal,S.D. and Nimbalkar,J.D. (1979). Effect of smut infection on sucrose metabolism in sugarcane. Indian Phytopath., 32(2): 306-307.
- ✓ Sankpal,S.D. and Nimbalkar,J.D. (1980). Physiological studies in smut infected sugarcane Var. Co-740. Indian J.Exp.Biol., 18(1) : 95-96.

- ✓ Sarkar,K.R. and Joshi,R.D. (1977). Effect of little leaf disease on the contents of certain essential nutrient elements in brinjal. Indian Phytopath., 30(3) : 429-430.
- ✓ Sastry,K.S.M. (1964). Effect of trace elements on multiplication and concentration of sunn hemp mosaic virus. Indian J.Exp.Biol., 2 : 185-189.
- ✓ Sastry,K.S.M. and Nariani,T.K. (1962). Effect of host nutrition on growth and susceptibility of tobacco plants to infection with tobacco leaf curl virus. Indian J.Agric.Sci., 32 : 288-293.
- ✓ Schlegel,D.E. (1957). The stimulatory effect of organic acids on tobacco mosaic virus multiplication. Virology, 4 : 135-140.
- ✓ Schmucker,Th. (1959). Hohere Parasiten. - In Encyclopaedia of Plant Physiology, 11 : 480-529. Springer.
- ✓ Sekine,T., Sasakawa,T., Morita,S., Kimura,T. and Kuratomi,K. (1965). Photometric Colometry in Biochemistry (Part-II). Nanko-do Publishing Co. Tokyo. pp.242.
- ✓ Sharma,S.G., Ram Narayan, Sangam Lal and Chaturvedi,C. (1983). Role of phenolic compounds in resistance of maize to leaf blight caused by Drechslera State of Cochliobolus heterostrophus. Indian Phytopath., 36(1) : 43-46.
- ✓ Shaw,M. (1963). The Physiology and host-parasite relations of the rusts. Ann.Rev.Phytopath., 1 : 259-294.

*Shevchenko,I.P. Golovchak,E.I.M., Serede,O. (1980). In : Seed treatment with trace elements. Eds. Vlasyuk, P.A.Izd. Naukova Dumka : Kiev, USSR. pp.189-193.

✓ Siddaramaiah,A.L., Vasuki,N., Barama Goudar,T.D., Lingaraju,S. and Hegde,R.K. (1979). Biochemical changes in rust infected leaves of groundnut. Indian Phytopath., 32 : 640-641.

*Silberschmidt,K. (1930). Der Einfluss der Mosaikkrankheit auf den Nikotengehalt der Tabakflanze. Ber Deutsch Bet.,Ges., 48 : 122-129.

✓ Sindho Mathar,A. and Vidhyasekaran,P. (1978); Physiology of resistance to rust in sunflower. Indian Phytopath., 31 : 289-293.

Singh,B.P. and Verma,A.K. (1974). Influence of hollyhock yellow mosaic virus on flowering and chemical composition of hollyhock leaves. Indian Phytopath., 27 : 72-77.

✓ Singh,C.T.N. and Prasad,M. (1989). Qualitative and quantitative changes in sugar contents of tomato fruit during pathogenesis by Corenbacterium michiganense pv. michiganese. Proceedings of the Seventy Six Session of The Indian Science Congress, Maturai, 86 : 49-50.

✓ Singh,Jai,N. and Singh, Jagdish,N. (1971). Studies on the physiology of host-parasite relationship in Orobanche. I. Respiratory metabolism of host and parasite. Physiol.Plant., 24 : 380-386.

✓ Singh, P. and Krishnan, P.S. (1971). Effect of root parasitism by Orobanche on respiration and chlorophyll content of Petunia. Phytochemistry, 10 : 315-318.

Singh, R. and Mall, T.P. (1978). Changes in chemical composition of pigeon pea fruits due to arhar (Pigeon pea) mosaic virus infection. Acta Phytopathol. Acad. Sci. Hung., 13(1/2) : 41-44.

Singh, R. and Singh, A.K. (1977). Influence of phosphorus nutrition on the growth and nodulation of arhar mosaic virus infected sunnhemp plants. Sci. and Cult., 43 : 480-483.

✓ Singh, H., Agarwal, P.D., Bhattacharya, G., Sethi, I. (1982). Changes in chloroplast pigments in wheat leaves infected with Puccinia graminis tritici. Phytopath. Z., 104(3) : 193-201.

✓ Singh, Jai.N., Singh, Jagdish N. and Rai, T.B. (1971). Studies on the Physiology of Host-parasite Relationship in Orobanche. II. Growth and Mineral Nutrition of Host and Parasite. Physio. Plant., 25 : 425-431.

✓ Singh, Jagdish.N., Rai, T.B. and Singh, Jai.N. (1972). Studies on the physiology of host parasite relationship in Orobanche. III. Carbohydrate and nitrogen metabolism of host and parasite. Physiol. Plant., 27 : 347-353.

✓ Singh, Jagdish, N., Singh, Jai, N. and Singh, D.P. (1979). Effect on removal of host shoot on carbohydrate and mineral changes in Orobanche. Indian J. Plant Physiol., 22(1) : 30-35.

- ✓ Singh,M., Beg,M.U., Singh,D.V., Tewari,K.K. and Krishnan,P.S. (1967). Biochemical aspects of angiosperm parasites. II. Phosphatase and phosphorylase activity in dodder and infected hosts. Ind.J.Biochem., 4 : 146-151.
- ✓ Singh,M., Singh,D.V., Misra,P.C., Tewari,K.K. and Krishnan,P.S. (1968). Biochemical aspects of parasitism by the angiosperm parasites : Starch accumulation. Physiol.Plant., 21 : 525-538.
- Singh,R.S., Chaube,H.S., Singh,N., Ashani,V.L. and Singh,R. (1970). Observations on the effect of host nutritions and seed, soil and foliar treatments on the incidence of downy mildew. I.A. Preliminary Report. Indian Phytopath., 23 : 209-215.
- ✓ Sivaprakasam,K., Jagadeesan,M. and Kannapiran,C. (1976). Effect of little leaf disease on the mineral metabolism of brinjal. (Solanum melongena L.). Annamalai Univ. Agric.Res.Annual, 6 : 146-150.
- ✓ Sivaprakasam,K., Vidhyasekaran,D. and Soumini Rajagopalan,C.K. (1974). Physiology of brinjal plants infected by Verticillium dahliae. Indian Phytopathol., 27(2) : 197-202.
- ✓ *Smith,D., Muscatine,L. and Lewis,D. (1969). Carbohydrate movement from autotrophs to heterotrophs in parasitic and mutualistic symbiosis. Biol.Rev., Cambridge Philos.Soc., 44 : 17-90.
- ✓ Sohi,H.S. and Rawal,R.D. (1977). Role of certain physiological factors responsible for resistance to rust and septoria rust spot disease in cowpea. All India Symposium on Physiology of host-pathogen interaction. pp.391-398.

- ✓ Span, (1958). Killing broomrape in tobacco. Span, 2 : 14-15.
- ✓ Srinivasan,K.V. (1968). Physiology of disease resistance in sugarcane with particular reference to red rot. Curr.Sci., 37 : 34-35.
- ✓ Srinivasan,K. and Chelliah,S. (1979). Preference of the leaf hopper vector Hishimonus phycitis (Distant) for egg plants infected with little leaf mycoplasma with reference to certain biochemical and histochemical changes. Workshop held at Univ.Agric.Sci., Bangalore (India). pp.14-22.
- ✓ Swain,B.N. and Prasad,J.S. (1988). Chlorophyll content in rice as influenced by the root-knot nematode, Meloidogyne graminicola infection. Curr.Sci., 57(16) : 895-896.
- ✓ Taniguchi,T. (1962). The amount of some organic acids found in TMV-infected tobacco leaves floated on water or various organic acids. Virology, 17 : 40-43.
- ✓ Tewari,J.P., Shukla,I.K. and Srivastava,S.K. (1983). Metabolic changes due to 3 strains of watermelon mosaic virus infection in Cucurbita maxima fruits. Biologia, 29(2) : 253-256.
- ✓ Thind,B.S., Sharma,R.R. and Singh,N. (1982). Changes in sugars, amino acids and phenols in Chilli (Capsicum annuum) leaves in response to Xanthomonas vesicatorio infection. Indian J. Mycol.Plant Patho., 11(2) : 207-210.
- ✓ Thompson,D.P. (1978). Free amino acid content of sweet potato roots infected with Rhizopus stolonifer. Indian Phytopath. 31(2) : 269-270.

- ✓ Tomlinson,J.A. and Webb,M.J.W. (1978). Ultrastructural changes in chloroplast of lettuce infected with beet western yellow virus. Physiol.Plant Pathol., 12 : 13-18.
- ✓ Toth,S.J., Prince,A.L., Wallace,A. and Mikkelsen,D.S. (1948). Rapid quantitative determination of eight mineral elements in plant tissues by systematic procedure involving use of flame photometer. Soil Sci., 66 : 459-466.
- ✓ Tsui,C. (1948). The role of zinc in auxin synthesis in the tomato plant. Am.J.Bot., 35 : 172-179.
- ✓ Tugnawat,R.K. (1977). Chlorophyll a/b ratio as influenced by plant pathogens. Sci.and Cult., 43(10) : 456.
- ✓ *Upadhyay,R.K. and Dwivedi,R.S. (1979). Biochemical changes in the leaf of Eucalyptus globulus infected with Pestalotiopsis funerae. Indian Phytopath., 32 : 434-437.
- ✓ *Vaucher,J.P. (1827). Monographie des Orobanches. J.J.Paschoud, Geneva, pp.72.
- ✓ Vayonis,G.C.(1954). Phosphorus disturbances in mosaic virus infected tobacco plants. Physiol.Plant., 7 : 687-697.
- ✓ Vidhyasekaran,P. (1974). Role of phenolics in leaf spot disease incidence in ragi infected by Helminthosporium tetrrema. Indian Phytopath., 27(4) : 583-586.
- ✓ Vidhyasekaran,P. and Kandaswamy,D. (1971). Effect of soil fertility on the physiology of corn plants in relation to helminthosporiose disease incidence. Phytopath., Z., 72 : 11-20.

- ✓ Vidhyasekaran, P. and Kandaswamy, D. (1972). Carbohydrate metabolism of Phaseolus aureus infected with obligate and facultative parasites. Indian Phytopath., 25 : 48-54.
- Vidhyasekaran, P., Krishnaswamy, V. and Parambaramani, C. (1974). Possible role of sugars in rust resistance in Setaria italica. Indian Phytopath., 27 : 291-293.
- Vidhyasekaran, P., Lalithakumari, D., Subramanian, J.M. and Govindaswamy, C.V. (1971). Studies on physiology of Helminthosporium setariae, the causal agent of leaf spot disease of Setaria italica. II. Carbon nutrition. Madras Agric.J., 58 : 253-259.
- ✓ Vijayakumar, C.S.K. and Rao, A.S. (1980). Physiological changes in Alternaria infected wheat leaves. Phytopath.Z., 98 : 76-83.
- Wallace, A., Romney, E.M. and Alexander, G.V. (1978). Mineral composition of Cuscuta nevadensis Johnston (dodder) in relation to its hosts. Plant Soil, 50(1) : 227-232.
- ✓ Walter, J.M. (1967). Hereditary resistance to disease in tomato. Ann.Rev.Phytopathol., 5 : 131-162.
- ✓ *Wen, T.C. (1983). Influence of heavy metal pollution on the growth, yield and quality of tobacco plant. IV. Studies on absorption and balance of copper-zinc cadmium in relation to pollution hazards to leaf tobacco. Bull.Tob.Res.Inst. Taiwan-Tobacco and Wine Monopoly. Bureau, 19 : 21-23.

- Whitney, P.J. (1972). The carbohydrate and water balance of beans (Vicia faba) attacked by broomrape (Orobanche crenata). Ann. Appl. Biol., 70 : 59-66.
- ✓ Whitney, P.J. (1973). Transport across the region of fusion between bean (Vicia faba) and brcomrape (Orobanche crenata) Proc. Eur. Weed. Res. Counc. Symp. Parasit. Weeds, pp.154-166.
- ✓ Wolswinkel, P. (1978). Accumulation of phloem mobile mineral elements at the site of attachment of Cuscuta europea L. Z. Pflanzenphysiol., 86(1) : 77-84.
- Wutscher, R.K. and Hardesty, C. (1979). Zinc and phenolics in the wood and incidence of citrus blight in a grove containing eight selections of 'Hamlin orange'. Proc. Fla. State Hortic. Soc., 92 : 70-72.
- ✓ Yarwood, C.E. and Jacobson, L. (1955). Accumulation of chemicals in diseased areas of leaves. Phytopathol., 45 : 43-48.
- *Yorganci, U. and Serin, S. (1986). Effects of viruses on the yield output, quality and chemical composition of tobacco. Journal of Turkish Phytopath., 15(1) : 1-19.
- ✓ Zuber, M. and Manibhushanrao, K. (1983). Physiology of host-parasite interaction in rice sheath blight disease with reference to changes in carbohydrate and nitrogen contents. Indian J. Bot., 6(2) : 117-124.

* Original not seen.