BIBLIOGRAPHY

Aparicio,P.J., Cardenas,J., Zumft,W.G., Vega,J.Ma., Harresa,J., Peneque,A. and Losada,M. (1971). Molybdenum and iron as constituents of the enzymes of the nitrite reducing system from Chlorella. Phytochemistry, 10 : 1487-1495.

Arnon,D.T. (1949). Copper enzymes in isolated chloroplasts : Polyphenol oxidase in Beta vulgaris. Plant Physiol., 24 : 1-15.

Avent,K.L. (1957). Some apple root stocks and potassium. J.Aust.Inst.Agric.Sci., 23 : 154-157.

Bates,L.S., Waldren,R.P. and Teare,I.D. (1973). Rapid determination of free proline for water stressed studies. Plant and Soil, 39 : 205-207.

Beevers,L. and Hageman,R.H. (1969). Nitrate reduction in higher plants. Ann.Rev.Plant Physiol., 20 : 495-522.

Begum,A. and Pereira,S.M.(1977). The Beta carotene content of Indian edible green leaves. Trop.Geogr.Med., 29 : 47-50.

Bhivare,V.N. and Nimbalkar,J.D. (1984). Biochemical changes leading to senescence in bean leaf. Biovigyanum, 10 : 31-34.

Bjorkman,E. (1953). Om "Granes gulspetssjuka" i plantskolor SV. Skogsvardsfoeren. Tidskr. 51 : 211-229.

Blahova,M. and Segata,V.(1980). Nitrate reductase activity in the course of cucumber leaf ontogenesis. Biol.Plant., 22(3) : 176-182.

Boggess,S.F., Stewart,C.R., Aspinall,D. and Paleg,L.G. (1976). Effect of water stress on proline synthesis from radioactive precursor. Plant Physiol., 58 : 398-401.

Bollard,E.G. (1958). Nitrogenous compounds in tree xylem sap. In 'The Physiology of Forest Trees' (K.V.Thimann, ed.). pp. 83-93, Ronald, New York.

Cavalieri,A.J. and Huang,A.H.C. (1979). Evaluation of proline accumulation acompanying necrosis of tobacco leaves systematically infected with tobacco mosaic virus. Phytopathology, 59 : 953-962.

Champion,H.G. and Seth,S.K. (1963). Indian silviculture, Govt.of India, New Delhi.

Chu,T.M., Aspinall,D. and Paleg,L.G. (1974). Stress metabolism. VI. Temperature stress and the accumulation of proline in barley leaves. Aust.J.Plant Physiol., 1 : 87-97.

Day,W.R. (1946). The pathology of beech on chalk soils. Quart.J.Forest., 40 : 72-82.

Dijkman,M.J. (1950). Leucaena a promising soil erosion control plant. Econ.Bot., 4 : 337-349.

Doddema,H., Stulen,I. and Hofstra,J.J. (1986). The distribution of nitrate reductase in tomato (L.e.) leaves as affected by age. Physiol.Plant., 68 : 615-619.

Evans,J. (1982). Nitrate reductase activity in chick-pea : A preliminary investigation. International chickpea News-letter., 6 : 22-23.

Folin,O. and Denis,W. (1915). A colorimetric estimation of phenols (and phenol derivatives) in urine. J.Biol.Chem., 22 : 305-308.

Gokhale,S.P., Karadge,B.A. and Rajmane,N.A. (1984). Biochemical changes during leaf development of Catharanthus roseus. Biovigyanum, 10 : 25-30.

Gregory,F.G.and Veale,J.A. (1957). A reassessment of the problem of apical dominance. Symp.Soc.Exp.Biol., 11:2-20.

Hardwick,R.C. (1983). Interactions between nitrogen and growth regulators in the control of plant development. "Why do seed legumes lose their leaves ?". British Plant Growth Regulator Group, Monograph, 9 : 61-73.

Haq,Q.N., Moillan,N.I. and Kiamuldin,M. (1974). Studies on polysaccharides of the bark of Litsaea chinesis Lamk. Bangladesh J.Sciindres., 9(3/4) : 95-100.

Hegde,B.A. and Karande,S.M. (1978). Effect of presowing treatment of sodium chloride on the incidence of green ear disease of Pennisetum typhoides (Burm) Stapf. and Hubb var. HB_3. Plant and Soil, 49(3) : 551-559.

Hepting,G.H. (1945). Reserve food storage in short leaf pine in relation to little leaf disease. Phytopathology, 35 : 106-119.

Hillis,W.E. ed. (1962). "Wood Extractives and their significance to the pulp and paper industries". Academic Press, New York.

Hillis,W.E. (1968). Chemical aspects of heart wood formation. Wood Sci.Technol., 2 : 241-259.

Holden,M. (1973). Chloroplast pigments in plant with the C_4 dicarboxylic acid pathway of photosynthesis. Photosynthetica, 7(1) : 41-49.

Jacobson,A. and Corcoran,M.R. (1977). Tannins as gibberellin antagonists in the synthesis of α- amylase and acid phosphatase by barley seeds. Plant Physiol., 59 : 129-133.

Jamale,B.B. and Joshi,G.V. (1976). Physiological studies in senescent leaves of mangroves. Indian J.Exp.Biol., 14 : 697-699.

Jamale,B.B. and Joshi,G.V. (1978). Effect of age on mineral constituents, polyphenols, polyphenol-oxidase and peroxidase in leaves of mangroves. Ind.J.Exp.Biol., 16(1) : 117-119.

Joshi,G.V. and Mishra,S.D. (1970). Photosynthesis and mineral metabolism in senescent leaves of Clerodendron inerme, Gaertn. Indian J.Exp.Biol., 8 : 41-43.

Judel,G.K. (1972). Peroxidase and catalase activity and the content of total phenols in the leaves of sunflowers as affected by copper and nitrogen deficiency. Z.Pflanzenernacher. Bodenke, 133 : 81-92.

Kalela,E.K. (1950). Mannikoiden fa Kuusikoiden juurisuhceista it. Acta.Forest.Fenn., 57 : 1-68.

Kannaiyan,J., Vidhyasekaran,P. and Kandaswamy,T.K. (1973-b). Amino acid contents of bajara in relation to ergot disease resistance. Indian Phytopath., 25 : 358-359.

Kar,M. and Mishra,D. (1976). Catalase, peroxidase and polyphenol oxidase activities during rice leaf senescence. Plant physiol., 57 : 315-319.

Karadge,B.A. (1981). Physiological Studies in succulents. A Ph.D.Thesis submitted to Shivaji University, Kolhapur (India).

Kira,T. (1975). Primary productivity of forests. In. "Photosynthesis and Productivity in Different Environments" (J.P. Cooper, ed.), IBP, Vol.3, pp.5-40. Cambridge Univ. Press, London and New York.

Kirk,J.O.T. and Allen,R.L. (1965). Dependence of chloroplast pigment synthesis on protein synthesis : Effect of actidione. Arch.Biochem.Biophys.Res.Commun., 21 : 523-530.

Koyama,H. (1978). Photosynthesis studies in Pasoh Forest. Malay.Nat.J., 30 : 253-258.

Kozlowski,T.T. and Winget,C.H. (1964). The role of reserves in leaves, branches, stems, and roots on shoot growth of red pine. Amer.J.Bot., 51 : 522-529.

Kozlowski,T.T. and Clasusen,J.J. (1966). Shoot growth characteristics of heterophyllous woody plants. Can.J.Bot., 44 : 827-843.

Kramer,P.J. and Kozlowski,T.T. (1960). 'Physiology of tree'. Mc Graw-Hill, New York.

Larcher,W. (1980). 'Physiological Plant Ecology'. Revised 2nd edition. Springer-Verlag, Berlin.

Leopold,A.C. (1961). Senescence in plant development. Science, 134 : 1727-1732.

Leopold,A.C. and Kriedemann,P.E. (1975). 'Plant Growth and Development'. McGraw Hill, New York.

Lianen-Jensen,S. and Jensen,A. (1971). Quantitative determination of carotenoids in photosynthetic tissues. In. "Methods in Enzymology", (A.Sen Pietro, Ed.) : Academic Press, Inc. Publishers, New York, pp.586-602.

Li,P.H., Weiser,C.J. and Van Juystee,R.B. (1965). Changes in metabolites of red-osier,dogwood during cold acclimation. Proc.Am.Soc.Hortic.Sci., 86 : 723-730.

Lowry,O.H., Rosebrough,N.J., Farr,A.L. and Randall,R.J. (1950). Protein measurement with the folin phenol reagent. J.Biol. Chem., 193 : 262-275.

Lunderstadt,J. (1976). Isolation and analysis of plant phenolics from foliage in relation to species characterization and to resistance against insects and pathogens. In. "Modern Methods in Forest Genetics" (J.P.Miksche, ed.), pp.158-164. Springer-Verlag, Berlin and New York.

Mayer,A.M. and Poljakoff-Mayber, A. (1963). "The Germination of seeds". MacMillan, New York.

Meyer,M.M. and Tukey,H.B. Jr. (1965). Nitrogen, phosphorus and potassium plant reserves and the spring growth of Taxus and Forsythia. Proc.Amer.Soc.Hort.Sci., 87 : 537-544.

Miflin,B.J. (1974). The location of Nitrate reductase and other enzymes related to amino acid biosynthesis in the plastids of root and leaves. Plant Physiol., 54(4) : 550-555.

Moller,C.M., Muller,D. and Nielsen,J. (1954). Graphic representation of dry matter production of European beech. Def. Forestl.Forsogsv.Danmark., 21 : 327-335.

Monteith,J.L. (1978). Reassessment of maximum growth rates for C_3 and C_4 crops. Expt.Agric., 14 : 1-5.

Murneek,A.E. (1942). Quantitative distribution of nitrogen and carbohydrates in apple trees. Mo.,Agric.Exp.Stn.,Res.Bull., 348.

Neish,A.C. (1960). Ann.Rev.Plant Physiol., 11 : 55 (1960). In. Formation of wood in forest trees. (M.H.Zimmerman, Ed.) p.219. Academic Press, New York, 1964.

Nelson,N. (1944). A photometric adaptation of Somogyi method for determination of glucose. J.Biol.Chem., 153 : 375-380.

Neyra,C.A. and Hageman,R.H. (1974). Dependence of nitrate reduction on electron transport in chloroplasts. Plant Physiol., 54(4) : 480-483.

Notton,B.A. and Hewitt,E.J. (1971). Incorporation of radioactive molybdenum into protein during nitrate reductase formation and effect of Molybdenum on nitrate reductase and diaphorase activities of spinach (Spinacia oleracea L.) Pl.Cell Physiol., Tokyo, 12 : 465-477.

Osborne,D.J. (1962). Effect of kinetin on protein and nucleic acid metabolism in Xanthium leaves during senescence. Plant Physiol., 37 : 595-602.

Ovington,J.D.(1962). Quantitative ecology and the woodland ecosystem concept. Adv.Ecol.Res., 1 : 103-192.

Owusu,J.K., Adomako,D. and Hutcheon,W.V. (1978). Seasonal changes in total free sugar content of field cocoa plants. Physiol.Plant., 44(1) : 43-47.

Pate,J.S. (1968). Inorganic and intermediate nitrogen metabolism. In. "Recent Aspects of Nitrogen Metabolism in Plants". (J.E.Hewit and C.V.Cutting, Eds.). pp.219-240. Academic Press, London.

Richardson,S.D. (1957). The effect of leaf age on the rate of photosynthesis in detached leaves of tree seedlings. Acta Bot.Neer., 6 : 445-457.

Rivas Nilo,R. and Carmen De M. (1979). Physical-chemical characterization of sapota fruit (Achras zapota) of the state of Zulia and study of its adaptibility to canning. Rev.Fac. Agron. (Maracay), 10(1-4) : 57-69.

Sable,A.B. (1983). Studies on photosynthesis in plants. Ph.D. thesis submitted to Shivaji University, Kolhapur (India).

Sacher,J.A. (1957). Relationship between auxin and membrane integrity in tissue senescence and abscission. Science, 125 : 1199-1200.

Scholtes,W.H. (1953). The concentration of forest tree roots in the surface zone of some piedmont soils. Proc.Iowa Acad.Sci., 60 : 243-259.

Schulman,E. (1958). Bristelecone pine, oldest known living thing. Nat.Geogr.Mag., 113 : 355-372.

Sestak,Z. and Catsky,J. (1962). Itensity of photosynthesis and chlorophyll content as related to leaf age in Nicotiana sanderae Hort. Biol.Plant., 4 :131-140.

Sestak,Z. (1963 a). Changes in the chlorophyll content as related to photosynthetic activity and age of leaves. Photochem. Photobiol., 2 : 101-110.

Shaner,D.L. and Boyer,J.S. (1976). Nitrate reductase activity in maize (Zea mays L.) leaves. I. Regulation by nitrate flux. Plant Physiol., 58 : 499-504.

Shetty,G.P. (1971). Physiology of growth and salt tolerance of plants. A Ph.D. thesis submitted to the Shivaji University, Kolhapur (India).

Sieckmann,S. and Bot.A.A. (1978). Low temperature increases reducing and total sugar concentrations in leaves of box wood (Buxus sempervirens L.) and cranberry (Vaccinium macrocarpon Ait.). Hort.Science, 13(4) : 439-440.

Siminovitch,D. and Briggs,D.R. (1949). The chemistry of the living bark of the black locust tree in relation to frost hardiness. I. Seasonal variations in protein content. Arch.Biochem., 23 : 8-17.

Sivaprakasam,K., Vidhyasekaran,D. and Soumini Rajagopalan,C.K. (1974). Physiology of brinjal plants infected with Verticillium dahliae. Indian Phytopath., 27 : 197-202.

Stone,E.L. (1968). Microelement nutrition of forest trees : A review, In. "Forest Fertilization - Theory and Practice". pp. 132-175. T.V.A., Muscles Shoals, Alabama.

Takashashi,M. and Ripperton,J.C. (1949). Kaohaole (LEUCAENA LGAUCA (L.) Benth.). Its establishment, culture and utilization as a forage crop. Hawaii Agric.Expt.Stn.Bull. 100:56.

Thimann,K.V. (1972). The natural plant hormones. In. "Plant Physiology" Vol.6B, (F.C.Steward, ed.) pp.3-145. Academic Press, New York.

SHIVAJI UNIVERSITY LIBRARY

Thombre,R.R. (1987). Studies in Physiology of leaf ontogeny in plants. Ph.D. Thesis submitted to Shivaji University, Kolhapur (India).

Troup,R.S. (1921). Silviculture of Indian Trees. Cf.Sagreiya, K.P. In. "Forests and Forestry" pp.214. (1967). National Book Trust, India.

Tukey,H.B. Jr. (1970). The leaching of substances from plants. Annu.Rev.Plant Physiol., 21 : 305-324.

Vega,J.M., Cardenas,J. and Losada,M. (1980). Ferredoxin-nitrate reductase. In. : Methods in Enzymology, 69 : 255-270.

Verner,J.E. (1961). Biochemistry of senescence. Annu.Rev.Plant Physiol., 12 : 245-264.

Vidhyasekaran,P. (1972). Studies in heminthosporiose of ragi in relation to disease resistance. Ph.D. Thesis. Tamilnadu Agric.University.

Wallace,J.W. and Mansell,R.L. eds. (1975). Biochemical Interactions between plants and insects. Plenum, New York.

Walker,L.C. (1956). Foliage symptoms as indicators of potassium deficient soils. Forest Sci., 2 : 113-120.

Walter,H. (1955). The water economy and hydrature of plants. Ann.Rev.Plant Physiol., 6 : 239-252.

Wareing,P.E., Khalifa,M.M. and Treharne,K.J. (1968). Rate limiting processes in photosynthesis at saturating light intensities. Nature (London), 220 : 453-457.

Wenger,K.F. (1953 a). The sprouting of sweetgum in relation to season of cutting and carbohydrate content. Plant Physiol., 28 : 35-49.

Went,F.W. (1942). Some physiological factors in the aging of a tree. Proc.18th Nat.Shade Tree Conf., pp. 330-334.

Whitfield,A.B. (1963). The effect of stock and scion on the mineral composition of apple leaves. Rep.E.Malling Res. Stn., 107-109.

Whittaker,R.H. (1970). The biochemical ecology of higher plants. In."Chemical Ecology" (E.Sondheimer and J.B. Simeone, eds.), pp.43-70. Academic Press, New York.

Wieckowski,S. (1958). Studies on the autumnal breakdown of chlorophyll. Acta.Biol.Cracov., Ser.Bot., 1 : 131-135.

Wolf,F.T. (1956). Changes in chlorophylls 'a' and 'b' in autumn leaves. Am.J.Bot., 43 : 714-718.

Ziegler,H. (1964). Storage, mobilization and distribution of reserve material in trees. In. "The formation of wood in Forest Trees". (CM. H.Zimmermann, Ed.). pp.303-320. Academic Press, New York.