
BIBLIOGRAPHY

REFERENCES

Agarwal, 1986. Economic plants of India.

Agharkar S. P. 1953. Gazetter of Bombay State Volume : General A. Botany Part-I Medicinal plants Executive
Editor and Secretary, Bombay.

Allgeir H. 1968. Structure of pachybiose (from root of Pachycarpus lineolatus) & asclepibiose (from root of Asclepias lilacina & seeds of Dregea volubilis & D. abyssinica).
Helv. Chim. Acta. 51 : 311-325.

Almedia S.M. 1989. Some rare endangered and threatened plant species from Ratnagiri district, Maharashtra
Jour. Bomb. nat. Hist. Soc. 86 : 478-479.

Ansari M. Y., 1968. A new species of Ceropegia Linn.
(Asclepiadaceae) from western ghats, Maharashtra.
Bull. Bot. Surv. India 10 : 95-97.

Ansari M. Y. 1969. Ceropegia media (Huber) Ansari Stat. Nov.
from Western Ghats. (Maharashtra.)
Bull Bot. Surv. India 11 : 199-201.

- Ansari, M. Y. 1971. Ceropegia vincaeefolia, Hook. (Asclepiadaceae) from Maharashtra-It's history and identity.
Bull. Bot. Surv. India 13 : 187-191.
- Ansari, M. Y. 1972. New Ceropegia, Linn (Asclepiadaceae) From Sahayadri Range in Maharashtra State.
J. Bomb. nat. Hist. Soc. 69 : 250-253.
- Ansari, M. Y., Kulkarni B. G. 1971. Ceropegia sahyadrica Ansari et. Kulkarni - A new species of Asclepiadaceae from Sahayadric range in Maharashtra State
Indian Forester 97 : 686-690.
- Ansari, M. Y. and Kulkarni B. G., 1980. A new species of Ceropegia Linn. (Asclepiadaceae) from the Western Ghat in Maharashtra state, India.
Bull. Bot. Surv. India 22 : 1-4.
- Ansari M. Y. 1980. Ceropegia maccannii a new species (Asclepiadaceae)
Bull. Bot. Surv. India 22 : 1-4.
- Ansari, M. Y. 1984. Fascicles of Flora of India
Bull. Bot. Surv. India 16 : 1-34.
- Arekal G. D. 1981. A new species of Brachystelma (Asclepiadaceae) from India.
Curr. Sci. 50 : 145.
- Arekal G. D. and Ramakrishna T. M. 1981. A new Brachystelma (Asclepiadaceae) from Kolar, Karnataka.
Proc. Indian Acad. Sci. Sci. 90 : 203-205.

- Arekal, G. D. and Ramakrishna, T. M. 1981. Extraloral nectaries of allotropies gigantea & Waittakaka Volubilis
Phytomorphology 36 303-306.
- Banerji, M. L. 1974. Petiole anatomy and minor viens as aids for the identification of Symplocos of Nepal.
Bull. Bot. Surv. India 16 : 89-100.
- Beri, R. M. and Sharma O. P. 1963. Chemical constituents of Sarcostemma acidum (Roxb.) Voight.
Indian J. Chem. 1 : 501.
- Bhatnagar, S. 1975. Floral polymorphism in sympatric population of Calotropis procera (Ait) R. Br. Acta Bot. Indica 3 : 43-46.
- Biswas, I. 1957. Embryological studies in Daemia extensa R. Br. J. Indian Bot. Soc. 36 : 207-222.
- Blatter, E and McCann, C. 1931. A new Ceropegia from the Western Ghats. Jour. Bomb. nat. Hist. Soc. 34 1936.
- Blatter, E and McCann, C. 1931. Another Ceropegia from the Western Ghats. Ibid 35 : 409.
- Blatter, E. and McCann, C. 1933. Revision of the flora of the Bombay Presidency (Asclepiad). Ibid 36 : 524-537.

Blatter, H. W. 1945. Anatomy of Cryptostegia grandiflora with special reference to the latex system. Amer. J. Bot. 32 :135-41.

Bolkovshish, Z., Grif, V. 1969. Chromosome number of flowering plants. Acad. Sci. USSR : 59.

Bookman, S. S. 1981. The floral morphology of Asclepias speciosa (Asclepiadaceae) in relation to pollination and a clarification in terminology for the genus. Amer. J. Bot. 68 : 575-679.

Borrl, M. 1961. Epidermal characteristic in the diploid sub-species of Dactylis glomerata L. J. Linn. Soc. London (Bot.). 56 : 453-458.

Brown, R. 1883. On the organs and mode of fecundation in orchidaceae and Asclepiadaceae. Trans. Linn Soc. Land. 13 :201-34.

Bruyns, P. 1985. The genus Ceropegia on the Canary Islands. (Asclepiadaceae-Ceropegieae) a morphological and taxonomic account. Beitr. Biot. Pflanzen 60 : 427-458.

Bruyns, P. V. 1985. Pollinaria and Ceropgia arabica and allies. Asclepias 36 : 65-67.

Bruyns, P. 1986. The genus Ceropegia on the Canary Island (Spain) (Asclapiadaceae-Ceropegieae) : A morphological and anatomical account. Beitr. Biot. Pflanzen. 60 : 427-458.

Chandra, V; Kapoor, S. L.; Sharma P. C. and Kapoor L. D. 1969. Epidermal and venation studies in Apocynaceae-I. Bull. Bot. Surv. India 11 : 286-289.

Chandra, V.; Mitra R.; Kapoor, S. L. and Kapoor L. D. 1972. Epidermal and venation studies in Apocynaceae-IV. Ibid. 14 : 76-82.

Char, 1978. A new species of Brachystelma R. Br. (Asclepiadaceae) from India. Curr. Soc. 17 : 965-966.

Chaturvedi, S. K. 1987. Pollination and pollen germination in Cynanchum canescens (Wild) Schum. Asklepias 40 : 93-96.

Chaturvedi, S. K. 1989. Abiotic pollination in Tylophora hirsuta, Weight. (Asclepiadaceae). Asklepias 45 : 58-62.

Chaturvedi, S. K. and Pant D. D. 1986. Further studies in the pollination of some Indian Asclepiads. Bull. Bot. Surv. India 28 : 23-30.

Chandra, R. N. and Chopra, I.C. 1958. Indigenous Drugs of India Dhur and Sons Pvt. Ltd., Calcutta-12.

Chopra et. al. 1956. Glossary of Indian Medicinal Plants, C.S.I.R.,

New Delhi : 222.

Christ P. and Schnepf E. 1886. The nectaries of Cynanchum vincetoxicum (Asclepiadaceae) ISR J. Bot. 34 : 79-90.

Cooke, T. 1904. The Flora of the Presidency of Bombay.

Colonval-Elenkov et. Malaisse, F. 1984. Recherches Sur Les Asclepiadaceae Du Shaba (Zaire)
Note. 2 : Apports et limites de l'anatomie follaire dans la systematique des Ceropegia L. Bull. Soc. Roy. Bot. Belg. 117 : 135-142.

Corry, T. H. 1883. On the structure and development of the gynostegium and the mode of fertilization in Asclepias cornutti Decaisne (A. syviaca) Trans. Linn. Soc. Ser. II Bot. 2 : 173-207.

Corry, T. H. 1884. On the structure and development of gynostegium and the mode of fertilization in Asclepias cornuti, Dcaine. Trans. Linn. Soc. Bot. 2 : 173-207.

Crete, P. 1950. Embryogeny. des Asclepiadaceæ, Dévelopement de l'embryon Asclepias curassavica C.R. Acad. Sci. Paris 230 : 172-173.

Cronquist, A. 1981. The evolution and classification of flowering plants. Hazell Watson & Viney Ltd. Aylesbury, Bucks.

Dassana M. D. and Jaysuriya A.H.M. 1975. A New species of Brachystelma (Asclezia) from Sri Lanka.
Ceylon J. Sci. Biol. Sci. 11 : 39-41.

Davis, G.L. 1966. Systematic embryology of the angiosperm.

John Wiley & Sons. INC.

Decker, J.M. 1967. Petiole vascularisation of Luxemburghiaeae (Ochnaceae)
Amer. J. Bot. 54 : 1175-1181.

Deshpande B. D. and Premlata Joneja, 1962. Morphology & embryology of Leptadenia Pyrotechnia Dcne. Phyton
19 : 73-84.

Devi, H. M. 1964. Embryological studies in Asclepiadaceae

Proc. Indian Acad. Sci. 60 : 54-65.

Devi, H. M. and Lakshminarayana K. 1979. Embryology of Oxystelma esculentum.
Phytomorphology. 27 : 59-67.

Dnyansagar, V. R. and Tijiar, V. R. 1979. Pollinia and pollination in Calotropis phytia. 23 : 97-106.

Dunber, et. al. 1986. Identification of laticifers in embryoides derived from callus. and suspension cultures of Asclezia species (Asclepiadaceae)
Am. J. Bot. 73 : 847-851.

Dyer, R. A. 1977. New species of Brachystelma (Asclepiadaceae)
Bothalia 12 : 55-57.

Dyer, R. A. 1977. New records of Brachystelma J. S. AFR 43 : 9-20.

Dyer, R. A. 1977. Asclepiadaceae : Two new species of Brachystelma
Bothlia 12 : 497-448.

Dyer, R.A. 1978. Asclepiadaceae : The identity of distribution of
Ceropegia zeyheri
Bothalia 12 : 446.

Dyer, R.A. 1978. Asclepiadaceae : New species of Ceropegia
Bothalia. 12 : 444-445.

Edwards, W. H. 1935. The systematic value of cuticular characters
in recent and fossil angiosperms.
Biol. Rev. 10 : 442-459.

Finn, W. W. 1925. Male cells in Angiosperms. I. spermatogenesis and
fertilization in Asclepias cornuta. Bot. Gaz.
80 : 1-25.

Francke, A. 1927. Zur kenntnis der. Exodermis der Asclepiadaceen.
Planta 3 : 1-26.

Frye, T. C. 1902. A morphological study of certain Asclepiadaceae
Bot. Gaz. 34 : 389-413.

Gaikwad D.K., Supate A.R., Yadave S.R., and Chavan P.D. 1989.
Occurrence of Crassulacean Acid Metabolism
in stem tissue of Ceropegia juncea, Roxb.
Photosynthetica 23 : 216-220.

Gujer, S.C. 1902. The development of the pollinium and sperm cells
in Asclepias cornuti

Ann. Bot. Lond. 16 : 123-148.

Galil J. and Zeroni M. 1965. Nectar system of Asclepias curassavica
Bot. Gaz. 126 : 144-148.

Gamble, J.S. 1923. The Flora of the Presidency of Madras

Gazzar, E. L. et. al. 1974. Pollen morphology and taxonomy of
Asclepiadaceae.

Pollen et. Spores 16 : 227-238.

Glabisz, J. 1908. Morphologische und physiologisch Untersuchungen on
Ceropegia woodii Schlechter.

Beih. Bot. Zbl. 23 : 65-136.

Gupta, R.C. 1985. Pharmacognostical studies on 'Jivanti' part IV -
Dregea volubilis (Linn F.) Benth.

Bull. Bot. Surv. India 27 41-47.

Gupta, R.C. Ansari M.S. and Kapoor L.D. 1970. Pharmacognostical
studies on Jivanti Part-I Desmotrichum
ibriatum Blume.

Bull. Bot. Surv. India 12 : 29-36.

Gupta, R.C. and Kapoor L.D. 1971. Pharmacognostical studies on Jivanti
Part II- Leptodenia reticulata wt. & Arn.
Csyn. Gymnema auranticum wall ex. Hook
F. and Asclepias tuberosa Roxb.
Ibid 13 : 53-63.

Gupta, R.C. and Wahl, A.K. 1978. Sarcostemma brevistigma W & A
 Ibid. 20 :103-112.

Hajarnavis, L.S. 1964. Free organic acids in some xerophytes
Curr. Sci. 33 : 584-585.

Handa, T. 1936. Anatomical observations on the internal cambium of
 the stem in Marsdenia tomentosa, Merr.
 et. Decne.
Jap. J. Bot. 8 : 59-64.

Harold, K. 1985. The worth of pollinarium
Asklepios 35 : 29.

Hassan, W.E. Youngken, H.W. and Quimby. M. W. 1951 Studies on
 the sps. of Asclepias I - VI Introduction
 and comparative study of the roots. J.
Amer. Pharm. Assoc. 41 : 6-8, 9, 86,
 320, 298.

Hassan W.E. R.A., Gosselin J.R., Vanderwyk, R.W. and Quimby M.W.
 1952 . Studies on the species of Asclepias
 VII. Determination of palisade ratio. J.
Amer. Pharm. Assoc. 42 : 666.

Hemadri, K. 1968. A new Ceropegia Linn. (Asclepiadaceae) from Western
 ghats, Maharashtra
Bull. Bot. Surv. India 10 : 123-125.

Hemadri, K. and Ansari, M. Y. 1971. Ceropegia mahabale A new species of Asclepiadaceae from Sahyadri range, Western Ghats.

Indian Forester 97 : 105-108.

Hildebrand, F. 1866. Über die Befruchtung von Asclepias cornuti.

Bot. Ztg. 24 : 376-378.

Hooker, J. D. 1883. The flora of British India.

Howard, R.A. 1962. The vascular structure of the petioles as a taxonomic character and their application
Adv. Horticultural Sci. III : 7-13.

Huber, H. 1957. Revision der Gattung Ceropegia

Mem. Soc. Broteriana 12 : 1-203.

Ito, et. al. 1978. Studies on the constituents of Marsdenia formosana Masumune. III. Isolation and structural elucidation of some new steroidal glycosides

Chem. Pharm. Bull. Tokyo 26. :3189-3194.

Jain, D.K. and Singh, V. 1974. Epidermal studies in Rosaceae-I.

Pyrus. Geobios L : 101-104.

Johansen, D.A. 1940. Plant microtechnique. McGraw Hill Book Co., Inc.

New York & London.

Kanodia, K.C. and Reddy B.V. 1964. Ceropegia fantastica
 (Asclepiadaceae) an imperfectly known species.

Bull. Bot. Surv. India 6 : 311-312.

Kapoor, S.L. Sharma, P.C. and Kapoor, L.D. 1969. Epidermal and venation studies in Apocynaceae-II,
Ibid 11 : 372-376.

Kapoor, S.L. Verma, C.L. and Trivedi, B.S. 1985. A study of the leaf cuticle of Indian clematis (Dill.ex.) (Ranunculaceae)
Ibid. 27 : 197-218.

Kirtikar and Basu. 1975. Indian medicinal plants. Vol-II.

Kunth, P. 1908. Handbook of flower pollination. Vol.I-III.

Transl. Davis J.R.A. Clarendon Press
Oxford.

Kupchan, S.M. John, R. Knox and Kelesy, J.E. 1964. Calotropin, a cytotoxic principle isolated from A. curassuvica L. Science : 146.

Krishnamurthy, K.H. and Sundaram. 1967. Foliar epidermis and Pharmacognosy in some members of Asclepiadaceae.

J. Indian Bot. Soc. 46 : 160-168.

Krishnamurthy and Kannabiran, 1970. Histo-morphophiology of foliar epidermis and pharmacognosy in Asclepiadaceae.

Ibid. 49 : 105-114.

Lange, O. and Zuber, M. 1978. Frerea indica, a stem succulent CAM plant with deciduous C₃ leaves.

O. Ecologia (BER) 31 : 67-72.

Lakshminarayana and Krishnappa D.G. 1988. Cytomorphological studies in a few species of the genus Ceropegia L Abstracts No. 184, Page. 100

Second All India Conference on Cytology and Genetics Warangal.

Lakshminarayana and Krishnappa D.G. 1990. Karyomorphological studies in some members of the South Indian Asclepiadaceae. Abstracts No. 44. Page.27
Third All India Conference on Cytology and Genetics Nagpur.

Lems, K. 1964. Evolutionary studies in the Ericaceae II Leaf anatomy as a phylogenetic index in the Andromedaeae
Bot Gaz. 125 : 178-186.

Lendner, A. 1924. L'anatomie due Solenostemma Arghel. Hayne.
Schweiz Apothzg. 62 : 9-13.

Lucansky, T.W. and Clough, K.T. 1986. Comparative anatomy and morphology of Asclepias perennis of A. tuberosa spp. rolfssi.
Bot. Gaz. 147 : 290-301.

Macior, L.W. 1965. Insect adaptation and behaviour in Asclepias pollination
Bull. Torry. Bot. Club. 92 : 114-126.

McCann, C. 1943. Light windows in certain flowers.

Journ. Bomb. Nat. Hist. Soc. 44 :182-184.

McCann, C. 1945. New species of Ceropegia and the synonymy of the Indian species.

J. Bombay. Nat. Hist. Soc. 45 : 209-211.

Malaisse, F. 1984. Recherches sur les Asclepiadaceae du Shaba (Zaire)
Note : I : Nouvelles observations sur le genre Ceropgia Bull. Jard. Bot. Nat. Belg. 54 :213-234.

Milburn, T.R. et.al. 1968. Crassulacean Acid Metabolism under natural tropical conditions.

New Phytol 67 : 883-897.

Mitra R., Mehrotra S., Mehrotra B.N. and Kapoor L.D. 1974.
Pharmacognostic study of Asclepias curassavica Linn.
Bull. Bot. Surv. India 16 82-88.

Mitra R., Kapoor S.L. and Kapoor L.D. 1978. Epidermal and venation studies on Apocynaceae-V.

Ibid 20 : 20-30.

Muller, F. 1977. Flowers and insects.

Nature 17 : 17.

Muller, H. 1883. The Fertilization of flowers

Trans. D' Arcy W. Thompson, London.

Mulay, B.N., Deshpande and Tolani U. 1965. Studies in Asclepiadaceae II Floral morphology & gametogenesis in certain members of the Asclepiadaceae.

J. Indian Bot. Soc. 44 :95-104.

Newton, L.E. 1980. Phytoogeographical Associations of the succulent plant flora of S.W. Arabia and the Horn of Africa.

Nat. Cact. & Succ. J. 35 : 83-88.

Nirula, R.L. 1954. On the development of the embryosac and the endosperm in Daemia extensa

Proc. Indian Acad. Sci. 21 :181-185.

Nuernbergk, E.L. 1961. Endogener Rhythmus und CO₂ Stoffwechsel bei Pflanzen mit diurnalem Sauerhythmus.

Planta 56 : 28-70.

Pant D.D. and Chaturvedi S.K. 1986. Pollination ecology of Asclepias curassavica L
Geophytology 16 :119-121.

Pant, D.D. Nautiyal D.D. and Chaturvedi S.K. 1982. Polllination ecology of some Indian Asclepiads.
Phytomorphology 32 :302-313.

Patnaik, G.K. and Dhawan, B.N. 1971. 'Asclepin' a new cardioactive glycoside from Asclepias currassavica
Ind. J. Pharm. 3 : 8.

Patwardhan, U.R. 1953. Embryological studies in the Asclepiadaceae-I
The female gametophytes
J. Univ. Poona 3 :11-18.

Percival, M.S. 1969. Floral Biology. 2nd ed. Pergaman ^OPress, Oxford.

Prat, H. 1960. Vers. une classification naturelles des graminees.
Bull. Soc. Bot. Fr. 107 :32-79.

Proctor, M. and Yeo, P. 1973. The pollination of flowers Collins.
St. J. Place London.

Puech, M. G. 1912. Etude anatomique de quelques especes d'
Asclepiadees aphyllles de l' ouest de
Madagascar Rev. Gen. Bot. 24 : 298-312
and 329-343.

Raghvan, R.S. and Ansary, M.Y. 1975. Chromosome number in the genus
Ceropegia Linn.
Curr. Sci. 44 : 863-864.

Ramayya, N. 1969. The development of trichomes in the compositae.

In Recent advances in the anatomy of tropical seed plants.

Hindustan Publishing Corporation, Delhi.

Ramayya, N. 1972. Classification and phylogeny of the trichomes of Angiosperms.

In Research Trends in Plant Anatomy. K.A. Chowdhury commemoration volume.

Tata McGraw Hill Publishing Company Ltd., New Delhi.

Ramayya, N. and Rajgopal, T. 1971. Foliar "Dermotypes" of the Indian Aizoaceae and their use in identification.

J. Indian Bot. Soc. 50 : 355-362.

Ramkrishna, T. and Govindappa, D. 1979. Pollination Biology of Calotropis gigantea (L)

R. Br. Curr. Sci. 48 :212.

Rao, C.V. and Rao, S.R. 1954. Embryology of Cryptostegia grandiflora and Caralluma attenuata J. Indian Bot. Soc. 33 : 453-472.

Rao, V. and Arati Ganguli. 1963. The floral anatomy of some Asclepiadaceae.

Proc. Indian Acad. Sci. 57 :15-44.

Reddi, B.V. 1964. Ceropegia fantastica sedwick (Asclepiadaceae) An imperfectly known species.

Bull. Bot. Surv. India 6 : 311-312.

Rintz, R.E. 1979. Three new species of Asclepiadaceae from peninusular Malaya. 17374 Millar Road, Mt. Clemens Mich 48043, U.S.A.

Sabet, Y.S. 1931. Development of embryo-sac in Calotropis procera with special reference to the endosperm formation.

Ann. Bot. Lond. 45 : 503-518.

Sabnis, T.S. 1919. The physiological anatomy of the plants of the Indian desert. J. Indian Bot. Sci. I : 42 and 65-83 I; 2:97-109 I; 4:97-113 I : 7-7, 33-43, 65-84, 183-205, 237-246 277-95. 2:1-19; 1920. 2: 61-79, 93-115, 157-73.

Sabnis and Bedi. 1971. Ceropegia odorata A little known plant of Western India.

Kew Bull. 25 : 57-60.

Safwat, F.M. 1962. The floral morphology of Secamone of the evolution of the pollinating apparatus in Asclepiadaceae

Ann. Mo. Bot. Gdn. 49 : 95-129.

Santapau, H. 1948. Miscellaneous notes 43. The genus Ceropegia further comments.

J. Bom. Nat. Hist. Soc. 47 :775-777.

Santapau, H. and Irani, N.A. 1958. The genus Ceropegia in Bombay.
Bull. Bot. Soc. Bengal 12 : 6-14.

Santapau, H. and Irani, N.A. 1960. The Asclepiadaceae and periplocaceae of Bombay

Saucer, H. et. al. 1965. Struktur der prävogenine (structure and drevogenine)

Helv. Acta. 48 : 857-878.

Saucer, H.H. and Weiss, E. 1966. The glycoside of the seeds of Dregea volubilis (L) Benthex Hook. Z. Isolation of more drevogenins.

Helv. Chin. Acta 40 : 1625-1632.

Saunders, E.R. 1939. Floral Morphology Vol. II Cambridge.

Sayeedud. and Saxena, 1940. On the anatomy of some of the Asclepiadaceae

Proc. Nat. Acad. Sci. Wash. 10:120-132.

Schill, R and Jakel, U. 1978. Beitrag Zur Kenntnis der Asclepiadaceen. Pollinatien tropische und subtropische. Pflanzenwelt : 22.

Schofield, E.K. 1968. Petiole anatomy of the Guttiferae and related families

Mem. N. Y Bot. Gard. 18 : 1-55.

Schnarf, K. 1931. Vergleichende Embryologie der Angiospermen Berlin.

Scott, D.N. and Sargent, E. 1893. On the pitchers of Dischidia rafflesiana Wall

Ann. Bot. Lond. 26 : 244-269.

Sedgwick, L.J. 1921. New Bombay species.

Journ. Indian Bot. 2 : 124-125.

Sharma P.C., Chandra V., Kapoor, S.L. and Kapoor, L.D. 1970.

Epidermal and venation studies in
Apocynaceae-II

Bull. Bot. Surv. India 14 : 76-82.

Sharma, H.P. and Misra, M.G. 1975. Tabulated phytochemical report.

Phytochem 14 : 1467-1468.

Singh, B. 1943. The origin and distribution of inter and intraxylary phloem in Leptadenia

Proc. Indian Acad. Sci. 18 : 14-19.

Singh, V. et/ al 1974. Trichomes in Salvia (Labiatae) and their taxonomic significance

Bull. Bot. Surv. India 16 : 27-37.

Singh, V. and Jain, D.K. 1975. Foliar Epidermal studies in Rosaceae-III: Prunus.

Ibid 17 : 137-146.

Small, J. 1913. The identification value of hairs

Pharm. J. 36 : 587-591

Sprengel, C.K. 1973. Das Entdeckte Geheimniss der Natur im Bau und inder Befruchtung der Blumen

Stace, C.A. 1961. Cuticular characters as an aid to the taxonomy of the Southwest African species of Combretum Mitt. Bot. Statssaml., Miinchen 4 :1-94.

Stace, C.A. 1965a. The significance of the leaf epidermis in the taxonomy of the Combretaceae

I. A general review of tribal, generic and specific characters.

J. Linn. Soc. London (Bot) 5 : 229-252.

Stace, C.A. 1965b. Cuticular studies as an aid to plant taxonomy
Bull. Br. Mus. Nat. Hist. (Bot) 4 :1-78

Stace, C.A. 1969a. The significans of the leaf epidermis in the taxonomy of Combretaceae.

II. The genus Combretum sub-genus Combretum in Africa.

J. Linn. Soc. London (Bot.) 62 : 131-168

Stace, C.A. 1969b. The significance of the leaf epidermis in the taxonomy of the Combretaceae.

III. The genus Combretum in America
Brittonia 21 :: 130-143.

Stace, C.A. 1973. The significance of the leaf epidermis in the taxonomy of the Combretaceae.

IV. The genus Combretum in Asia
Bot. J. Linn. Soc. 66 : 97-715.

Starr, A.M. 1912. Comparative anatomy of dune plants.

Bot. Gaz. 54 : 265-305.

Supate A.R., Gaikwad D.K., Yadav S.R. and Chavan P.D. 1990.

Crassulacean Acid Metabolism features
 occurred in leafy forms of Ceropegia.

Photosynthetica 24 : 270-272.

Szareek, S.R. 1979. The occurrence of crassulacean Acid Metabolism : A supplementary list during 1976-1979.

Photosynthetica 11 : 330-342.

Szarek, S.R. and Ting, I.P. 1977. The occurrence of Crassulacean Acid Metabolism among plants.

Photosynthetica 11 330-342.

Trochain, J. 1932. Note sur l'anatomie de la feuille dune Asclepiadaeae : le Leptadenia lancifolia Decne.

Bull. Soc. Bot. Fr. 79 : 28-32.

Tschesche, et. al. 1958, 1959. Sited in Pharmacognostic study
of Asclepias curassavica Linn. by Mitra,
R., Mehrotra S., Mehrotra B.N., and
Kapoor L.D.

Bull. Bot. Surv. India 16 : 82-88.

Van, L. 1937. Itallia Pharm. Weekblad 24 : 5-9

Venkatreddi, B. 1968. Ceropegia lawii concepts
J. Wiley & Sons Inc., New York.

Vajravelu, E. 1987. A note on the Brachystelma glabrum Hook.
(Asclepiadaceae) from South India.
J. Bomb. Nat. Hist. Soc. 84 : 262-263.

Wadhwa and Ansari M.Y. 1968. A new species of Ceropegia Linn.
(Asclepiadaceae) from Western Ghats,
Maharashtra.

Bull. Bot. Surv. India 10 : 95-97.

Walters, S.M. 1953. Eleocharis mamillata. Linbol. Fil. and allied
species.

Ber. Schweiz Bot. Ges. 62 : 271-286.

Wilson K.J., Peterson B.H. and Biesoer D.D. 1984.

Immunocytochemical identification of
laticifers.

Protoplasma 122 : 86-90.

Woodson, R.E. 1954. The North American species of Asclepias
L. Ann. Mus. Bot. Gard. 41 : 1-211.

Wyatt, R. 1976. Pollination and Fruit-set in Asclepias, a
reappraisal.
Amer. J. Bot. 63 : 845-851.

Waytt, R. 1980. The impact of nectar robbing ants on the
pollination system of Asclepias
curassavica Bull. Torrey Bot. Club.
107 : 24-28.

Yadav, S.R.; Salunkhe, C.B.; and, G.B. Dixit 1989 . Two
new records of Asclepiadaceæ from
Maharashtra.

J. Bom. Nat. Hist. Soc. 86 : 480-482.

Zemke, E. 1915. Anatomische Untersuchungen on Pflanzen der
Nambiwuste. (Deutsch-Sudwestafrika).
Flora Jena 33 : 665-416

Zornig, H. and Buch, O. 1926. Beitrag zur Anatomie des Blattes
pharmazeutisch gebrauchlicher Labiaten-
Drogen.

Arch. Pharm. Berl. 264 : 301-321.