BIBLIOGRAPHY

Abeles, F.B.; Dunn, L.J.; Morgans, P., Callahan, A.; Dinterman, R.E. and Schmidt, J. (1988). Induction of 33-KD and 60 KD peroxidase during ethylene enhanced senescence of cucumber cotyledons. Plant Physiol., 87 : 609-615.

Aberg, B. (1958). Ascorbic acid. In : Encyclopedia of plant physiology (W. Ruhland ed.) Springer. Berlin, 6: 479.

Abraham, P.G.; Dave, I.G.; Pandya, R.B. and Saxena, O.P. (1968). A study of the rate of respiration and Ascorbic acid metabolism during flowering and senescence of wheat. Proc. 37th Session Nat. Aca. Sci. pp.48.

*Adams, P.A. (1969). Studies of Gibberellic acid. Induced growth in Avena stem segments. Ph.D. Dissertation University of Michigen, Ann. Arbor.

Adaams, C.A.; Broman, T.H.; Norby, S.W.; Rinne, S.W. and Rinne, R.W. (1981). Occurrence of multiple forms of α-amylase and absence of starch phosphorylase in soybean. Annals of Botany, 48(6) : 895-903.

Ali AL-Ani; Francoise Bruzao; Philippe Raymond; Verorique Sainteges; Jean Maro Leblanc and Alain Pradet (1985). Germination, respiration and Adenylate energy charge of seeds at various oxygen partial pressures. Plant Physiol, 79 : 885-890.

Amer, M.A.; Mbdel-Rasoul, M.; Kandil; S.A. and Raafat, A. (1988). The effect of seed soaking with Ancymidol or CCC on the accumulation of dry matter, aminoacids and nucleic acids at early stage of maize growth under different water regims. Proceeding of International Congress of Plant Physiology, New Delhi. pp.123.

Anderson, J.D. (1979). Purine nucleotide metabolism of germination soybean embryonic axes. Plant Physiology, 63(1) : 100-104.

*Anderson, M.B. and Cherry, J.H. (1969). Proc. Natl. Acad. Sci., 62 : 202-209.

Arrigoni, O.; DeGara, L.; Tommasi, F. and Liso, R. (1992). Changes in the ascorbate system during seed development of *Vicia faba* L. Plant Physiol., 99 : 235-238.

Arnon, D.I.; Whatley, F.R. and Allen, M.B. (1957). Photosynthesis by isolated chloroplasts II. Photosynthetic phosphorylation, the conversion of light into phosphate bound energy. J.Am. Chem. Soc., 76 : 6324-6329.

Arora, S.K., Banerjee, M.K., Pratap, P.S. and Srivastava, V.K. (1990). Hormonal regulation of plant growth and development Vol.V. Ed.S.S. Purohit, pp.125-148. Agro Botanical Publisher (India) Bikaner.

Asada, K. (1992). Production and scavenging of active oxygen in chloroplasts. In : Scandalics, J.C. (ed.). Molecular Biology of free radical scavenging systems p.173. Cold Spring Harbar Laboratory Press, New York.

Aslam, M.A., Oak, S. and Huffaker, R.C. (1976). Effect of light and glucose on the induction of nitrate reductase and on the distribution of nitrate in etiolated barley leaves. Plant Physiol., 58 : 588-591.

*Asthana, J.S. and Srivastava, H.S. (1978). Indian J. Plant Physiol., 21 : 150.

Austin, R.B.; Ford, M.A. and Blackwell, R.D. (1978). Relationships between nitrate reductase activity. Plant weight and nitrogen content in seedlings of *Triticum, Aegilops* and *Triticale*. Ann. Bot., 42 : 429-438.

Baba,I., Inada, K. and Tajima, K. (1965). Mineral nutrition and the occurrence of physiological disease in the mineral nutrition of the rice plant. Proceeding of a symposium at the international rice research institute. Pub. John. No. Kins Press Baltimore, Meryland.

Bae, Sookyung, Hahn shik Kwak and Chungnoojoo (1986). Biochemical studies on ginseng saponin XXV. The effect of gingseng saponin fraction on the biosynthesis of Ascorbic acid during germination of soybean (*Glycine max*). Korean Biochem J., 19(1) : 1-5.

Balma, D. (1988). Study of germination capacity in soybean seeds produced in Tropical Africa. Oleagineux, 43(6) : 255-259.

Barbaro, A.DE. and Kuchmierz. T. (1980). The effect of temperature on the effectiveness of selected pesticides during the germination and growth of soybean (*Glycine max* (L.)Merrill) Part II.Biuletyn Instituta Hodowli Aklmatyzacji Roslin No.140, 51-65.

Barbaro, A.DE. and Szalonck, K. (1980). The effect of temperature on the effectiveness of selected pesticides during the germination and growth of soybeans (*Glycine max* (L.) Merill). Part I. Biuletyn Instituta Hodowli Aklmatyzacji, Roslin No.140, 39-50.

Barclay, L.R.C. (1988). The co-operative antioxidant role of glutathione with a lipid soluble and a water soluble antioxidant during peroxidation of liposomes initiated in the aqueous phase and in the lipid phase. J. Biol. Chem., 263 : 16138-16142.

Barker, G.R.; Bray, C.M. and Water,T.J. (1974). Development of Ribonuclease and acid phosphatase during germination of *Pisum ariense*. Biochem. J., 142 : 211-219.

Bate's L.S.; Waldren, R.P. and Teare, I.D. (1973). Rapid determination of proline for water stress studies. Plant Soil, 39 : 205-207.

Beck, J.L.; McConachie, L.A.; Summors, A.C.; Arnold, W.N.; DeJersey, J. and Zermer, B.C. (1986). Properties of a purple phosphatase from red kidney bean : a zinc-Iron-metalloenzyme. Biochem-Biophys. Acta., 869 : 61-68.

Beevers, H. (1969). Glyoxysomes of castor bean endosperm and their relation to gluconeogenesis. Ann. N.X. Acad. Sci., 168 : 313-324.

Bendich, A.; Machlin, L.J.; Scandura, O.; Burton, G.W and Wyner, D.D.M. (1986). In 'Advances in Free Radical Biology and Medicine" (W.A. Pryor ed.) Vol.2, pp.419-444, Pergamon, New York.

Bensal, R.C.; Tiwari, R.K.; Singh, U.M. and Kiran, Ravi (1981). Biochemical changes during germination of muskmelon (*Cucumis* melo) seeds. Plant Biochem. J., 8(1) : 28-33.

Bernstein, L. (1974). Crop growth and salinity in "Drainagr for Agriculture". Agronomy, 17 : 39-54.

Bergmann, L. and Ulbrich, J. (1981). In biology of inorganic nitrogen and sulfur (Bothe. H. and Trebst A. eds.) Springer, Berlin.

Bewley, J.D. and Black,M.(1978)."Physiology and Biochemistry of Seeds in relation to germination Vol.I, Development, germination and growth". Springer, Verlag, Berlin.

Bhatnagar, P.S. (1994). Soybean seed and its production techniques. National research centre for soybean, ICAR, publication. pp. 1-10, Khandwa Road, Indore.

Bhadre, S.K. (1983). Physiological studies in cotton plant. Ph.D. thesis submitted to the Shivaji University, Kolhapur (India).

Bhandari, M.C. and Sen, D.N. (1975). Effect of growth retardants on seedling growth excised cotyledons, chlorophylls, proteins and sugar content in *Citrulus species*. Biochem. Physiol. Pflanz (BPP) 167 : 135-140.

Bhargava, R. and Sachar, R.C. (1987). Induction of acid phosphatase in cotton seedlings. Enzyme purification, sub unit structure and kinetic properties. Phytochemistry, 26 : 1293-1297.

*Bick, M.D.; Liebke, H.; Cherry, J.H. and Strehler, B.L. (1970). Biochem. Biophys. Acta., 204 : 175-182.

Biswas, T.K. and Candiff, C. (1991). Multiple forms of acid phosphatase in germinating seeds of *Vigna sinensis*. Phytochemistry, 30 : 2119-2125.

Bisaria, A.K. and Paliwal, N.K. (1982). Effect of ethephon on seed germination, seedling growth and sugars in triticale. Acta. Bot. Indica., 9(1) : 148-150.

Bond, H.M., Bowles, D.J. (1983). Characterization of soybean endopeptidase activity using exogenous and endogenous substrate. Plant Physiol., 72 : 345-350.

Bourris, S. (1978). Hormonal regulation of phosphatase activity in after ripened and dormant *Agrostemma cithago* L. embryos. Biochem. Physiol (Pflanza) 172(5) : 453-474.

Boutin, J.P.; Provot, M. and Roux, L. (1981). Effect of cycloheximide and renewal of phosphorus supply on surface acid phosphatase activity in phosphorus deficient tomato roots. Physiol Plant, 51 : 353-360.

Bray, C.M.; Water, T.J. and Barker, C.R. (1979). Development of ribonuclear and acid phosphatase during germination *Pisum orverner*. Biochem. J., 142(2) : 211-219.

*Burlison, W.L.; J. Doran, C.A. and Hackleman, J.C. (1940). Agri. Expt. Sta. Bull., 462 : 123-127.

Calero, E.; West, S.H. and Hinson, K. (1981). Water absorption of soybean seeds and associated causal factors. Crop. Sci., 21 : 926-923.

Cartter, J.L. and Hopper, T.H. (1942). Influence of variety, environment and fertility level on the chemical composition of soybean seed.USDA.Tech.Bull.,7 : 87-89.

Cartter, J.L. (1940). Effect of environment on composition of soybean seeds. Soil. Sci. Soc. Am. Proc., 5 : 125-131.

*Carnegie, P.R. (1963). Biochem J., 89 : 459.

*Catsimpoolas, N.; Campbell, T.G. and Mayer, E.W. (1968). Plant Physiol., 43 : 799-805.

Chabot, J.F. and Leopold, A.C. (1985). Ultra structural aspects of chiling injury in the soybean (*Glycine max*). Seed radicle. Am. J. Bot., 72(7) : 1120-1126.

Chakravorti, M. and Burma, D.P. (1959). Enzymes of the pentose phosphatase pathway in the mungbean seedlings. Biochem. J., 73 : 48-53.

Chanprasert, W.; Jintakanon, S. and Wongmaneeroj, A. (1992). The effect of pregermination and ageing on germinability under different salt concentrations of soybean (*Glycine max* (L.) Merrill). Seeds. Karsert Journal, Natural Science, 26(1) : 90-94.

Chao, S.E. (1970). Biochemical genetics and developmental studies of starch degradation enzymes in maize. Ph.D. dissertation Michingham state Univ.

Chavan, S.R. (1987). Physiological studies in nitrogen metabolism of groundnut (*Arachis hypogea* L.) M.Phil. Thesis submitted to the Shivaji University, Kolhapur (India).

Cheng, C.L.; Acedo, G.N.; Christinsinm and Cronkling, M.A. (1992). Sucrose mimics the light induction of Arabidopsis nitrate reductase genetranscription. Proc. Natl. Acad. Sci., USA, 89 : 1861-1864.

*Chikasne, M. (1953). Bull. Fac. Agric. Mic. Univ., 7, 188, 1953.

Chinoy,J.J. and co-workers (1969). A.PL480 project report submitted to Gujarath University, Ahmedabad, India.

Chinoy, J.J. and co-workers (1984). The role of Ascorbic acid in growth, differentation and metabolism in plants (Eds. N.J. Chinoy). Martinus Nijhoff/DR/W. Junk Publishers.

Chinoy, J.J. and Saxena, O.P. (1972). Inductive effect of ascorbic acid on RNA, amylase, protease and RNAase. Proc. Int. Seed Test ASS., 37 : 903-910.

Chinoy, J.J.,Saxena, O.P., Abraham,P.G., Pandya, R.B., Dave, I.C., Gurumurthi, K. and Sastri, K. (1969). Physiology of seed germination in relation to the early production of AA and Ascorbic acid oxidase in wheat (*Tricticum*) and peanut (*Arachis*). Final Res. Prog. Report of Pl.480.480, Res. Project.

Ching, J.M. (1972). Metabolism of germinating seeds. In 'Seed Biology', Vol.11 (T.T. Kozlowski ed.), Academic Press, New York.

Chrispeels, M.J. and Varner, J.E. (1967). Hormonal control of enzyme synthesis on the mode of gibberellic acid and abscission in aleurone Layers of barley. Plant Physiol, 42 : 1008-1016.

Christina,W.N. and Leopold,A.C. (1983). Dynamics of inhibition by soybean embryos. Plant Physiol., 72 : 190-193.

Citterios, S.; Sergio Sgorbati; Stefania Scippana and Elio Sparvoli (1994). Ascorbic acid effect on the onset of cell proliferation in pea root. Physiologia Plantaram, 92 : 601-607.

Colson (1979). Ultra structural localization of acid phosphatase activity and phosphatase activity and phosphatase in radical of *Allium cepa* in ungerminated seeds and at the beginning of germination effects of short treatments at low temperature (-196°C). Ann. Sci. Nat. Biol. Veg., 1(1) : 21-38.

Covell, R.H.; Ellis,R.H.; Roberts,E.H. and Summerfield, R.J. (1986). The influence of temperature on seed germination rate in grain legumes. Journal of Experimental Botany., 37, No.178, pp.705-715.

Cruz, R. and Avelino, L.D. (1986). Soybean (*Glycine max*), endogenous and galactosidase and invertase during germination process. AQ Biol. Technol., 29(3) : 435-443.

Dashek, W.V. and Erickson, S.S. (1981). Isolation, assay, biosynthesis, metabolism, uptake and translocation and function of proline in plant cells and tissues. Bot. Rev., 47 : 349-385.

Davis (1964).Disc electrophoresis II. Method and application to human serum protein. Ann. Ny. Acad. Sci., 12 : 404-427.

Daubert, B.F. (1950). In : "Soybean and soybean products". Vol.1. pp.157. Interscience, New York.

Daussant, J., Neucere, N.J. and Conkerton, E.J. (1969). Plant Physiol., 44 : 480-489.

Dawson, M.J. (1965). Effect of seed soaking on the growth and developments of crop plants I. Finger Millet (*Eleusine coracana*, Goer). Indian J. Plant Physiol., 8 : 52-56.

Deng, M.; Moureaux, T.; Cherel, I.; Boutin,J.P. and Caboche, M. (1991). Effects of nitrogen metabolites on the regulation and circadian expression of tobacco nitrate reductase. Plant Physiol. Biochem., 29 : 239-247.

*Denny, F.E. (1924,a). Bot. Gaz., (Chicago) 7 : 322.

*Denny, F.E. (1924b). J. Agric Res., 27 : 757.

Dandsay, J.P.S. and Sachar, R.C. (1978). Hormonal control of peroxidase activity in germinating mungbean cotyledons. Phytochemistry, 17(6) : 1017-1019.

*deKok, L.J., deKan, P.J.L., Tanezos, O.G. and Kuiper, P.J.C. (1981). Physiol. Plant, 53 : 435.

Deleo, P. and Sacher, J.A.C. (1970). Senescence : Association of synthesis of Acid phosphatase with Banana ripening. Plant Physiol., 46(2) : 208-211.

*Delouche, J.C. (1953). Proc. Assoc. off Seed Annals., 43 : 117-126.

Delouche, J.C.; Mathes, R.J.; Dougherty, G.M. and Boyd, A.H. (1973). Storage of seed in subtropical and tropical regions. Seed. Sci. Technol., 663-692.

Delouche, J.C. (1974). Maintaining soybean seed quality soybean production, marketing and use. Tenn. Val. Auth. Bull. Y., 69.

De Oliveira, J.C.; Dasilva, D.M.; De Amorim, H.V. (1976). Enzymic activity of polyphenol oxidase and catalase in *Coffea Arabica* L. seeds and seedlings. Cientifica 4(1) : 68-70.

Dev Prakash, V. (1970). Nitrate reductase activity in cotton seedlings treated with gibberellic acid and 2-chloroethyl trimethyl ammonium chloride. Indian J. Plant Physiol., 13 : 67-71.

Dhopte, A.M. and Phadnawis, B.N. (1989). Estimation of ascorbic acid (AA) with dye method. Biochemical analysis of plant scientist (eds. A.M. Dhopte and Livera M.) Forum for plant physiologist, Akola (India), pp.243.

Dies, E.J. (1992). Soybean Gold from the soil. New York, Macmillan

Dizengremel, P. and Tuquet, C. (1984). Changes in respiration and mitochondrial activities in cotyledons from imbibition to senescence. Physiol. Veg., 22 : 687-700.

Dornbos, D.L. and Jr. Mullen, R.E. (1991). Influence of stress during soybean seed fill on seed weight, germination and seedling growth rate. Canadian Journal of Plant Science. 71(2) : 373-383.

*Drapron,R.; N. Guyan X.U. an Anh; Launay, B. and Guibot, A. (1969). Cereal Chem., 46 : 647-655.

Dracup M.N.H.; Barrett-Lennard, E.G.; Greenway, H.; Robson, A.D. (1984). Effect of phosphorus deficiency on phosphatase activity of cell walls from roots of subterranean clover. J. Exp. Bot., 35 : 466-480.

Dua, S.; Sareen, K.N. and Amma, M.K.P. (1983). Hormone induced changes in acid phosphatase activity during imbibition of 96 hr. germinated castor seed. Indian J. Biochem Biophys., 20(2) : 118-120.

Duff, S.M.G.;Lefebvre,D.D. and Plaxton,W.C.(1989). Purification and characterization of phosphoenol pyruvate phosphatase from *Brassica nigra* suspension cells. Plant Physiol., 90 : 734-741.

Duff, S.M.G.; Gautam, S. and William, C. Plaxton (1994). The role of acid phosphatase in plant phosphorus metabolism. Physiol. Plant, 90 : 791-800.

Duke, S.H.; Kakefuda, G.; Hensen, C.A.; Loeffler, N.L. and Vanhulle, N.M.C. (1986). Role of the testa epidermis leakage of intracellular substances from imbibing soybean seeds and its implications for seed survival. Physiologia Plantarum, 68 : 625-631.

*Dzikowski, B. (1936). Pamiet. Panst W. Inst. Nauk. Gospod Wiejske, 16 No.253, pp.69-100.

Eastwell, K.C. and Spencer, M.S. (1982). Effect of ethylene on the gibberellic acid enhanced synthesis and release of amylase by isolated barley aleurone layers. Plant Physiol., 69 : 557-562.

Eastwell, K.C. and Spencer, M.S. (1982b). Ethylene effects on amylase activity from isolated barley aleurone layers; possible modification by proteolytic enzymes. Plant Physiol., 70 : 849-852.

Egley, G.H.; Paul,R.N.; Vaughan,K.C. and Duke,S.O. (1983). Role of peroxidase in the development of water

impermeable seed coats in *Sida spinosa* (L.). Planta, 157 : 224-232.

ELLman, G.L. (1959). Tissue sulfadryl groups. Archives of Biochemistry and Biophysics., 82 : 70-77.

*Fahey, R.C.; Brody, S. and Mikolajazyk, S.D. (1973). J. Bacteriol., 121 : 144.

*Fahey, R.C.; Distefano, D.L.; Meier,G.P. and Bryan, R.N. (1980). Plant Physiol., 65 : 1062.

*Fahey, R.C.; Newton, G.L.; Dorian, R. and Kosower, E.M. (1980). Analyt Biochem., 107 : 1.

Fang, S.C.; Bourke, J.B.; Stevens, V.L. and Butts, J.S. (1960). Influence of Gibberellic acid on metabolism of indole acetic acid acetate, and glucose in roots of higher plants. Plant Physiol., 35 : 251-255.

Fehr, W.R. (1978). Breeding in soybean physiology, agronomy and utilization ed. A.G. Norman, New York, Academic Press.

Feng, D.S. and Egli, D.B. (1994). Germination of fresh, immature soybean seed : Effect of the pregerminational light environment. Seed Science and Technology, 22 (1) : 33-42.

Ferguson, J.M.; Tekrony, D.M. and Egli, D.B. (1990). Changes during early soybean seed and axes deterioration : I seed quality mitochondrial respiration. Crop Sci., 30 (1) : 175-179.

Ferrari, T.E.; Yoder, A.C. and Filner,P. (1973). Anaerobic nitrite production by plant cells and tissues. Evidence of two nitrite pools. Plant Physiol., 51 : 423-431.

*Fiske, C.H. and Subba Raw, Y. (1925). J. Biol. Chem., 66 : 575.

Flin, A.M. and Smith, D.L. (1987). The localization of enzymes in cotyledons of *Pisum arvense* L.during germination. Planta, 75 : 10-22.

Fox, L.P.; Puryes, W.K. and H.I. Nakade (1965). The role of horse radish peroxidase in indole-3-acetic acid oxidation. Biochemistry, 4 : 2754-2763.

Foyer, C.H. and Halliwell, B. (1976). The presence of glutathione and glutathione reductase in chloroplasts : a proposed role in ascorbic acid metabolism. Planta, 133 : 21-25.

Gaff, D.F. (1966). The sulphydryl-Disulphide hypothesis in relation to desiccation injury of cabbage leaves. Aust. J. Biol. Sci., 19 : 291-299.

Gardon, A.G. (1973). The rate of germination in seed ecology, Ed. W. Heydecker, pp.284-311. Publ. Butterwortho, London.

*Gasper, T., Verbeek, R. and Khan, A.A. (1971). Planta, 83 (4), 387.

Gasper, T.; Penel, C.; Thorpe, T. and Greepin, H. (1982). 'Peroxidases' Publ. Universite P. Genue, Switzerland.

Gelmond, H. (1978). Problems in crop seed germination in crop physiology, pp.1-78, U.S. Gupta (Ed.) Oxford Loid. Publishing Co., New Delhi.

Genkel, P.A. (1961). Drought resistance in plants, methods of recognition and intensification. In Proc. Madrid Symp. Vol.XVI UNESCO, Paris, pp.167-174.

Genkel, P.A. and KoLotova, S.S. (1934). O. Predposevnoi zakolke restenii Kzasukhe Vustouyakh Vegetataionnogo Opyta. (Izv. Parmskogo biol. Nautonoissled). Bull. Perm. Biol. Sci. Inst., 9 : 1-3.

George, A.A.; De-Lumen, B.O. (1991). A novel methyonine rich protein in soybean seed : Identification, aminoacid composition and N-terminal sequence. J. Agric Food. Chem., Washington, 39(1) : 224-227.

Gibson, D.M. and Ullah, A.H. (1988). Purification and characterization of phytase from cotyledons of

germinating soybean seeds. Arch. Biochem. Biophys., 260 : 503-513.

Gilman,D.F.; Fehr, W.R. and Burris, J.S. (1973). Temperature effects on hypocotyl elongation of soybean. Crop. Sci., 13 : 246-249.

Goldstein, A.H.; Baertlein, D.A. and Danm, A. (1989). Phosphate starvation stress as an experimental system for molecular analysis. Plant Mol. Biol. Rep., 7 : 7-16.

Gopalchari, N.C. (1963). Changes in the activities of certain oxidizing enzymes during germination and seedling development of *Phoseolus mungo* and *Sorghum vulgare*. Indian J. Exp. Biol., 1(2) : 98-100.

Goyal, M.R.; Drew, G.L.; Nelson and Logan, T.J. (1980). Soybean seedling emergence force. Transactions of the Asae, 23(4) : 836-839.

Goswami, A.K.; Jain, M.K. and Pauls, B. (1977). B-amylase in seed germination. Soil Plant, 19(6) : 469-471.

Grabe, D.F. and Metzer, R.B. (1969). Temperature induced inhibition of soybean hypocatyl elungation and seedling emergence. Crop Sci., 9 : 331-333.

*Grable, A.R. and Danielson, R.E. (1965). Soil. Sci. Soc., Am. Proc., 29 : 12-18.

Granstedt, R.C. and Huffaker,R.C. (1982). Identification of leaf vacuole as a major nitrate storage pool. Plant Physiol., 70 : 85-88.

*Grinberg, A.A. (1971). Physiocochemical basis of the protective function of catalase. Latv. PSR ZI Akad. Veslis., 2 : 60-62.

Guerero, M.G.; Vega, J.M. and Losada, M. (1981). The assimilation of nitrate reducing system and its regulation. Ann. Rev. Plant. Physiol., 32 : 169-204.

Guang-hua, Z.; Zhen-hui, G. and Ben-mei, X. (1981). A physiological study on the chilling injury in the germinating soybean seeds. Scientia Agricultura Sinica 2 : 65-72.

Gunning, B.E.S.; Steer, M.W. and Cochrane, M.P. (1968). "Occurrence molecular structure and induced formation of the stroma centre" in plastids. J. Cell. Sci., 3 : 445-456.

Hadas, A. (1976). Water uptake and germination of leguminous seeds under changing external water potential in osmotic solutions. J. Expl. Botany, 27 : 480-489.

Hadley, P., Roberts, E.H.; Summerfield, R.J. and Minchin, F.R. (1984). Effects of temperature and photoperiod on flowering in soybean (*Glycine max* (L.) Merill) : A Quantitative Model, Ibid, 53 : 669-681.

Hailstones, M.D.; Smith, M.T. (1991). Soybean seed invigoration by ferrous sulphate : change in lipid peroxidation, conductivity. Tetrazolium reduction DNA and protein synthesis. Journal of Plant Physiology, 137(3) : 307-311.

Halvey, A.N. (1964). Effect of gibberellic and growth retarding chemicals on respiration and catalase activity in various organs of cucumber seedlings. J. Expt. Bot., 15 : 546-555.

Hang, J.T.; Nam, M.H.; Lee, S.K. and Park, W. (1983). Changes in peroxidase activity and its isozymes of soybean, red bean and mung bean during germination. Hanguk Nonghwa Hakhoechi, 26(3) : 151-156.

Harper, J.E. and Hageman, R.H. (1972). Canopy and seasonal profiles of nitrate reductase in soybean (*Glycine max* (L.) Merr.). Plant Physiol., 49 : 146-154.

Harper, J.E.; Nicholas, J.S. and Hageman, R.H. (1972). Seasonal and conopy variation in nitrate reductase

activity of soybean (*Glycine max* (L.) Merrill.) varieties. Crop. Sci., 12 : 382-386.

HanishTenGate, C.H. and H. Bretcler (1982). Effect of plant growth regulators on nitrate utilization by roots of nitrogen depleted dwarf bean. J. Expt. Bot., 132 : 36-46.

Hasson, P.E. and Poljakoff Mayber,A. (1981). Germination of pea seeds exposed to saline stress. Israel J. Bot., 29 : 98-104.

*Hatfield, J.L. and Egliy, D.B. (1974). Crop. Sci., 14 : 423-429.

Howell and Collins (1957). In : Book World Oil Seeds, pp.12-13.

Herbert, D. (1955). Catalase from bactreia, In "Methods in enzymology", Vol.II, Eds,Colowick, S.P. and Kaplan, N.O. Academic Press Inc. Publishers, New York, pp.784-786.

Hendricks, J.D. and Taylorson, R.B. (1974). Promotion of seed germination by nitrate, nitrite, hydroxylamine and ammonium salts. Plant Physiol., 54 : 304-309.

Hidalgo, A.; Garcia-Herdugo G.; Gonzalez-Reyes J.A.; Morre, D.J. and Navas, P. (1991). Ascorbate free radical stimulates onion root growth by increasing cell elongation. Bot. Gaz., 152 : 252-288.

Hideaki, K. (1977). Acid phosphatase isozymes in the endosperm of rice seeds. Akita Daigaka Kyoikygakuba Kenkya Kiyo Shizen Kagaka, 27 : 72-77 (Japan).

*Hildebrand, A.A. (1959). Can. J. Bot., 37 : 927-957.

Hildebrand, P.F.; Hymowitz, T.C. (1981). Role of β-amylase in starch metabolism during soybean seed development and germination. Physiologia Plantarum, 53(4) : 429-434.

Hill, H.J.; S.H. West and K. Hinson (1986). Soybean seed size influences expression of the impermeable seedcoat trait. Crop. Sci., 26 : 634-637.

Hoagland Robert, E. (1987). Glyphosphate effects on proteolytic enzyme activity in soybean axes. Bot. Gaz., 148(1) : 7-11.

Howell, R.W. (1963). In "The Soybean" (A.G. Norman, ed, pp.75-124. Academic Press, New York.

Hopkins, W.G. and Hillman, W.S. (1965). Phytochrome changes in tissues of dark grown seedlings representing various photoperiodic classes. Am. J. Botany, 52 : 427-432.

HODTH, Keates, RAB, Varner JE (1977). Effect of ethylene on the release of GA_3 induced α-amylase from barley aleurone layers. Plant Physiol. (Suppl.), 59 : 56.

HODTH, Varner, JE (1976). Response of barley aleurone layers to abscisic acid. Plant Physiol., 57 : 175-178.

*Huang,A.H.C. and Moreau,R.A. (1978). Planta, 141 : 111-116.

Hudson, S.J. (1985). Phytic acid mobilization in germinating soybean seeds. M.Sc. Thesis, University of Illinois, Urbana.

Hutton, D. and Stumpf,P.K. (1969). Characterization of the ß-oxidation systems from maturing and germinating castor bean seeds. Plant Physiol., 44 : 508-516.

Hunter, J.R. and Erickson, A.E. (1952). Agron. J., 44 : 107-109.

Hymavaty, R.K. (1983). Germination studies in finger millet (*Eleusine coracana* Gaertn). M.Phil. Thesis submitted to Shivaji University, Kolhapur.

Imeson, H.C.; Rood, S.B.; Weselake, R.J.; Zanewich, K.P.; Bullock, B.A.; Stobbs, K.E. and Koesis, M.G. (1993). Hormonal control of lipase activity in oil seed rate germinants. Physiol Plant, 89 : 476-482.

Innocenti, A.M.; Bitonti, M.B.; Arrigoni, O. and Liso, R. (1990). The size of quiescent centre in roots of *Allium cepa* L. grown with ascorbic acid. New Phytol., 114 : 507-509.

*Inouye,C. (1953). Proc. Crop. Sci. Soc. Jpn., 21 : 276-277.

Ioana, M. (1969). Physiological changes during the germination of some legume seeds. *Luerari ofmt*. Inst. Agron (N. Balecescusera), 12 : 299-306.

Ishinaga, M.; Yamauchi, T; Egusa, K. and Kuroda, K. (1992). Changes of dolichol fatty acyl esters during the germination and development of soybeans. Biochemistry and Cell Biology, 70(6) : 466-469.

Islam, M.T. (1978). Effects of sodium chloride on soybean seed germination. Bangladesh journal of Bot. 7(1) : 107-109.

Ivanova, T.M.; Devydove, M.A. and Rubin, B.A. (1967). On catalytic functions of peroxidase. Biochemiya, 32 : 607-611.

*Jackson, G.A.D.; Wood, R.B. and Prosser, M.V. (1961). Nature (London), 191 : 282-283.

Jacobsen, J.V.; Scandalics, J.G.; Varner, J.E. (1970). Plant Physiology, 45 : 367-371.

Janowski, R. and Iguacio (1977). Relation between and phosphatase content and germinating capacity in seeds of different species. Circ. Farm., 35(254) : 33-44.

Jaworski, E.G. (1971). Nitrate reductase assay in intact plant tissues. Biochem. Biophys. Res. Commun., 43 : 1274-1279.

Johnson, C.B.; Holloway, C.B.; Smith, H. and Grierson, D. (1973). Isozymes of acid phosphatase in germinating peas. Planta, 115(1) : 1-10.

Joshi, G.V. and Hegde, B.A. (1978). Ascorbic acid and stress

physiology. Proc. of symposium organised by Indian Soc. Plant Physiology, Ahmedabad, India. pp.21.

*Jones, E. and Hughes, R.E. (1983). Phytochemistry, 22 : 2493-2499.

*Jones, E. and Hughes, R.E. (1984). Phytochemistry, 23 : 2366-2367.

Jones, R.L. and Carbonell, J. (1984). Regulation of synthesis on Barley Aleurore α-amylase by gibberellic acid and calcium ions. Plant Physiol., 76 : 213-218.

Jones, R.S. and Mhcheli,C.A. (1989). Calcium ion involvement in growth inhibition of mechanically stressed soybean (*Glycine max*) seedlings. Physiologia Plantarum, 76(4) : 598-602.

Justice, O.L. and Louis N.B.(1978). Principles and practices of seed storage. USDA Agr. Hand Book, 506, 298 p.

Kaloyereas, S.A. (1958). A new method of determining drought resistance. Plant Physiol., 33 : 232-233.

Kamb oj, R.K. (1977). Some biochemical aspects of germination in soybean. Thesis abstract (1977) 3(½) 49 [En]. Dep. of Biochem. Naryana Agric. Univ. Hissar, 125004, Haryana, India.

Kamoj, R.K. and Nainawatee, H.S. (1978). Peroxidase isoenzyme changes in germinating soybean varieties. Seed Res., 6(2) : 140-144.

*Kamynina, L.M. (1965). Agrokhimiya, 10 : 123.

*Kanabus, J., Cherry, J.H.(1971). Proc. Natl. Acad. Sci., 68 : 873-876.

Kasai, T. and Larson, P.O. (1980). In progress in the chemistry of organic products (Herz. W. Griselach, H. and Kirby, G.W. Eds.) Vol.39, No.243, Springer, Wien.

Katsumi, M. and Fukuhara, M. (1969). The activity of α-amylase in the shoot and its relation ot gibberelin induced elongation. Physiol Plant, 22 : 68-75.

Kaul, K. and Farooq, S. (1992). Kinetin induced changes in growth and activity of some enzymes in morning glory hypocotyl segments. Indian J. of Plant Physiol., 37(4) : 214-216.

Kaur, M. (1987). Effects of some phenols on nitrogen metabolism in two peanut (*Arachis hypogaea* L.) varieties Ph.D. Thesis, PAU, Ludhiana.

Khan, A.A. (1978). Incorporation of bioactive chemicals into seeds to alleviate environmental stress. Acta. Hort., 83 : 225-234.

Khan, A.A.; Tao, K.L.; Knypl, J.S.; Borkowaska, B. and Powell, L.E. (1978). Osmotic conditioning of (vegetable) seeds : Physiological and bio-chemical changes. Acta. Hort., 83 : 267-278.

Kim, D.H.; Yum. C.A. and Kim, W.J. (1990). Kinetic study of hydration and volume changes of soybeans during soaking. Hanguk Nonghwahak Hoechi, 33(1) : 18-23.

*Kirsop, B.H. and Pollock, J.R.A. (1957). European brewery convention, p.84.

Kittoch, D.L. and Law, A.G. (1957). Relationship of seedling vigour to respiration and tetrazolium reduction by germinaing wheat seedling. Agron. J., 60 : 286-288.

Knypl, J.S. and Knypl, M.A. (1993). Increase chilling stress tolerance in soybean seeds treated with polyacrylic acid ethyl acrylate vinyl acetate, a superabsorbent. Acta Physiologiaeplantarum, 5(1), 3-9.

KoslaNund, R., Delouche, J.C. (1987) Factors affecting imbibitional injury in soybean seeds. News letter of the association of official seed Analysis, 61(2) : 13.

Kosower, E.M. (1976). In Glutathione : metabolism and function (Arias, J.M. and Jako by W.B. eds.) p.1, Raven Press, New York.

Kosower, N.S. and Kosower, E.M. (1974). In Glutathione (Flohe, L. Benohr, H. Ch., Sies. H. Walloc, H.D. and Wordel A. eds). p.276, George Thomas Stuttgart.

Kosower, N.S. and Kosower, E.M. (1976). Glutathione metabolism and function (Arias, I.M. and Jakoby, W.B. eds.), p.159. Raven Press, New York.

Kosower, N.S. and Kosower, E.W. (1978). The glutathione status of cells. Int. Rev. Cytol., 54 : 109-160.

Kremer, M.L. (1970). Peroxidatic activity of catalase, Biochem. Biophys. Acta., 198 : 199-209.

Krober,O.A. and Cartter,J.L. (1962). Quantitative interrelationships of protein and nonprotein constituents of soybeans. Crop. Sci., 2 : 171-175.

Kubota, Y. Shojis; Yamaraka, T. and Yamato, M. (1976). Carboxypeptidase from germinating soybeans. I. Purification and propertie of two carborypeptidase. Yakugaku Zasshi, 96 : 639-647.

Kuvaera, E.B. and Kretovich, V.L. (1978). Phytase of germinating pea seeds. Fiziol. Rast., 25(2) : 366-372.

Ku, H.S.; Yang, S.P. and Pratt, H.K. (1970). Ethylene production and peroxidase activity during tomato fruit ripening. Plant Cell Physiol., 11 : 241-246.

Kutwal, D.N. (1989). Influence of chloro choline choride (CCC) pretreatment on physiology of groundnut (*Arachis hypogaea* L.). M.Phil. Thesis submitted Shivaji University, Kolhapur, India.

Kwon, E.O. and Gun, J.O. (1987). Effect of presowing treatment and growth regulators on different growth and yield contributing parameters in soybean (*Glycine max* (L.) Merrill). Korean Journal of Crop Science, 32(4) : 392-402.

*Laughland and Laughland (1939). Advances in Agronomy American Society of Agronomy, Vol.I, Academic Press, I. Nc. Publ., New York,1 pp.107 [Edited by A.J. Norman].

LeBansky, B.R.; Meknight, T.D. and Lawrence, L.R. (1992). Purification and characterization of asecreted purple phosphatase from soybean suspension culture. Plant Physiol, 99 : 391-395.

Lefebvre, D.D.; Duff, S.M.G.; Fife,C.; Singh, J.I. and Plaxton, W.C. (1990). Response to phosphate deprivation in *Brassica nigra* suspension cells. Enhancement of intracellular, cell surface and secreted phosphatase activities compared to increase in P1 absorption rate. Plant Physiol., 93 : 504-511.

Leopold, A.C. (1980). Temperature effects on soybean imbibition and leakage. Plant Physiol., 65 : 1096-1098.

Lee, P.L.Y. (1974). Three hour single column sodium citrate, physiological analysis - Durrum Application Notes No.10, Dioney, Group., Synnyvale, CA.

Lee, C.K. and Karunanithy, R. (1990). Effects of germination on the chemical composition of Glycine and phaseolus Beans. J. Sci. Food Agric., 51 : 437-445.

Levitt, J. (1962). A sulphydryl-disulfide hypothesis of frost injury and fesistance in plants. J. Theoret, Biol., 3 : 355-391.

Levitt, J. (1980). Response of plants to environmental stress 2nd ed., Vol.2, Academic Press, New York.

Lillo, C. (1994). Light regulation of nitrate reductase in green leaves of higher plants. Physiol Plant, 90 : 616-620.

Liener, I.E. (1978). Nutritional value of food protein products in soybeans. Chemistry.

Li, Y.J., Chang, R.Z. and Zho, Y.T. (1991). Imbibitional injury in cultivars of soybean. Soybean genetics News

Letter, 18 : 68-73. Chinase Academi of Agricultural Science.

*Lips, H. and Roth-Bejerano, N. (1969). Science, 166 : 109.

Liso, R.; Calabrese, G.; Bitonti, M.B. and Arrigoni, O (1984). Relationship between ascorbic acid and cell division. Exp. Cell Res., 150 : 314-320.

Liso, R.; Innocenti, A.M.; Bitonti, M.B. and Arrigoni, O. (1988). Ascorbic acid induced progression of quiescent centre cells from G_1 to S phase. New Phytol, 110 : 469-471.

Liu, M.C. and Hadley, H.H. (1971). Relationship of Nitrate reductase activity to protein content in related nodulating and nonnodulating soybeans. Crop. Sci., 11 : 467-471.

Loewus, F.L. (1988). Ascorbic acid and its metabolic products. The biochemistry of plants, pp.85-107, Academic Press, Inc. SanDiego, California 92101.

Longden, P.C. (1973). Washing sugar beet seed. Institute International de Researches betteravieres, 6 : 154-162.

Lowry, O.H.; Rosenbrough, N.J.; Furr, A.L. and Randall, R.J. (1951). Protein measurement with the folin phenol reasent. J. Biol. Chem., 193 : 262-275.

Luedders, V.D. and Burns, J.S. (1979). Effects of broken seed coats on field emergence of soybeans. Agron J., 71 : 877-879.

Mackintosh Carol, Pauline Douglas and Cahrine Lillo (1995). Identification of a protein that inhibits the phosphorylated form of nitrate reductase from spinach (*Spinacia oleracea*) leaves. Plant Physiol., 107 : 451-457.

Madaan, T.R.; Aruna, Tyagi and H.C. Kapoor (1990). Effect of Gibberellic acid and Adenosine 3'5'-cyclic monophosphate on lipase activity in muskmelon (*Cucumis*

melo L.) seedlings. Indian J. of Exptl. Biology, 28 : 382-384.

Maehly, A.G. (1954). Methods in biochemical analysis. Ed. Glick, D.T. Interscience Publishers. Inc. New York, pp.385-386.

*Magdon, E. (1967). Physiol OK 101, Biochem. Keimung, Matev. Inst. Symp. (Publ. 1967), 1963 : 191.

*Mahadevapp and Raina (1975). In : World oil Seeds, pp.12.

Malik, C.P.; Singh, P.; Setia,R.C.; Setia,N. and Bhatia,D.S. (1986). Controlling plant biology and enhancing food production with phenolic compounds In : Hormonal Regulation of Plant Growth and Development Vol.III pp.371-415 (Ed.), S.S. Purohit, Bikaner, Agro Botanical Publishers.

*Mandal,N.C. and Biswas, B.B. (1970). Plant Physiol., 45, 4.

Mandal, N.C.; Burman, S. and Biswas, B.B. (1972). Isolation, purification and characterization of phytase from germinating mung beans. Phytochemistry, 11 : 495-502.

McDonald, M.B.; Jr. Vertucci, C.W. and Ross, E.E. (1988). Seed coat regulation of soybean seed imbibition. Crop. Sci., 28 : 987-992.

McDonald, M.B.; Jr.Vertucci, C.W. and Roos, E.E. (1988b). Soybean seed imbibition water absorption by seed parts. Crop Sci., 28 : 993-997.

*McCleary, B.V. and Matheson, N.K. (1974). Phytochemistry, 13 : 1747-1757.

*McCleary, B.V. and Matheson, N.K. (1975). Phytochemistry, 14 : 1187-1194.

*McCleary, B.V. and Matheson, N.K. (1976). Phytochemistry, 15 : 43-47.

Mclean, R.C. and Ivimey-cook, W.R. (1968). Text Book of Practical Botany. Longmans, Green and Co. Ltd., Edinburgh, Britain pp. 347-348.

MaLachlan, K.D. (1980). Acid phosphatase activity of intact roots and phosphorus nutrition in plants I. Assay conditions and phosphatase activity. Aust. J. Agric. Res., 21 : 429-440.

*Mapson, L.W. and Moustafa, E.M. (1957). Biochem. J. 62 : 248.

Marriott, K.M. and Northcote, D.H. (1975). The induction of enzyme activity in the endosperm of germinating castor bean seeds. Biochem. J., 152 : 65-70.

Marty, F.; Branton, D. and Leigh, R.A. (1980). Plant vacuoles. In EE. Conn. P. Stumpf eds. Biochemistry of Plants. Vol.I, Academic Press, New York. pp.625-658.

Masatoshi, N. and T. Asshi (1973). Increase in activities of membrane bound hydrolase in pea cotyledons during germination. Plant Cell Physiol., 14(6) : 1205-1208.

Matile, P. (1975). The lytic compartment of plant cells. Springer Verlag, Berlin.

Maxey, A.M. and J.C. Delouche (1980). Storage and viability of impermeable soybean seed. Agron. Abstr. American Society of Agronomy, Madison, WI. p.111.

Mayer, A.M. and Poljakoff-Mayber, A. (1963). The germination of seeds. MacMillan Co. New York, 71 : 81-88.

Mayer, A.M. and Poljakoff-Mayber, A. (1978). "Ther Germination of Seeds" Second Edition Pergamon Press Ltd., New York, pp.117-123.

Mehta, P.M. and Subhadradevi, M. (1974). Effect of ascorbic acid, maleic hydrazide and combination of Gibberellic acid, maleic hydrazide on germination and catalase activity in soybean. Botanique, 5(2) : 99-108.

Mehta, P.M.; Patel, R.J. and Rao, B.V.N. (1974). Effect of Gibberellic acid, Ascorbic acid, Morphactin and other combinations on seedling growth and enzymes of *Cicer arietinum* L. Geobios, 1 : 130-133.

Melzer, J.M.; Kleinhofs, A. and Warner, R.L. (1989). Nitrate reductase regulation. Effects of nitrate and light on nitrate reductase mRNA accumulation. Mol. Gen. Genet. 217 : 341-346.

Miha, Lyfi, J.P. (1968). Studies on catalase activity in almond seeds. Ann. Uni. Sci. Budapest Rolando Eotvos Nominate Sect. Biol., 9/10 : 305-308.

Miles, D.F.; D.M. Tekrony and Egli, D.B. (1988). Changes in viability, germination and respiration of freshly harvested soybean seed during development. Crop Sci., 28 : 700-704.

Minor, H.C. and Paschal, E.H. (1982). Variation in storability of soybeans under stimulated tropical conditions. Seed Sci. Technol., 10 : 131-139.

Mlodianowski, F. (1972). The occurrence of acid phosphatase in the thylakoid of developing plastids in lupin cotyledons. Z. Pflanzen. Physiol., 66 : 362-365.

Mohr. H.; Neininger, A. and Seith, B. (1992). Control of nitrate reductase and nitrite reductase gene expression by light, Nitrate and plastidic factor. Bot. Acta., 105 : 81-88.

Morse, W.J. (1950). In : Soybean and Soybean Products, Vol.1, p. 135, Interscience, New York.

Morohashi, Y. (1986). Patterns of mitochondrial development in reserve tissues of germinating seed : A survey. Physiol. Plant, 66 : 653-658.

Morohashi, Y.; Beweley, J.D. and Yeung, E.C. (1981). Biogenesis of mitochondria in imbibed peanut catyledons : Influence of the axis. J. Exp. Bot., 32 : 605-613.

Morita, Y. and Vagi, F. (1979). Characterization of Bamylase isozymes in soybean seeds by isoelectric focussing. Plant Cell Physiology 20(4) : 797-802.

Morre, R.P. (1973). Tetrazolium staining for assessing seed quality, "In Seed Ecology" Ed. Heydecker, W. Butterworths, London, 1973, pp.347-366.

Moustafa Seham, M. EL-Gandour, M.A. and Ghazi, Safia, M. (1982). Effect of seedtreatment with cycocel and Alar on growth and development of *Vicia faba* plant grown at different levels of soil salinity. Ain shams Uni. Fac. Agric. Res. Bull., 1625 : 1-17.

Mukherjee, J.J. and Paul, A.K. (1971). Respiration in germinating rice seeds. Proc. 58th Ind. Sci. Cong. Part-III, 23-9.

Mukherji, R.K.; Singh, R.A. and Khan, R.A. (1971). Tetrazolium staining as anindex of germination and vigour in soybean seeds. Bull. Grain Technol., 9 : 211-213.

Mukhiya, Y.K.; Shrotriya, N.; Joshi, J.K. and Singh, V.P. (1981). Salt tolerance of wheat, barley and soybean in respect of germination and pigment concentration. The Indian Journal of Agriculatural Sciences, 51(12) : 881-885.

Muntz, K.; Bassuner, R.; Lichtenfeld, C.; Scholz, G. and Weber, E. (1985). Proteolytic cleavage of storage proteins during embryogenesis and germination of legume seeds. Physiol. Veg., 23 : 75-94.

Muto, S. and H. Beevers (1974). Lipase activities in castor bean endospern during germination. Plant Physiol., 54 : 23-28.

Murphy, J.B.; T.L. Noland (1982). Temperature effects on seed imbibition and leakage mediated by viscosity and membranes. Plant Physiol., 69 : 428-431.

*Nanda, K.K. (1950). Curr. Sci., 19 : 22.

Nicholas, J.C.; Harper, J.E. and R.H. Hageman (1976). Nitrate reductase activity in soybean (*Glycine max* (L.)

merr) II. Energy limitations. Plant Physiol., 58 : 736-759.

Navarrow, F.; Rodriguez-Anguilera, J.C.; Alcain, F.J.; Burnon, M.I. and Navas, P. (1992). Ascorbate potentiates serum-induced S-phase entry of quiescent 3T3 cells. Protoplasma, 169 : 85-87.

Nielsen, N.C. (1984). The chemistry of legume storage proteins. Phil. Trans. R. Soc., (London) B : 287-304.

Nielsen, N.C. (1985). The structure and complexity of the 11S polypeptides in soybean. J. Am. Oil Chem. Soc., 62 : 1680-1686.

Nishikata, M. (1984). Trypsin like protease from soybean seeds. Purification and some properties. J. Biochem., 95 : 1169-1177.

Noat, G.; Crasnier, M. and Ricard, J. (1980). Ionic control of acid phosphatase activity in plant cells. Plant Cell Environ., 3 : 225-229.

Nonmi, H. and Boyer, J.S. (1990). Wall extensibility and cell hydraulic conductivity decrease in enlarging stem tissues at low water potentials. Plant Physiology (1990) 93(4) : 1610-1619.

Norris, W.E.Jr. and Foulds, E.C. (1961). Effect of gibberellic acid and 3 indole acetic acid and respiration of onion roots and seedling. Physiol. Plant., 14 : 293-299.

Nsowah, G.F. (1980). Effect of temperature on emergence, hypocotyl growth and dry matter production in soybeans [Abstract]. In world soybean research conference II abstract [Corbin, F.T. (editor)] Bowder, Colorado West View Press (1980). 59-60 [Ex]. Sci and Technology Univ., Kumasi, Ghana.

Oaks, A.M. Aslam and Boesel (1977). Ammonium and Aminoacids

at regulators of nitrate reductase in corn roots. Plant Physiol., 59 : 391-394.

Oathout, C.H. (1928). The vitality of soybean seed as affected by storage conditions and mechanical injury. J. Am. Soc. Agron., 2 : 837-855.

Okuda, Shin-ichi; Junkaneko and Toshiya Ogawa (1987). Increase in enzyme activities in embryonic axes of soybean seeds during germination. Agric. Biol. Chem., 51(1) : 109-114.

Ovcharov, K.E. and A. Akhmedov (1972). Alcohol Lactate malate and pyruvate dehydrogenase activity in maize seeds germinating under anaerobic conditions. Soviet Plant Physiol., 19 : 296-301.

Owen, F.V. (1927) Inheritance studies in soybean I. Ibid., 12 : 441-448.

Padma, A. and Reddy, G.M. (1975). Genetic and peroxidase isoenzyme studies of certain dwarf in *O. sativa*. Abst. from Genetics (Suppl.), 80(3) : Part I : 62.

Padmawati, T., Pratibha Devi and Kiranmali,V. (1992). Induced variability for different biological programme in soybean. J. Cyt. and Gen., 27 : 175-177.

Paleg, L.G. (1965). Physiological effects of gibberellins. Ann. Rev. Plant Physiol., 16 : 291-315.

Paleg, L.G. and Aspinal, D. (1981). The physiology and biochemistry of drought resistance inplants. Publ. Academic Press, Australia, pp.446.

Paleg, L.G.; Douglas, T.J.; Van Daal, A. and Keech, D.B. (1981). Proline, Betaine and other organic solutes protect enzymes against heat inactivation. Aust. J. Plant Physiol., 8 : 107-114.

Paliwal, K.V.; Maliwal, G.L. and Sharma, B.P. (1970). 1st annual report of ICAR scheme of salt tolerance of field crops, Udaipur Univ., Udaipur.

Palmer, C.E. (1985). Influence of ethephon on nitrate reductase, protein amino nitrogen and nitrate content of *Solanum tuberosum* L. Plant Cell Physiol., 26(3) : 407-417.

Palmino, E.P. and Juliano, P.B. (1972). Biochemical changes in the rice grain during germination. Plant Physiol., 49 : 751-756.

Papavizas, G.C. and Lewis, J.A. (1976). Acetone infusion of pyroxichor into soybean seed for the control of *Phytophthora megasperma* var. *Sojae*. Plant Dis. Rep., 60 : 484-488.

Papp, E.and Szabo, L. (1979). Germination temperature of soybean. Botanikai Kozlemenyek, 66(4) : 263-267.

Pardeshi, V.F.; Reddy, V.G. and Nalwandikar, P.K. (1989). Effects of different fungicides on seedling vigour and seed viability in soybean. Journal of Maharashtra Agricultural Universities, 14(1) : 33-36.

Parrish, D.J. and Leopold, A.C. (1977). Transient changes during soybean imbibition. Plant Physiology, 59 : 111-1115.

Parrish, D.J. and Leopold, A.C. (1978). On the mechanism of ageing in soybean seeds. Plant Physiol., 61: 365-368.

Patnum, F.W. (1953). Protein denaturation. The proteins Vol.8 (Ed. H.Nearath and K. Bailey) pp.807-892. Academic Press, New York.

Paul, A.K. and Mukherji, S. (1972). Changes in respiration rate of rice seedlings as affected by storage and viability and its possible relation with catalase and peroxidase activities during germination. Biol. Plant., 14(6) : 416-419.

Paul, A.A. and Southgate, D.T. (1978). The composition of foods (McCance and Widdowson, eds.). 4th edn. HMSO London and Elsevier/North Holland, Amsterdam.

Pazur, J.H.; Shadaksharaswamy, M. and Meidell, G.E. (1962). Arch Biochem. Biophys., 99 : 78-85.

Pelvi, Y.A.V. Narovo, N.N. and Drohysilliva, N.I. (1972). Nitrate reducing activity of plant peroxidase. Fiziologia Rasteini, 19 : 340-347.

Perl, M. (1986). ATP synthesis and utilization in the early stage of seed germination in relation to seed dormancy and quality. Physiol. Plant., 66 : 177-182.

*Peer's, F.G. (1953). Biochem. J., 53 : 102.

Pereira, L.A.G. and Andrews, C.H. (1985). Comparison of nonwrinkled and wrinkled soybean seed coats by scanning E Microscopy. Seed Sci. Technol., 13 : 853-860.

Pollock, B.M. (1969). Imbibition temperature sensitivity of lima beans controlled by seed moisture. Plant Physiol., 44 : 907-911.

Potts, H.C. (1978). Hard seeded soybeans. p.33-42. In Proc. 8th soybean seed Res. Conf., Chicago, 1L.12-14. Dec. American Seed Trade Association, Washington, DC.

Preseley, H.J. and Fowden, L. (1965). Acid phosphatase and isocitrase production during seed germination. Phytochem., 1(1) : 169-176.

Price, C.A. and Thimann, K.V. (1954). Dehydrogenase activity and respiration : a quantitative comparison. Plant Physiol., 29 : 495.

Pristley, D.A. and Leopold, C.C. (1979). Absence of lipid oxidation during accelerated ageing of soybean seeds. Plant Physiol., 63 : 726-729.

*Price, C.A. (1927). Nature (London), 180 : 148.

Price, C.E. and Ey, P.L. (1970). The distribution of some hydrolytic enzymes in the wheat seeds. J. Expt. Bot., 21(66) : 228-235.

Pristley, D.A. and Leopold, A.C. (1980). Alcohol stress on soybean seeds. Annals of Botany, 45(1) : 39-45.

Qi, Xiaoqun; Karla. Wilson and Anna L. Tan-Wilson (1992). Characterization of the major protease involved in the soybean ß-conglycinin storage protein mobilization. Plant Physiol., 99 : 725-733.

*Radin,J.W. and Trealese,R.N.(1976).Plant Physiol.,57 : 902.

Radin, J.W. (1977). Aminoacids interactions in the regulation of nitrate reductase induction in cotton root tips. Plant Physiol., 60 : 467-469.

Ram, C.; Balasimha, D. and Tewari, M.N. (1976). Effect of gibberellic acid and auxin on growth, sulphydryl content and peroxidase activity in *Phaseolus radiatus* L. seedlings. The Plant Biochem. Journal., 4(1) : 28-77.

Rao, J. (1973). Cytochemical localization of peroxidase in plant cells. Physiol. Plant, 28 : 132-133.

Rassini, J.B. and Lin, S.S. (1981). Efeitos de periodos de estiagens artificals durante estadios de desenvolvimen-toda planta no rendimento qualidade semente desoja (*Glycine max* (L.) Merr.). Argon., 17 : 225-237.

Raskin, T., (1992). Role of salicylic acid in plants. Annu. Rev. Plant. Physiol. Plant Mol. Biol., 43 : 439-463.

Rathore, T.R.; Ghildyal, B.P. and Sachar, R.S. (1981). Germination and emergence of soybean under crusted soil conditions. I effect of crust impedance to seedling emergence. Plant and Soil., 62(1) : 97-105.

Rayboy Victor; Stephen J. Hudson and David, B.Dickson (1985). Reduced phytic acid content doesnot have an adverse effect on germination of soybean seeds. Plant Physiol., 79 : 323-325.

Record, R.D. and Griffing L.R. (1988). Covergence of the endocytic and lytic compartments in soybean protoplasts. Planta, 176 : 425-432.

Renninberg, H. (1982). Glutathione metabolism and possible biological roles in higher plants. Phytochem., 21 : 2771-2781.

Roberts, E.H. (1972). Dormancy a factor affecting seed survival in the soil. In "Viability of seeds" Ed. Roberts, E.H. Chaponan and Hall London, pp.321-359.

Roberts, E.H. and Smith, R.D. (1977). Dormancy and the Pentose Phosphate Pathway in "The Physiology and Biochemistry of Seed Dormancy and Germination". (A.A. Khan ed.) pp.385-441. Elsevier North Holland, Biomedical Press, Amsterdam.

Roberts, E.H. and Ellis, R.H. (1980). Seed physiology and seed quality in soybean. In : R.J. Summerfield and A.H. Bunting (Eds.) Advances in Legume research Royal botanic garden, Kew p. 297-311.

Robinson, R.J. and Gibbs, M. (1982). Hydrogen peroxide synthesis in isolated spinach chloroplast lamellae. Plant Physiol., 70 : 1249-1254.

*Robertson, D.W.; Lute, A.M. and Kroeger, H. (1943). J. Am. Soc. Agron., 35 : 786-795.

*Roth-Bejerano, N. and Lips (1970). New Physiol., 69 : 165.

Roye, R.H., Vandana Sharma and Ram Narayan (1981). Hexokinase activity of germinating soybean (*Glycine max* L.). Indian Journal of Pl. Physiol., 24(2) : 109-112.

Rychter, A.;Rudnicki, R. and Lewak, S. (1971). Regulation of acid phosphatase activity by abscisic acid and gibberellin in apple seeds during stratification. Bull. Acad. Polon., 19 : 211-212.

Rychter, A.; Szkuthicka, K. and Zewak, S.C. (1972). Acid phosphatase and the low temperature requirement of apple seed stratification. Physiol. Veg., 10 : 671-676.

*Sachs, J. (1860). Physiologiche interauchungen liber der

Abhang 9 Keit der Keimung von der temperature. Jahrb. Wiss. Bot., 2 : 338.

Saha, H. and Gupta, K. (1993). Effect of LAB-150978 A plant growth retardant on sunflower and mung bean seedlings under saline stress. Indian J. Plant Physiol., 36(3) : 151-154.

Saio, K. and Watanabe, T. (1968). Observation of soybean food under electron microscope. J. Food Sci. Technol. (Japan), 15 : 290-298.

Saliim, N.H. and Todd, O.W. (1968). Seed soaking as a presowing, drought hardening treatment in wheat and barley seedlings. Agronomy J., 60 : 179-182.

Salunkhe, D.K.; Sathe, S.K. and Reddy, N.R. (1982). Legume lipids. In Chemistry and Biochemistry of food legumes. ed. S.K. Arora. p.52, New Delhi, Oxford and IBH.

*Sankhala, N. and Huber, W. (1975). Z. Pflanze Physiol. 75 : 392.

Sangeeta and Varshney, K.A. (1991). Effect of gibberellic acid, Maleichydrazide and cycocel on early growth and activities of some oxidoreductase in *Avena sativa* L. J. Exp. Biol., 29(1) : 80-82.

Sengupta, T.; Ghosh, B. and Sircar, S.M. (1977). Effect of growth substances on the IAA oxidase-peroxidase, activity during seed germination of rice *(Oryza sativa* L.). The plant biochemical journal, 4(1) : 28-33.

Sathiyamorthi, P. (1990). Identification of vanillic and p-coumaric acid as endogenous inhibitors of soybean seeds and their inhibitory effect on germination. Journal of Plant Physiol., 136(1) : 120-121.

Saxena, O.P.; Arya, V.; Pandey, N. and Maheshwari, D.C. (1988). Performance of the aged seeds under the influence of bioregulators. In proceeding of the plant

growth regulator society of America. [edited by Cooke, A.R.], 194-204.

Sayer, M.W. (1933). Soybean (*Glycine hispida*). Agric and Live Stock in India, 3(5) : 471-474.

Scandalios, J.G. (1969). Genetic control of multiple molecular forms of enzymes in plants. Ann. Rev. Biochem., Genet, 3 : 37-79.

Schopfer, P.; Bajracharya, D. and Plachy, C. (1979). Control of seed germination by abscisic acid. I. Time Course Action in *Sinopis alba*. Plant Physiol., 64(5) : 822-827.

Schobert, B. and Tschesche, H. (1978). Unusal solution properties of proline and its interaction with proteins. Biochem. Biophys. Acta., 541 : 270-277.

Sediyama, J.; Cardoso, A.A. and Vieirac, C. (1972). Testes prelimimars sobre of efeitos do returdamento da colheita da soja cultivar 'Vicoja'. Rev. Ceres., 19 : 306-310.

Seong, R.C.; Chung, H.J. and Hong, E.H. (1988). Varietal responses of soybean germination and seedling elongation to temperature and polyethylene glycol solution. Korean Journal of Crop. Sci., 33(1) : 31-37.

Senaratna, J. and B.D. Mckersie (1983). Dehydration injury in germinating soybean (*Glycine max* L. Merrill) seeds. Plant Physiol., 72 : 620-624.

Senaratna, Tissa, Bryan, D. Mckersie and Robert, H. Stinson (1985). Antioxidants levels in germinating soybean [(Glycine max) CV (maple arrow)] seed axes in relation ot free radical and hydration tolerance. Plant Physiol., 78(1) : 168-171.

Setty, T.A.S., Jagannath, V. and Rao, K.B. (1991). Effect of chemical seed treatment against seed borne fungi on

germination of soybean. Current Research University of Agricultural Science (Bangalore), 20(5) : 86-87.

Shannon, M.C. and Francois, L.E. (1977). Influence of seed pretreatments on salt tolerance of cotton during germination. Agronomy Journal, 69(4) : 619-622.

Sharma, S.M. and Tayal, M.S (1984). Effects of some phenols and GA_3 on, germination early seedling growth and α-amylase and protease activity of *Cicer arientinum* L. J. Indian Bot. Soc., 63 : 261-265.

Sharma, R.; Grewal, M. Kaur and Singh Gurbaksh (1986). Effect of etheral on lipid metabolism in soybean germinating under moisture stress. Indian J. Plant Physiol., 29(3) : 207-210.

Sharma Rajinder and Mohinder, K. Grewal (1987). Effect of etheral on some morphological and biochemical parameters in soybean germinating under moisture stress. Indian Journal Plant Physiol., 30(1) : 91-95.

Shiraishi, N.; Sato, T.; Ogura, N. and Nakagawa, H. (1992). Control by glutamine of the synthesis of nitrate reductase in cultured spinach cells. Plant Cell. Physiol., 33 : 727-731.

Shrivastava, H.S. (1980). Regulation of nitrate reductase activity in higher plants. Phytochem., 19 : 725-733.

Shull, C.A. (1920). Temperature and rate of moisture intake in seeds. Bot. Gaz., 69 : 361.

Siddique, S. and Krishnamoorthy, H.N. (1985). Effect of salinity and B-nine on *Vigna unguiculata* (cowpea) var. HFC42-1) In national seminar on plant physiology Varanasi, 21-23 Feb. 1985, pp.66-67.

Silva, H.C.; Braga, G.L.; Bianchi, M.D. and Rossi, E.A. (1990). Effect of germination on oligosoccharide and reducing sugar contents of Brazilian soybean cultivars. Alimentos Nutricao, 2 : 13-19.

Simon, E.W. (1974). Phospholipid and plant membrane permeability. Newphytol, 73 : 377-420.

Simotacchi, M.; Caro, A.; Fraga, G.G. and Puntarulo, S. (1993). Oxidative stress affect α-Tocopherol content soybean embryonic axed upon imbibition and following germination. Plant Physiol., 103(3) : 949-953.

Singh G. and Singh H. (1992). Maintenance of germinability of soybean (*Glycine max*) seeds. Seed Research, 20(1) : 49-50.

Singh, R.A. and Thakur, R.P. (1979). Effect of presowing soaking treatments on germination of soybean under moisture stress. Indian Journal of Ecology, 6(2) : 98-102.

Singh, R.A.; Tomar, C.S. and Thakur, R.S. (1981). Effect of cobalt and molybdenum on germination of soybean seeds. Seed Research, 9(1) : 73-75.

Sing, V.K.; Misra, H.S. and D.P. Misra (1987). Behaviour mitochondria in soybean (*Glycine max* L. var. Bragg) seeds during early stage of germination. Indian J. Exp. Biol., 25(4) : 270-272.

Sinha, S.K. and Nicholas, J.D. (1981). 'Nitrate Reductase' in physiology and biochemistry of drought resistance in plants. Academic Press, Australia, pp.167.

Smiciklas, R.D.; Mullen, R.E.; Carlson, R.E. and Knapp, A.D. (1989). Drought induced stress effect on soybean seed calcium and quality. Crop. Sci., 29 : 1519-1523.

Smith, C.R. (1954). Seed protein source-amino acid composition and total protein content of various plant seeds. Econ. Bot., 13(2) : 132-150.

Smith, A.K. and Circle, S.J. (1978). Soybean Chemistry and Technology, Vol.I. Proteins West Port, CT. AVI.

Smith, I.K.; Vierheller, T.L. and Thorne, C.A. (1989).

Properties and functions of glutathione reductase in plants. Physiol. Plant., 77 : 449-456.

Smolenska, G. and Lewak, S. (1974). The role of lipase in the germination of dormant apple embryos. Planta, 116 : 361-370.

Solomos, T. and Laties, G.G. (1975). The mechanism of theylene and cyanide action in triggering the rise in respiration in potato tubers. Plant Physiol., 55 : 499-504.

Srivastava, H.S. (1980). Regulation of nitrate reductase activity in higher plants. Phytochem., 19 : 725-753.

Srivastava, A.K.; Azhur, S. and Krishnamurti, C.R. (1972). Inhibition of germination in *Cicer arietinum*. Phytochem, 11 : 3181-3185.

Stafford, H.A (1965). Factors controlling the synthesis of natural and induced lignin in Phleum and Elodea. Plant Physiol., 40 : 844-851.

Stahmann, M.A. and Demorest, D.M. (1972). Changes in isozyme of host and pathogen following some fungal infections. Symp. Biol. Hung., 13 : 355-365.

Startseva, A.V. (1963). Kazanskogo filiola Akad Nauk SSSR, Ser. Biol. Nauk., 9 : 68.

Stewart, G.R. and Lee, J.A. (1974). The role of proline accumulation in halophytes. Planta, 120 : 279-289.

Stoddard (1945). Soybeans Advances in Agronomy. American society of agronomy, Vol.I, (ed. A.J. Norman), pp.107, Academic Press, Inc. Pub. New York.

*Stucky, D.J. (1976). Agron. J., 68 : 291-294.

Sugimoto, T.; Tanaka, K. Monma, M.; Kawamura, V. and Saio, K. (1989). Phosphoenolpyruvate carboxylase level in soybean seed highly correlates toits contents of protein and lipid. Agric. Biol. Chem. Tokyo., 53(3) : 885-887.

Sugiura, Y.; Kawabe, H.; Tanaka, H.; Fugimoto, S. and Ohra, A. (1981). Purification, enzymatic properties and active site environment of a novel manganese (III). containing acid phosphatase. J. Biol. Chem., 256 : 10664-10670.

Summerfield, R.J. and Roberts, E.H. (1985). Photothermal regulation of flowering in soybean. In proceeding world soybean research conference III. Ed. R. Colorado, pp.848-857.

Sushma Rani, Kuhad, M.S. and Nandwal, A.S. (1985). Effect of growth retardants on germination, early growth and biochemical changes in Raye (*Brassica juncea*) under stimulated stress. National Seminar on Plant Physiol., December, 15-17.

*Switzer, C.M. (1957). Plant Physiol., 32 : 42.

Szyrmer, J. and Szczepanska, K. (1982). Screening of soybean genotypes for cold tolerance during germination. Zeitschrift fur Pflanzenzuchtung, 88(3) : 255-260.

Tajima, Shigeyuki, Takeshi, Kanazawa, Ekotakeuchi and Vukioyamamoto (1985). Characteristics of ureate degrading diamine oxidase, peroxidase system in soybean (*Glycine max*) radicle. Plant Cell. Physiol., 26(5) : 787-796.

*Tavener, R.J.A and Laidman, D.L. (1972). Phytochem., 11 : 981-987, 2nd, 989-997.

Takashi, Y.; Hirofumi, V. and Miroshi, S. (1981). Gibberellic acid induced increase in activity of a particular isozyme of acid phosphatase in wheat half seeds. Plant Cell Physiol., 22(6) : 1023-1028.

Tamura, Jakeyoshi, Minamikawa, Takao, Koshiba, Tomokazu (1982). Acid phosphatase in cotyledons in germinating *Vigna mungo*. Plant Cell Physiol., 23(1) : 155-159.

Tang, W.; Diar, P.; Wilson, K.A. and Tan-wilson, A.L. (1993). Spatial distribution of protein accumulation in soybean seed, Phytochemistry-Oxford. Oxford, New York, pp.1289-1295.

*Tazakawa, Y. and Hirokawa, T. (1956). J. Biochem. Tokyo., 43 : 785.

Thanh, V.H. and Shibaski, K. (1978). Major protein of soybean seeds subunit structure of B-conglycinin. J. Agric Food Chem., 26 : 692-698.

*Theimer, R.R. and Rosnitschek, I. (1978). Planta., 139 : 249-256.

Tilden, R.L. and West, S.H. (1985). Reversal of the effects of aging in soybean seeds. Plant Physiol., 77 : 584-586.

*Todd, G.W. and Yoo, B.Y. (1964). Phyton, 21 : 61.

Tomes, L.J.; Teksomy, D.M.; Egli, D.B. (1981). The relationship of time, temperature and moisture content during accelerated aging to soybean seed germination. In Agronomy abstract 73rd annual meeting American Society (USA).

*Toole, E.H.; Hendricks, S.B.; Borthwick, H.A. and Toole, V.K. (1956). Annual Review Plant Physiol., 7 : 299-324.

Toole and Toole (1946). Soybeans Advances in agronomy, American Society of Agronomy, Vol.I, pp.107, Academic Press, New York.

Tregubenko, M.Y.; Fillippov, G.L. and Vishulvski, N.Y. (1973). Dykhatelnyl gazoobmen Kukuruzy Priuluchsheni uslovii Vyrocchivania (Respiratory metabolism of cor following improved growing conditions). DOKL. Ver. (Ordena lenia) Akad. S. Kh. IMV. lenina, 12 : 4-6.

Tully, R.E.; Musgrave, M.E. and Leopold, A.C. (1981). The seed coat as a control of imbibitional chilling injury. Crop. Science., 21(2) : 312-317.

Tu, S.I.; Brouillete, J.N.; Nagahashi, G.; Kumosinski, T.F. (1988). Effects of multivalent cations on c ell wall associated acid phosphatase activity. Plant Physiol., 88 : 514-520.

Ullah, A.H. and Gibson, D.M. (1988). Purification and characterization of acid phosphatase from cotyledons of germinating soybean seeds. Arch. Biochem. Biophys., 260 : 514-520.

Vaidehi, M.P. and Kadam, S.S. (1989). Soybean. In : Hand Book of world Food Legumes Nutritional Chemistry, Processing Technology and Utilization ed. D.K. Salunkhe and S.S. Kadam, Vol.III, pp.1-32. Boca Raton, FL. CRC Press.

Valio, I.F.M. (1986). The role of seed coat in early stage of soybean germination (*Glycine max* cultivar santrosa). Biol. Plant., 28(4) : 258-264.

Van Toai, T.T. and Bolles, C.S. (1991). Postanoxic injury in soybean (*Glycine max*) seedling. Plant Physiol., 97(2) : 588-592.

Vancnckelen, H.A.; Caubergs, R. and Degreef, J.A. (1976). Light controlled regulationof and ß-amylase activity incotyledons of *Phaseolus vulgaris*. Int. Physiol. Biochem., 84(1) : 202-204.

Vertucci, C.W. and Leopold, A.C. (1983). Dynamics of imbibition of soybean embryos. Plant Physiol., 72 : 190-193.

Vieira, R.D.; Tekrony, D.M. and Egli, D.B. (1992). Effect of drought and defoliation stress in the field on soybean seed Germination and Vigor. Crop. Sci., 32 : 471-475.

Vieira, R.D.; Tekrony, D.M. and Egli, D.B. (1991). Effects of drought stress on soybean seed germination and vigor. J. Seed Technol., 16 : 12-21.

Vincent, J.B., Crowder, M.W. and Averill, B.A. (1992). Hydrolysis of phosphate monoesters, a biological with multiple chemical solution. Trends Biochem. Sci., 17 : 105-110.

Vora, A.B.; Patel, H.C.; Vyas, A.V.; Patel, B.R. and Patel, J.N. (1975). Ascorbic acid turnover in water stressed Sesamum seedlings. Geobios, 2 : 186-187.

Vora, A.B.; Dehal, K.S. and Vyas, A.V. (1976). Effect of GA on catalase and peroxidase activity of *Pearl millet* (Bajra) seedling grown under restricted moisture level. Sci. and Cult., 42(9) : 479-481.

Vyas, D.N.; Patel, K.C. and Patel, R.D. (1965). Influence of pregermination chemical treatment on ascorbic acid and ascorbic acid oxidase of peanut. Naturwissenschaflen, 52 : 166-167.

Waggoner, P.I. and J.Y. Parlange (1976). Water uptake and water diffusivity of seeds. Plant Physiol., 57 : 153-156.

*Waley, S.G. (1966). Adv. Protein Chem., 21 : 1.

*Wanner, G. and Theimer, R.R. (1978). Planta, 140 : 163-169.

Webster, B.D. and Leopold, A.C. (1977). The ultra structure of dry and imbibed cotyledons of soybean. Am. J. Bot., 64 : 1286-1293.

*Wegg, S.M. and Townsley, P.M. (1983). Plant Cell Rep., 2 : 78-83.

Weston, Lesliea and Alan, R. Putnam (1986). Inhibition of legume seedling growth by residues and extracts of quackgrass (*Agropyron repens*). Weed Sci., 34(3) : 366-372.

Weil, J.; Pinsky, A. and Grossman, S. (1966). The proteases of the soybean. Cereal Chem., 43 : 392-399.

Will and deSaussure (1983). Seed soaking as a presowing, drought hardening treatment in wheat and barley seedlings. Agronomy J., 60 : 179-182.

Wilson, A.M. (1971). Amylase synthesis and stability in crested wheat grass seeds at low water potentials. Plant Physiol., 48(5) : 541-546.

Wilson, K.A.; Rightmire, B.R.; Chen, J.C. and Tan-Wilson, A.L. (1986). Differential proteolysis of glycinin and ß-conglycinin polypeptides during soybean germination and seedling growth. Plant Physiol., 82 : 71-76.

Willing, R.P.; Leopold, A.C. (1983). Cellular expansion at low temperature as a cause of membrane lesions. Plant Physiol., 71 : 118-121.

Wolf, F.T. (1968). Enzymatic activities in the leaves of wheat seedlings. Z. Pflanzenphysiol., 59-5 : 39-44.

Wolf, W.J. and Sly, D.A. (1967). Cryoprecipitation of soybean 11S protein. Cereal Chem., 44 : 653-658.

Wolf, W.J. and Cowan, J.C. (1975). Soybean as a food source. Boca-Raton, FL : CRC Press.

Woodstock, L.W. (1969). "Seedling growth" as a measure of seed vigour". proc. Int. Seed Test. Ass., 34 : 273-280.

Woodstock, L.W. and Combs, M. (1961). Application of the respiration test for seed vigour in peas. Proc. Ass. off. Seed Analyst. N. Am., 57 : 144-148.

Woodstock, L.W. and Grabe, D. (1967). Relationships between respiration during imbibition and subsequent seedling growth in *Zea mays* (L.). Plant Physiol., 42 : 1071-1076.

Woodstock, L.W. and Tao, K.J. (1981). Prevention of imbibitional injury in low vigour soybean embryonic axes by osmotic control of water uptake. Physiol. Plant., 51 : 133-139.

Woodstock, L.W. and Taylorson, R.B. (1981). Ethanol and Acetaldehyde in imbibing soybean seeds in relation to deterioration. Plant Physiol., 67 : 424-428.

*Wright, R.D.; Kanabus, J. and Cherry, J.H. (1974). Plant Sc. Lett., 2 : 347-355.

Yaklich, R.W. (1984). Moisture stress and soybean seed quality. J. Seed Technol., 90 : 60-67.

Yamuchi, F.; Sato, M.; Sato, W.; Kamala, Y. and Shibasaki, K. (1981). Isolation and identification of a new type of ß-conglycinin in soybean globulins. Agric. Biol. Chem., 45 : 2863-2868.

Yomo, H. and Varner, J.E. (1973). Content of the formation of α-amylase and protease in the cotyledons of germinating peas. Plant Physiol., 51 : 708-713.

Zairov, S.Z.; Erezhepor, A.E. and Valikhanova, G.Zh. (1983). Activation of isoperoxidase of wheat in the early of seed germiantion. Izv. Akad. Nauk. Kaz. SSR Biol., 4 : 16-20.

Zarska-Maciejewska, B.; Sinska, J.; Witkowska, E. and Lewak, S. (1980). Low temperature, gibberellin and acid lipase activity in removal of apple seed dormancy. Physiol. Plant, 48 : 532-535.

* Original not seen.