

BIBLIOGRAPHY

BIBLIOGRAPHY

- Abrams, R. (1975): Status of Research on Pigeon peas in Puerto Rico. International Workshop on Grain Legumes, pp. 141-147. Hyderabad, India. Int. Crop. Research Inst., Semiarid Trop. p. 350.
- Adams, C.A., Rinne, R.W. and Fjestad, M.C. (1980): Starch Deposition and Carbohydrase Activity in Developing and Germinating Soybean Seeds. Ann. Bot., 45: 577-582.
- Ahmad, T. (1938): The tur pod fly (Agromyza obtusamall), a pest of Cajanus cajan. Indian J. Agric. Sci., 8: 63-76.
- Akamine, T., Minamikawa, T. and T. Murata (1964): Enzymic Mechanism of Starch Synthesis in Ripening Rice Organ. Plant Physiology, 39: 371-78.
- Allen, O.N. and E.K. Allen (1940): C.F.W.O. Sutton (1983): Nodule Development and Senescence. In 'Nitrogen Fixation 3 Legumes' Ed. W.J. Broughton, Clarendon Press, Oxford. pp. 144-200.
- Allen, M.B. and D.I. Arnon (1965): Studies on Nitrogen Fixing Blue Green Algae II. Na requirement of Anabaena cylindrica. Physiol. Plantarum, 8: 653-660.
- Alten, Mineralsalzemburg, Kahlenso urea (1938): Slides Pflanze, 34: 73-75.
- Andrews, A.C. and D. Manjuli (1980): Effects of Plant Density and P & S Fertilizers on the yield of pigeon pea Cajanus cajan at Pakia. Thi. J. of Agric. Sci., 13(1): 9-14.

- ✓ Anonymous (1950): Cajanus cajan wealth of India (Raw Material, New Delhi Council of Scientific and Industrial Research. Wealth of India, 11(6):11.
- ✓ Arnon, D.T. (1949) Cu Enzymes in isolated chloroplast, polyphenol oxidase in Beta vulgaris. Plant Physiol., 24: 1-15.
- ✓ Asana, R.F., Riaz, B.K., Zain and M. Zain. UL Abedin (1968): Nutritional Studies on Some pulses commonly consumed in Pakistan. Pakistan Jour. Bot., 16(20): 24C.
- Asada and Kazi, Z. (1962): Formation of myo-inositol and phytin in ripening rice grains. Plant Cell Physiology. 3, 397-406.
- Austin, R.H. (1930): Effect of soil type and fertilizer treatment on the composition of the soybean plant, J. Am. Soc. Agroc., 22, 136-56.
- Ayub, A.T., (1974): Effect of calcium or sodium salinity of beans (Phaseolus vulgaris L.) J. Exp. Bot., 25: 245-252.
- Ayub, A.T. (1977): Some primary features of salt tolerance in Senna (Cassia autifolia). J. Exp. Bot., 28: 484-492.
- ✓ Bagley, B.W., Cherry, J.H., Rollins, M.L. and Altschul, A.M. (1963). Am. J. Botany, 50: 523. (Title?)
- Bangal, D.B., Deshmukh, S.N. and Patil, V.A. (1983): Contribution of pod wall in grain development of chickpea (Cicer arietinum L.) as influenced by foliar application of growth regulator and urea, Indian J. Pl. Physiol., 26: 292-295.
- Batt, T. and H.W. Woolhouse (1975): Changing activities during senescence and sites of synthesis of photosynthetic enzymes in leaves of the labile Perilla frutescens (L.) Bitt. J.

Exp. Bot., 26(93): 569-579.

Bacharach, A.L. (1940): The Distribution of nicotinic acid in human and animal foods, Nutr. Abstr. and Rev., 10: 459-565.

Barrat (1921): The tropical crops cited by Pathak, G.N. Pulse crops of India (1970): Chapter 2, red gram 46.

Bathurst, N.O. (1954): J. Exp. Bot. 5: 253-254.

Barnett, N.M. and A.W. Naylor (1966): Plant Physiology, 47: 591-594.

Basu, K.P. and M.C. Malakar (1939): Biological value of proteins of Bangal grams (Cicer arietinum) and arhar (Cajanus cajan) J. Indian Chem. Soc. 16: 4272-432.

Bates, L.S., Waldren, R.P., Teare, I.D. (1973): Rapid determination of free proline for water stress studies, plant and soil, 39, 205-207.

Babu, V.R. and S. Kumar (1979): Seed germination and early seedling growth of Cicer arietinum cultivar stressed condition. J. Indian Bot. Soc., 58 (2): 140-148.

Bangerth, F., Dilley, D.R. and Dewey, D.H. (1972): J. Am. Soc. Hort. Sci., 97: 679-682.

Bachroodi, V. (1973): Function of naturally occurring polyamines Academic Press, New York, 96:107.

Bain, J.M. and Mercer, F.V. (1966): Subcellular organization of the developing cotyledons of pisum sativum L. Aust. Biol. Sci., 19: 49-67.

Baumeister and Schinidth (1962): On the role of sodium in plant metabolism. Flora Allg. Bot. Zeitung, 152: 24-26.

- Baker, J.G. (1987): Flora of Mauritius and Seybelles L. reeve and Co., London.
- Bair, D. and Latzko, E. (1976): Biophys. Acta. 396: 141-148.
- Beever, L. and Poulson, R. (1972): Protein Synthesis in Cotyledons of Pisum sativum L.I. Changes in cell-free acid incorporation capacity during seed development and maturation. Pt. Physiol. Lancaster, 49: 476:481.
- Beever, L. and R. Storey (1976): Glutamate synthetase in developing cotyledons of Pisum sativum Plant Physiol., 57: 862-866.
- Beohar, A.B.L. and P.K. Nigam (1972): Correlation Studies in Arher, (Cajanus cajan L.) Millsp. JNKW.Res.J., 6(1): 58.
- Ben-Zioni, A.: Vaadia, V. and Zips, S.H. (1971): Physiol. Plant., 24: 255-290.
- Bengton, C.G.B., Klockare, Klockare, R., Larsson, S. and C. Rundquist (1978). Physiol. Pt., 43: 205-212.
- Beever, L. and R.H., Hageman (1969): Nitrate Reduction in Higher Plants A Rev. Pl. Physiol., 20: 495-522.
- Bharucha, F.R. and M.V. Dabholkar (1959): Physiological Studies on calcicolous plant of Bombay. IV. J. Univ. Bombay, 27: 46-73.
- Biswas, H.C. (1943): Investigation on the saponin content of Cajanus indicus. Sci. and Culture., 9: 165.
- Bhambri, Neeru; Malik (1983): Biochemical changes in developing seeds and pods of phaseolus vulgaris. Chemical Abstract, 98: 50520 'P'.

- / Bhivare, V.N. (1984): Physiological studies in pulses: A Ph.D. Thesis submitted to the Shivaji University, Kolhapur (India).
- / Biale, J.C. (1960): In 'Handbuch der Pflanzen Physiologie', C.W. Ruhland, edi. Vol. XII. pp. 347-359.
- / Bidhan Chandra (1987): Studies on the physiology of seed development in Urad bean at different days after anthesis. Ind. J. Pl. Phys., 30(2): 199-201.
- / Bission, C.S. and Jones, H.A. (1932): Changes accompanying fruit development in the garden pea. Plant Physiology, 7: 91-105.
- Biswas, H.C. (1943): Investigation on the Saponin content of C. indicus. Sci. and culture, 9: 165.
- / Black, A. (1965): Methods of soil analysis part 2 chemical and microbial properties Am. Soc. Agron. Inc. Medison, Wisconsin.
- Black, C.C. (1973): Photosynthetic 'C' fixation in relation to net CO_2 uptake. Ann. Rev. Plant Physiology, 24: 253-286.
- Boulter, D. and Davis, O.J. (1968): Flinn and Pate, 1968; Mumtaz, 1973; Mumtaz (1976), Pate (1977).
- / Bogges, Samuel, F.; D.E. Koeppe, C.R. Stewart (1978): Oxidation of proline by plant mitochondria Biol. Abst., 66(12): 73454.
- / Boulter, D. (1986): Regulation of storage protein synthesis and deposition in developing pea seeds. Crop. Plant Physiol. Abst., 12(12): 4417.
- / Brag, H. (1972): Plant Physiol., 26: 250-257.
- / Broyer and Stout (1959): The micronutrient elements. Ann. Rev. Plant Physiol., 10: 277-300.

- Brownwell and Crossland (1972): The requirement for Na as a micronutrient by species having the C₄ dicarboxylic photosynthetic pathway. Pt. Physiol., 49: 794-797.
- Brownwell, P.F., and J.G. Wood (1957): Na as an essential element for Atriplex vesicaria Heward, Nature, 179: 635-636.
- Broyer, T.C.; A.B. Carlton; C.M. Johnson; P.R. Stout (1958): Chloride micronutrient element for higher plants. Plant Physiology 29: 526-532.
- Broadbent, F.E. (1973): Sources and sinks of nitrate in soils. In process of the 1st Annual Trace contaminants conference. Oak Ridge National Laboratory, National Science Foundation, Washington D.C. 88-119.
- Broadbent, F.E. (1968): Turnover of nitrogen in soil organic matter Pontificiae Scientiarum Scripta Varia. 32: 61-88.
- Bryan, J.K. (1976): Amino acid biosynthesis and its regulation in 'Plant Biochemistry', Ed. J. Bonner, J.E. Varner. Academic New York. pp. 525-597.
- Britikov, E.A.; Schrauern, J. and H.F. Linskens (1970): Acta Bot. Neerl., 19: 515-520.
- Britikov, E.A.; Vladimirtseva, S.V. and N.A. Musatova (1965): Eiz. Rast., 12: 953-967.
- Burridge (1964): The levels of N,P,K, Ca and Mg in leaves of cocoa (*Theobroma Cacao L.*) as affected by shade fertilizer, irrigation and season. Ann. Bot., 28: 401-417.
- Burns, R.C. and R.W.F. Hardy (1975): Nitrogen Fixation in Bacteria and Higher Plants. Springer-Verlag, New York.

- Burstrom, H. (1943), 45). The nitrate nutritor of plants: A general survey of occurrence and assimilation of nitrate. Kgl. Lantbruks Hoegsk. Ann., 13: 1-89.
- Carl Erik Danielson (1960): Sugars and Starch in ripening peas. Chemical Abst., 54: 1-2824.
- Cadolle (1904): Committee on Technology Innovation Board on Sci. and Tech. for International Development, Commission of International Relations, Washington.
- Campbell, W.H. (1988): Nitrate reductase and its role in nitrate assimilation in plant. Physiol. Plantarum., 74: 214-219.
- Casterel (1981): Centre for the Application of Science and Technology for Rural Development (1981), Review of experimental cultivation and propagation of winged bean during 1979-80.
- Gerry, K. (1978): Comparative nutritional and clinical aspects of the winged bean. In Proc. 1st International Sympon. Developing Potential of the Winged Bean, Jan. 1978, Philippines, Manila, p. 281.
- CENA (1987) - Piracicaba, Sao Paulo, Brazil. Crop Physiol. Abst., 13(6).
- Chavan, J.K., Kadam, S.S., Ghonsikar, C.P. and Salunkhe, D.K. (1979b): Removal of tannins and improvement of in vitro protein digestibility of sorghum seeds by soaking in alkali: J. Food Sci. 44: 1319.
- Chakravarti, S.C. (1964): Vernalization and growth Indian Agric., 8: 68-70.

Chattarjee, S.R., Pokhriyal, T.C. and Abrol, Y.L. (1977): Proline content and amino acid composition of developing seeds of bengal gram (*Cicer arietinum*) Plant Biochem. J., 4: 62-71.

Chang, Hsin-Hsing and Ching-Yung Zicu (1984). Natl. Sci. Counc. Mon. 11 (12 Part 3): 503-572.

Choudhari (1968): Pulses research in India. Indian Foramina. Nov. 25-26..

Chavan, V.M. and N.R. Bhat (1949): Increased production of pulses through breeding, introduction and cultural methods. Poona Agric. College Mag. 40: 18-28.

Chavan, P.D. (1980): Physiological Studies in Plants (Physiological studies in *Eleusine coracana* Goertin.) Ph.D. Thesis submitted to Shivaji University, Kolhapur (India).

Clements, R.L. (1970): Protein patterns in fruits in biochemistry of fruits and their products, Academic Press, London, 1: 159-178.

Claydon, A. (1975): Winged bean, a food with many uses. Plant food for man, 2: 49.

Clo-ver (1937): Pulses make a comeback Indian Foramina June 26-27.

Claydon, A. (1978): The role of the winged bean in human nutrition. In: Poc. 1st International symposium on developing the potential of winged bean. Phillipines, Manila. pp. 263-280.

Crookston et al., 1974; Flinn and Pate (1970); Schlesier 1977; Pate et al. (1977) - Characterization of the bean pod as a photosynthetic organ - Crop. Sci. 14, 708-713.

- Campbell, N.A. and Thomson, W.W. (1977): Plant Physiol. 60: 635-639.
- ✓ Daniel (1956): Stoffwechsel und Mineralsalzernährung einzelliger Grünalgen
III. Atmung und oxydative assimilation von chlorella flora
(Jena), 143: 31-66.
- ✓ Data, E.S. and H.K. Pratt (1980): Trop. Agric. (Trinidad) 57(4):
309-317.
- Dalal, R.C. (1980): Nutrient element concentration and dry matter production by field grown pea (Cajanus cajan (L.) Mill s.p.) Tropical Agriculture, 57(2): 107-112.
- ✓ Davis, S.D.; Bill, D. (1978): Occurrence of A-amylase in the axis of germinating peas. Plant Physiol., 60(4): 513-517.
- ✓ Danuta, W. (1976): Functional and Structural changes in chloroplasts of senescent tobacco leaves. Acta Soc. Bot. Pol., 45(3): 341-352.
- ✓ Darlington, C.D. (1969): The silent millennia in the origin of agriculture. In the Domestication and Exploitation of Plants and animals, eds. P.J. Veko and G.W. Dimbleby. Duckworth, London, pp. 67-72.
- Delwiche, C.C. (1970): Sci. Amer. 223(3): 136 - cited by Bonner, J. and J.E. Vamer (1976). In plant Biochemistry Third Edition Academic Press, New York.
- ✓ Deshpande, R.S. (1981): Physiological Studies in Pulses of Maharashtra. Ph.D. thesis submitted to the Shivaji University, Kolhapur (India).
- Dekort, C.A.D., Bartelink, A.K.M. and R.R. Schurmans (1973): Insect. Biochem. 3: 11-17.

and C.C. Mack 184

- Yian-Peng. Sci. Jean Sung (1989): Sucrose metabolism in lime bean seed. Plant Physiol. 89(4): 1106-1116.
- Dodge, J.D. (1970): Changes in chloroplast fine structure during the autumnal senescence of Betual leaves. Ann. Bot. (London), 34 (137): 816-824.
- Drosdoff, M. and C.D. Nearpass (1948): Quantitative micro-determination of magnesium in plant tissue and soil extracts. Annual Chem. 20: 673-674.
- Dubroca, E. and Bory, G. (1983): Biochem. Syst. Ecol. 9(4): 283-288, G.M. (1982), Crop Science 22: 459-463.
- Durie, R.A.; Schaffer, H.M.S. and D.J. Swaine (1965) - Technical Communication of Commonwealth Sci. Industr. Res. Org. 47: 33-36.
- Doughall, D.K. and J. Bloch (1976): A survey of presence of glutamate synthetase in Plant Cell Suspension Cultures. Can. J. Bot. 54: 2294-2297.
- Dunphy, E.J. (1972): Soybean, water-soluble carbohydrate, nutrient, yield and growth pattern responses to phosphorus and potassium fertility differences. Ph.D. Thesis. Iowa State University Library, Ames.
- Danielson (1959): Sugars and Starch in Ripening Peas. Chemical Abst., 54: 692.
- Dunn, G.C. (1974): A model for starch breakdown in higher plants. Phytochemistry, 13: 1341-11346.

✓ Dutt, B.K. and A.G. Thakurta (1939): Catalase activity of cajanus seed at different stages pre-resting, resting and post-resting. Bibliograph Pigeonpea., 93 (103-1152).

EK Penyong, T.E. and R.L. Barchers (1978): Nutritional aspects of the winged bean. In Proc. 1st International Symposium on Developing Potential of the Winged Bean, Philippines, Manila, p. 300.

Elias, L.G.; Dernades, D.G. and Bressani, R. (1979): Possible effects of seed coat polyphenols on the nutritional quality of bean protein. J. Food Sci., 55: 524.

Ellis, M.A.; Paschal and P. Bwell (1977): The effect of maturity and foliar fungicides on pigeonpea seed quality. Plant Dis. Res. 16(12): 1006-1008.

Embleton, T.W., Jones, W.W. and Platt, R.G. (1975): Hort. Science, 10: 48-50.

Engler (1915): Paternoster, Square Pf Lanzenwelt Afrikas 3: 665-666.

Epstein, E. (1965): Mineral metabolism. In: Plant Biochemistry. J. Bonner and J. Varner (Eds.) pp. 438-456. Academic Press, New York.

Epstein (1972): Mineral Nutrition of Plants. Principles and Perspectives, John Wiley and Sons Inc., New York, London, Sydney, Toronto.

Erdman, J.W. Jr. (1981): Bioavailability of trace minerals from cereals and legumes. Cereal Chem. 58, 21.

✓ Evans (1982): Nitrate reductase activity in chickpea: A Preliminary investigation, International Chickpea News Letter, 6: 22-23.

- Evans, C.E., Lathwell, D.J. and Mederski, H.J. (1950): Effect of deficient or toxic levels of nutrients in solution on foliar symptoms and mineral content of soybean leaves as measured by spectrographic methods. Agron. J. 42, 25-32.
- Fairchild, D. (1944): Garden islands of the great East. C. Scribner's Sons, New York.
- Favoretto (1979): Effect of cutting time on the yield and chemical compound of pigeon pea (*Cajanus cajan*) dentifica Jabaticabal 7(3): 505-510.
- Fader, G.M. (1986): Transport of assimilate into the soybean fruit and seed growth. Dissertations Abst. International, 12(6): 2013.
- Fernandez Tarango (1979): Starch degradation during germination of *Cicer arietinum* L. seeds. Rev. Esp. Fisial., 67(1) 522.
- Ferry, J.F. and Ward (1959) - Fundamentals of plant Physiology. Macmillan Company, New York.
- Fidler, J.C. (1960): Pflanzen Physiol. Vol. XII pp. 347-359.
- Flinn and Pate (1968): Biochemical and Physiological changes during maturation of fruit of the field pea (*P. arvensis*). Ann. Bot. 32: 479-495.
- Flinn and Pate (1970): A quantitative study of Carbon transfer from pod and subtending leaf to the ripening seeds of the field pea, 21: 71-82.

Fox et al. (1964): Studies on the proteins of seeds of the leguminase - I. Albumin. Phytochemistry, 3: 417-419.

Folin and Denis (1915): A colorimetric estimation of phenols (and phenol derivatives) in urine J. Biol. Chem., 22: 305-302.

Fowden, L. (1954); F.D.G. Hill; Cottingham and G.P. Lloyd Jones (1977): Translocation of Nitrogenous compounds in plants. Academic Press New York, pp. 397-405.

Fountain, D.W.; H.A. Outred; J.M. Holdsworth; R.G. Thomas (1988): Seed development in Phaseolus Vulgaris L. LV Seminole Plant Physiol., 89(1) 333-340.

Firewood crops Pub. NAS (1980) shrubs, trees, species for energy production. Report of Hoc. Panel of the Advisory Committee.

Garcia, V.V. and J.K. Palmer (1980): Proximate analysis of five varieties of winged bean Propocarpus tetragonolobus (L) DC J. food Technol. (U.K.) 15: 469.

Gavazzi, G.; Nava-Racchi, M. and C. Tonelli (1975): Theor. Appl. Genet., 46: 339-346.

Givan, C.F.; Givan, A and R.M. Leech (1970): Photoreductions α -Ketoglutarate to glutamate by Vicia faba Chloplasts. Plant Physiol., 45: 624-630.

Govindrajan, S.V. (1954): Role of micronutrient in causing diseased condition in plants and (b) increasing crop product, 10th meeting Board of Agri. and Animethesh crop and soil wing. Indian Council Agric. Res. Bangalore.

- Gopal, G.R. and G.R. Rao (1982): Salt effect on in vivo activity of nitrate reductase in peanut (Arachis hypogaea L.) Seedlings Turrialba, 32 (2) 216-218.
- Goldstein, J.L. and T. Swain (1963): Changes in Tannins in ripening fruits. Phytochemistry, 2: 371.
- Griffiths, L.A. (1960): Comparative study of seed polyphenols of Theobroma. Chemical Abst., 54: 6205-9372.
- Hartt, C.E. (1972): Plant Physiol., 44: 1461-1469.
- Hartt, C.E. (1929, 1934): Some effect of Potassium upon the amounts of protein and amino forms of N. sugars and enzyme activity of sugarcane, Plant Physiol., 10: 277-300.
- Haskins, F.A. and S.C. Downs (1961): Activities of acid phosphatase, peroxidase and polyphenols in etiolated shoots from control and irradiated maize seeds. Chemical Abst., 55: 11945-15000.
- Harris, J. (1976): Plant Physiol. (Suppl.), 67 (4): 115.
- Hawk, P.B.; B.L. Oser and W.H. Summerson (1948): Practical physiological chemistry pb. The Blankiston Company, U.S.A.
- Hageman, R.H.; C.F. Crosswell and E.J. Hewitt (1962): Reduction of nitrate, nitrite and Hydroxylamine to ammonia by enzymes extracted from higher plants. Nature (Lond.), 193: 247-250.
- Hardy, R.W.F. and M.D. Havelka (1975): Nitrogen fixation research. A Key to World Food Science, 188: 633-643.
- Harvey and Hadley, 1975a, 1976b; Harvey et al. (1976): Sugars in seedling organs and isolated cotyledons of Lupinus luteus L. Bull Soc. Amis. Sci. Lett. Poznan SER D Set Biol. 67 (2): 11601.

- Herklotz (1972): Kobely and Steels (1976).
- Helbeak (1966): Pre-Pottery neolithic farming at Beidha Palest Exploro.
98: 61-66.
- Hewitt, E.J.; S.C. Agarwala and A.H. Williams (1957): Regulation
of nitrite assimilation. Academic Press, New York, pp.
255-282.
- Hewitt, E.J. (1966): Sand and water culture methods used in the
study of plant nutrition, 2nd edn. Common W. Bur. Host.
Plant Crops. Tech. Comm. 22.
- Heber, V. and P. Purchold (1977): Substrate and product fluxes
across the Chloroplast envelope during bicarbonate and
nitrate red. Proc. 4th Int. Corner, on photosyphiosis pp.
107-118.
- Hilgeman, R.H.; Dunlop, J.A. and Sharples, G.C. (1967): Proc.
Am. Soc. Hortic. Sci., 90: 1101-115.
- Hill, J.E. and R.W. Breidenbach (1974): Accumulation of the major
protein component during seed development and maturation,
Pt. Physio. 53: 747-751.
- Hildbrand, D.F. and T. Hynowitz (1981): Electrophoretic separation
and properties of winged bean seed trypsin inhibitor. J.
Sci. Food Agri., 32: 443.
- Hipsley, E.H. and N.E. Kirk (1965): Studies on dietary intake and
expenditure of energy by new Guineans Tech. Paper No.
147.
- Hipsley, E.H. and Clement, F.W. (1947): Report of the New Guinea

Nutrition Survey Expedition, 1947, Canberra, Deptt. Ext.
Terri. pp. 120-274.

Hocking, P.J. and J.S. Pate (1978): Mobilization of minerals to developing seeds of legumes Biol. Obst., 65(7): 37249-49664.

Hocking, P.G. and J.S. Pate (1977): Ann. Bot., 41: 1259-1278.

Hocking, P.G., Kudo, J. and J.S. Pate (1980): Aust. J. Bot. 29: 1-18.

Hoffmannowa, Aleksandra and Lubomira Domka (1979): Soluble Sugars in Seedling Organs and Isolated cotyledons of Lupinus Luteus L. Bull. Soc. AMIS. SCI. Lett. Poznan ser. D. Sci. Biol., 67(2): 1145.

Holden, M. (1973): Chloroplast pigments in plants with C₄ dicarboxylic acid pathway of photosynthesis, Photosynthetica, 7(1): 41-49.

Humble, G.D. and Roschke, K. (1971): Plant Physiol. 48: 447-453.

Huang, A.H.C. and A.J. Cavalieri (1979): Pt. Physiol. 63: 531-535.

Hutchinson, J. (1964): The genera of flowering plants (Angiospermae) Dicotyledons Vol. 1, Oxford, Clarendon Press, p. 442.

Hyde, E.O.C. (1954): The function of the hilum in some Papilionaceae in relation to the ripening of the seed and the permeability of the testa Ann. Bot. 18, 241-256.

Hymowitz, T. and Boyd, J. (1977): Origin, ethnobotany and agricultural potential of winged bean P. tetragonolobus. Eco. Bot. 31: 180-188.

- Ingle, J. Beitz, D. and Hageman, R.H. (1965): Changes in composition during development and maturation of maize seeds. Pt. Physiol., Lancaster, 40: 48-53.
- ✓ Iskander, F.Y. (1987): Maturation State and mineral content of soybean. Crop. Physiology Abst., 13(9), 2930.
- Jaffe, W.G. and R. Korte (1976): Nutritional characteristics of the winged bean. Int. Nutr. Rept. 14: 449.
- Jambunathan, R. and Singh, R.V. (1980): Studies on desi and Kabuli Chickpea (Cicer arietinum L.) cultivars 1. Chemical Composition. In Proc. International Workshop on Chickpea Improvement, ICRISAT. Hydraba. India, pp. 61-66.
- Jambunathan, R. and Singh, U. (1981): Studies on desi and Kabuli chickpea (Cicer arietinum L.) cultivars. Mineral and trace element composition J. Agric. Ford Chem. 29: 1091-1093.
- Jacobson, L.R.; Hamapel, R.J.; Schaedle, M. and More (1961): Effects of root to solution ratio in iron absorption expt. Plant Physiology, 36: 65.
- ✓ Jepson, R.C. and R.R. Johnson, Hadley, H. (1978): Variation in mobilization of plant nitrogen to grain in nodulation and non-nodulation soybean genotypes. Crop. Sci. 18: 1058-1062.
- Jones, R.W. and R.W. Sheard (1972): Nitrate reductase activity, Phyto chrome mediation of induction in etiolated peas. Nature, New Biol. 238: 221-222.
- ✓ Joshi, G.V. (1976): Studies in photosynthesis under saline conditions,

P.L. 480 Project report submitted to Shivaji University, Kolhapur (India).

Joseph, A. Lazslo (1990): Developmental Changes in mineral contents at soybean seed coats and embryo. Plant Physiology, 93:1.

Kathiiresan, K.; Gnamarethinam, J.L. and Daniel. I (1985): National Seminar on Plant Physiology, Varanasi, Abstr.No. 107: p. 65.

Kartz and J. Mayers (1965): Nutritional growth of several blue-green algae. Am. J. Bot., 42: 282-287.

Kalenov, P.A.; Blokhim; Baslichn; Kult; Zadachami gel (1983): Changes in protein complex in ripening seeds of soybean varieties with different vegetative periods. Chemical Abst., 98: 1816100-197181.

Kadam, S.S., Salunkhe, D.K. (1981): Nutritional aspects of winged bean Legume Research, 4: 33.

Kaplan (1965): Archaeology and domestication in America. Phaseolus. Econ. Bot., 10: 358-368.

Kamat, M.N.; P.G. Patwardhan; V.G. Rao and A.V. Sathe (1971): Fungi of Maharashtra. The Mahatma Phule Krishi Vidyapeeth, Maharashtra State, pb. No. 1.

Kadam, S.B., L.S. Kuste, K.M. Lawande and D.K. Salunkhe (1982): Changes in Chemical Composition of winged bean (Psopbocarpus tetragonolobus) during seed development. J. Food Sci. 47: 2051.

Karadge, B.A. and P.D. Chavan (1980): Physiological studies in salt tolerance in Groundnut. UGC. Project (1978-80) (8828/77), p. 87

- Katsumi, M. and M. Fukuhara (1969): The activity of α -amylase in the shoot and its relation to Gibberelin induced elongation. Plant Physiology., 22: 68-75.
- Khokar, P.K. and Warsi, A.S. (1987): Nutrient uptake studies in gram (C-235) seeds and forms CAB Prompts Series Chickpea and Pigeonpea., 1(2): 178.
- Khan, A.A. (1983): The physiology and biochemistry of seed development Chem. Abst. 96(6): 28251-89935.
- Khatra, S.S.; Kumar, V.; Sindhu, P.S. (1987): Pattern of dry matter accumulation in developing fruit parts of early and late maturing Pigeon pea. Biologia Plantarum, 13(5): 1471.
- Khanna, Chopra and Sinha (1976) - Importance of fruit wall in seed yield of pea (Plumus sativum L.) and mustard (Brassica). Indian J. Exp. Biol. 14: 159-162.
- Kidd, F. and West (1930): Respiration of fruits. Nature (London), 135: 327-330.
- Kidd, F. (1935): Respiration of fruit. Nature (London) 135: 327-330.
- King, N.J. and S.T. Bayley (1965) - J. Exp. Bot. 16: 294-303.
- Kirolos, M.A.; El-Hafeez, S.A. (1986): Effect of orobancheccrenate parasitizing Vicia faba plants on the activity of peroxidase and polyphenol oxidase in different parts of the host parasite. Crop. Physiol. Abst., 12 (5): 1708.
- Konno, H.; Yamaya, T.; Yamasaki, Y. and Katrumoto, H. (1984): Plant Physiol. 76: 633-637.

Kokro, J.G. (1976): Effect of seed soaking in water and concentrated sulphuric acid on emergence and growth of winged bean (*P. tetragonolobus*). I.T.T.A. Trainee report No. 56, Ibadan, grain Legume, Improvement Programme, International Institute of Tropical Agriculture.

Kramer, P.J. and Kozlowski, T.T. (1979): Physiology of woody plants, Academic Press, New York.

Krishna, T.; Gopala, R.K. Mitra and C.R. Bhatia (1978) Seed glutelins of *C. cajans*, Plant Foods Hunutritive Biol. Absts. 65(10): 49665-62080.

Krishna, T.G., R.K. Mitra and C.R. Bhatia (1977): Seed Globulins of *Cajanus cajan*, Qul. Plant Food Hum. Nutrition., 27(3/4): 313-326.

Kumar; K. Ganesh, L.V. Ventataraman (1978): Chickpea seed proteins, Modification during germination. Phytochemistry (OXF), 66(7): 37249-49664.

Kumar Rao, J.V.D.K. and P.J. Dart (1987): Nodulation, Nitrogen fixation and nitrogen uptake in pigeonpea (*Cajanus cajan* (L) Mills p) of different maturity groups. Plant and Soil 99, 255-266.

Kumar, V., Singh, M., Narwal, R.P. (1987): Effect of different sources and leaves of Zn with 'S' on uptake and utilization of 'P' in Soybean (*Glycin max* (L) Merr). Crop. Physiol. Abst. 13(7): 2355.

Kumar, J. Gavda, C.L.L.; Saxena, N.P., Sethi, S.C., Singh, V.

- and Sabrowat, K.L. (1983): Effect of salinity on protein content and seed size of Chickpea (Cicer arietinum L.) 52: 82-83.
- Kumar, K.O., V.G. and A.D. Berezovik (1962): N. containing compounds of the vegetative pods and seeds of V. faba in course of development, Chemical Abstract, 56: 5126h.
- Lovell, P.H. and Lovell, P.J. (1970): Fixation of CO_2 and export of photosynthate by the carpel in P. sativum - Physiol. Plant., 23, 316-322.
- Loureiro, J. (1988): Hocker (1978): Flora Cochinchinensis Tomus I. Ulyssipone.
- Lolas, G.M. and P. Markakis (1975): Phytic Acid and other Phosphorus Compounds of Beans (P. vulgaris). J. Agric. Food Chem., 23: 13.
- Larson, H. (1967): Biochemical aspects of extreme halophilism. Advan. Microb. Physiol. 1, 97-132.
- Lauer, M.J.; Blevins, D.G. (1990): Dry matter accumulation and phosphate distribution in soybean grown on varying levels of phosphate nutrition. Crop Physiol. Abstract., 16(4), 1196.
- Lal, B.M. and Verma, S.C. (1968): Physiology of Bengal gram seed III. Changes in the Phosphorus compounds of the seed parts during ripening of seed. J. Sci. Food Agri., 19: 113-116.
- Lal, B.M.; Rohewal, S.S.; Verma, S.C. and Ved Prakash (1963): Chemical composition of some sure strains of Bengal gram.

- Ann - Biochem. Exp. Med., 23: 543-548.
- Lee, T.S.G. (1987): Protein pattern changes in the developing seeds of soybean (P. vulgaris L.) Crop Physiol. Abst., 1316:
1880.
et al?
- Leon, G.A. (1968): Botanical fundamental of tropical crops. San Jose, Costa Rica, Institute Interamericana de Gencias Agricolas de la O. E.A. p. 324-325.
- Losada, M.A., J.M.R. Paneque and F.F. Del Gampo (1963): Mechanism of nitrate reduction in Chloroplasts, Biochem. Biophys. Res. Commun., 10: 298-303.
- Lodha, M.L.; N.D. Sharma; R.P. Johari and S.L. Mehta (1984): Nitrate reductase activity in nodules and leaves of groundnut (A. hypogaea L.) and green (V. radiata) Indian J. Expt. Biol. 22: 615-619.
- Loneragan, J.F., Snowball, K. and Simmons, W.J. (1968): Aust. J. Agric. Res., 19: 845-857.
- Martin, F.W. (1977): Winged bean P. tetragonolobus fact sheet, Mayaguez, Puerto Rico, Mayaguez Institute of Tropical Agriculture 1.
- Marre, E. and Murneek, A.E. (1953): Carbon Metabolism in tomato fruit as affected by pollination, fertilization application of growth regulator. Pt. Physiol., 28, 255-266.
- McAllister, D.F. and Krobar, O.A. (1951): Translocation of food reserves from soybean cotyledons and their influence on the development of the Plant, Pt. Physiol., 26: 525-38.
- Malewar, G.V.; Jadhav, N.S. and Budhewar, L.M. (1982): Possible

- role of Zinc in solution and other growth attributes of groundnut. J. Maharashtra Agri. Univ. 7: 241-242.
- Mangel, K. and Kirkby, B.A. (1982): Plant nutrients. In: Principles of Plant Nutrition, International Potash Institute, Bern, 1982.
- Mascarenhas, J.P. and Machlis (1964): Plant Physiol. 39: 70-77.
- Marme, D. (1983): "Encyclopaedia of Plant Physiology, New Series", Vol. 15B, pp. 599-626.
- Manam, R.M.; Teak, I.D., Powers, W.L. and Skidmore, E.C. (1977): Phyton, Rev. Int. Bot. Exp., 35: 189-194.
- Mehta, B.V. and P.D. Khatri (1962): Accumulation and movement of minerals in Pigeon pea (*C. cajan* Mills p.). J. Maharaja Sayajirao Univ. Ann. Bibliography of Pigeon pea II 109-122.
- Meyer, V. and E. Schulze (1984): Über die Einwirkung von Hydroxylaminsalzen auf Pflanzen Ber. dt. chem. Ges. 17: 1554-1558.
- Miflin, B.J. and P.J. Lea (1975): (F.P.J. lea and B.J.M. flin (1977) - The assimilation of ammonium nitrogen by Chlorophyllus tissue In 'Nitrogen Assimilation of Plants' Press, New York, pp. 475-499.
- Millard, A.M.; Simon and Stern (1971): Legume synthesis in developing cotyledons of V. faba. Pt. Physiol., 148-425.
- Morilla, C.A., Boyer, J.S. and Hageman, R.H. (1973): Plant Physiol. 51: 817-824.
- Muflin, B.J. and Lea, P.J. (1974): Alternative route for N assimilation in higher plants. Phytochemistry, 15: 873-885.
- Murneek, A.E. (1926): Physiology of reproduction in host plant.

- Reproduction and Metabolism Efficiency in the tomato Mo.
Agr. Exp. Sta. Res. Bull., 90: 1-19.
- Murthy, G.S. and D.J. Bagyaraj (1978): Rhizosphere Microflora of C.cajan in relation to fusarium wilt resistance. Plant Soil., 5012.
- Muflin, B.J. and P.J. Lea (1976): The pathway of nitrogen assimilation in plants. Phytochemistry, 15: 873-885.
- Muntz et al. (1976), Muntz, 1973 ab - Biochem. Physiol., 164: 357-360, 370-382. (Table 4)
- Murray, B.J.K. (1990): The regulation of soybean pod maturation Crop Physiol. Abst. 16(2): 405.
- Munjal, S.N.; Kadam, S.S. and Salunkhe, D.K. (1983): Indian J. Plant Physiol. 26(1): 21-26. (Table 9)
- Murphy, M.J.; L.M. Siegel; S.R. Tove and H. Kamin: Siroheme (1974): New prosthetic group participating in six electron reduction reactions, Cotyledons by both sulfite and nitrite reductases. Proc. Nath. Acad. Sci. U.S.A., 71: 612-616.
- NAS (1975): The winged bean, a high protein crop for the tropics. National Academy of Science, Washington, D.C.
- Nelsch, A.C. (1939): Studies on chloroplast II. Their chemical composition and distribution of certain metabolites between the chloroplasts and the remainder of the leaf. Biochem. J., 33: 300-308.
- Nelson, N. (1944): A photometric adoption of the Somogyi method for the determination of glucose. J. Biol. Chem., 153: 375-380.

- Nightingale, G.T. (1937): Nitrate metabolism, Plant Biochemistry, Ed. Bonner J. and J.E. Varnen Academic Press (Inc.) (Lond) pp. 633-672.
- Norman, M.J., P.G.E. searte, N. Dankittipakul; K.C. Ingram and J.De.B. Balsoro (1980): Evaluation of pigeon pea (*C.cajan*) as an outmn forge for coastal New South Wales. Australis. Aust. J. Exp. Agric. Anim. Husb. 20(102): 55-62.
- Oaks, A.; Wallace, W. and Stevens, D. (1972): Plant Physiol. 50: 649-654.
- Oaks, A.; Aslam, M. and Boeset, I. (1977): Plant Physiol., 59: 391-394.
- Oches, J.J. (1931): Vegetables of the Dutch East India cited by Pathak, G.N. Pulse Crops of India (1970, Chapter 2: Red gram, p. 45.
- Osborn, T.B. (1924): The vegetable protein, Longmans Green, New York.
- ✓ Paul, E. Staswick (1989): Preferential loss of an abundant storage protein from soybean pods during seed development. Pt. Physiol. Jr. 90(4): 1252-1255.
- ✓ Patnaik, S. (1955): Role of micronutrients on the growth and metabolism of rice plant Bull. Nat. Inst. Sci. India, 8: 65-68.
- ✓ Pandey, R.K.; Saxena, M.C. and Singh, V.B. (1978a): Assimilation and utilization of carbon-14s assimilates in Chickpea. J. Nucl. Agric. Biol., 7: 56-60.
- Pandey, R.K.; Singh, V.B. and Singh, B.K. (1980): Effect of reduced

sunlight on growth and yield of Chickpea Indian J. Agric. Sci., 50: 405-411.

Pandey, R.K.; Saxena, M.C., Kalubarma, M.H., Singh, V.B.S. and Prasad, V.S.S (1976): Genotypic variation in photosynthetic rate and respiratory losses in some grain legumes. Plant Biochem. J., 3: 72-80.

Palfi, G. and Jubasz, J. (1969): Zeitrchrift for Pflanzenernabeung and Bodenkunda 124: 36-47.

Parkar, J. (1958): In "Water deficit and plant growth" (T.T. Kozlowski ed.), Vol. I, pp. 195-235. Academic Press, New York.

Patil 1964): Physiological Studies in Parthenium hysterophorus: A Ph.D. Thesis submitted to the Shivaji University, Kolhapur (India).

Pate, J.S., P.J. Sharkey and C.A. Atkins (1977): Nutrition of a developing legume fruit. Functional economy in terms of carbon, nitrogen, water - Plant Physiol. 59: 506-510.

Pathak, G.N. (1970): In Pulse crops of India, 2 Red Gram, Publ. by Pruthi, T.S. Under Secretary for ICAR, New Delhi, pp. 14-53.

Peterburgski, A.V.; Chechetkina, V. (1986): Light and mineral nutrition of P. vulgaris. Crop Physiol. Abst., 12(8): 2740.

Peat, S.; Whelan, W.J.; Pirt, S.J. (1949): The amylolytic enzymes of soybean. Nature, 164: 499-500.

Pfluger, R. and Wiedemann (1977): Proc. 13th of Colloq. Int. Potash Instt., Bern., pp. 95-100.

- ✓ Pinegina, R.I. and V.G. Klimenko (1961): Variability of the content of proteins, carbohydrates and fats in seeds and vegetative mass of certain types of peas. Chemical Abst., 55: 6053-8948.
- ✓ Pickering (1979): Chronological history of plants. Little, Brown and Company, Boston.
- ✓ Pospisil, S.K.; Karlikar and E. Boamah-Mensah (1971): Investigations on winged bean in Ghana. World Crops, 23: 260-264.
- Pokhriyal, T.C.; Raturi (1986): Indian J. 111(12): 1071-1076.
- ✓ Prell, W. (1977): Z. Pflanzenzuechte, 79: 224-237.
- ✓ Raccake (1957b): Protein synthesis in ripening pea seeds. 1. Analysis of whole seeds, 2. Development of embryos and seed coats. 3. Study of pods. Biochemistry, 66: 101-116.
- Rao, K.V.; Madhava and Rajeshwar Rao (1979): N₂ and Protein metabolism of developing and germinating seeds of Pigeon pea (C. Indicus speng). Biol. Abstr., 67(5): 30877.
- ✓ Rasumov, V. (1931): The localization of photoperiodic stimulation. Genet. Plant Breed., 27: 249-282.
- ✓ Rauf, A. (1980): The free amino acid in the developing seed parts and pod of certain legumes. Acta Bot. India, 8: 196-200.
- ✓ Rajendran, R.A., Satyanarayana, V., Selvaraj, and B.S. Bhargava (1978): Some investigations on the indigenous and exotic collections of winged bean in India. In 1st International symposium on the development of potentials of the winged bean, 72-83.

- Rauf, A. and Banerji, D., 1978; Saxena and Sheldrak (1980c) and Singh (1981b): Comparative growth and biochemical studies on seed development (2) Biochemical changes in developing seed parts of certain legumes. Plant Bioch. 5: 167-173.
- Rauf, A. 1978: A comparative growth and biochemical studies on seed development 1 - biochemical changes in developing seed and pods of certain legumes. Plant Biochem. J., 5: 157-166.
- Rao, P.U. and Deosthale, Y.G. (1988): Polyphenol oxidase activity in germinated legume seeds. C.A.B. International, Chickpeas and Pigeon peas, 1(2): 158.
- Rathi, U.P.S.; Anandi Bhatt and U.S. Singh (1987): Biochemical changes in pigeon pea (*C.cajan* L. Mills. P.) leaves in relation to resistance against sterility mosaic disease Biol. Abst. AB., 84 (3): 29040.
- Reiss, H.D. and Herth, W. (1979): Planta: 145: 225-232.
- Rensing, L. and Cornelius, G. (1980): Biol. Rundsch. 18: 197-209.
- Reiss, H.D. and Herth, W. (1979): Planta, 145: 225-232.
- Reddy, N.R.; C.V. Balkrishnan and D.K. Salunkhe (1978) Phytate Phosphorus and mineral changes during germination and cooking of black gram (*P. mungo*) seeds. J. Food Sci. 43: 540.
- Reid, P.H. and F.R. Cox, 1973: Soil properties, mineral nutrition and fertilizer. In peanut culture and uses. Published by American peanut Research and Education Association, Inc. p. 271-288.

- Reid, P.H. and E.T. York Jr. (1958): Effect of nutrient deficiencies on growth and fruiting characteristics of peanuts in sand cultures Agron.J., 50: 63-67. *et al 8*
- Reddy, S.C.S. and S.V. Patil (1981): Effect of growth retardants on the yield and yield attributes of groundnuts (A. hypogea L.) Mysore Journal of Agriculture Sciences, 15(2): 238-241.
- Renfrew (1966): A report on recent findings of carbonized cereal grains and seeds from prehistoric Thessaly, Thessalike 5-21-36.
- Rijven, A.H.G.C. (1958): Effect of some inorganic nitrogenous substance on growth and nitrogen assimilation of young embryo in vitro. Aust. J. Biol. Sci. 11: 142-154.
- est*
Robson, A.D. (1988): The mineral nutrition of Lupins. Crop Physiol. Abst., 14(7): 2173.
- Rochat, Boutin (1990): Carbohydrates and nitrogenous compound changes in the hull and in the seed during the pod development of pea. Crop Physiol. Abst., 16(4): 1135.
- Rochland, L.B.; Zaragosa, E.M. and Oracca Tettch, R. (1979): Quick cooking winged bean (P. tetragonolobus). J. Food. Sci., 44: 1004.
- Russell, R.S. and D.A. Barber A. (1960): Annu. Rev. Plant Physiol., 11: 127-140.
- Saxena, N.P.; Narayanan, A. and Sheldrake, (1980) and A.R. Singh (1981a): Effect of seed grading on yield of chickpea and pigeon pea. Indian Agri. Sci. 51: 99-102.

- Sangwan, R.S.; Poplis, S. and Singh, R.C. (1983): Carbohydrate oxidising enzymes and rate of oxy. consumption in developing chickpea (*C. arietinum*) seeds. Indian J. Exp. Biol., 21: 37-39.
- Sathe, S.K.; S.S. Deshpande and Salunkhe, D.K. (1984): Dry beans of *Phaseolus*. Food. Sci. Nutrition, 21: 41.
- Salunkhe, D.K. (1975): Induction and regulation of some enzymes involved in the assimilation of inorganic nitrogen in higher plants. Biol. Chemistry, 40(1): 15-35.
- ✓ Sacher, J.A. (1973): 'Senescence and Post-harvest Physiology'. In Ann. Rev. Plant Physiol., 24: 197-224.
- Savita, Gurmit Singh (1988): Assessing the physiology; Col. maturity of pigeon pea (*C. cajan*) seeds. Pt. Physiol. CAB. Prompts series, chickpea and P. pea., 1(4): 432.
- Schorkert (1977): Z. Pflanzen Physiology, 85: 451-461.
- ✓ Schlesier, G. (1976, 1977): Nitrate reductase activity in leaves and fruits of various legumes. Biochem. Physiol. Pflanzen., 171: 511-523.
- Ryan, C.J. (1973): Annu. Rev. Plant. Physiol., 24: 173-196.
- ✓ Sajjan, V.S. and Wankhede, D.B. (1981): Carbohydrate composition of winged bean (*P. tetragonolobus*). J. Food Science, 46: 601.
- Salonen, M.L. and L.K. Simola (1977): Physiology Plant 41: 55-58.
- Sashidhar, V.R.; Mekhri, A.A. and K.S. Shastry (1977): Indian J. Agric. Sci., 47: 595-598.

- Sangappa, H.K.; T.M.M. Ali and E.S. Balraju, A.: Efficiency of insecticides in controlling the pod borers of red gram (C. cajan sprengel) Mysore J. of Agri. Sci. 11(3): 390-400.
- Savita Singh, Gurmit Singh and Harising (1987): Assessing the physiological maturity of pigeon pea (C. cajan) seeds. CAB propts series. Chickpea and pigeon pea, 1(4): 437.
- Salunkhe, D.K.; Jadhav, S.J.; Kadam, S.S.; Chavan, J.K. (1982): ✓ Chemical, biochemical and biological significance of polyphenols in cereals and legumes. Crit. Rev. Food Sci. Nutri. 17: 277.
- Schlesier, G. and K. Muntz (1974): ✓ The function of pod at protein storage in the seeds of V. faba L. III. Nitrate reductase in developing pods and seeds of Leguminosae. Biochem. Physiol. Aflanzen. 166: 87-93.
- Schweinfurth, F. (1966): ✓ The origin of cultivated plants. Harvard University Press, Cambridge, Mass.
- Sestak, Z.J.; Cotsky and P.G. Jarvis (1971): Plant photosynthetic production manual and methods. Dr. W. Junk, N.V. Publ. The Hague Netherland, pp. 343-384.
- Sestak, Z. (1966): ✓ Limitations for finding a linear relationship between chlorophyll content and photosynthetic activity. Biol. Plant Acad. Sci. Biochemistry, 8(5): 336-346.
- Sengupta, U.K. (1990): ✓ Changes of phenolic compounds during seed development and germination of groundnut seed cultivars. Crop physiology Abst., 16(5) 1337.

- ✓ Sekine, T.; Sasakawa, T.; Morita; Kimara, T.; and Karatomi, K. (1965): Laboratory manual for physiol. studies for rice. Eds. S. Yoshida; D. Forne; J.N. Cook and K.A. Gome, Pb. International Rice Research Inst. Manila.
- Sen, D. and R.W. Weaver (1980): Plant Sci. Lett., 18: 315-318.
- Sen, D. and R.W. Weaver (1981): J. Expt. Bot., 32 (129): 713-716.
- Sheldrake, A.R. and A. Narayan (1979): Growth, Development and Nutrient uptake in pigeon pea (*C. cajan* L.) J. Agri. Sci. 92(3): 513-528.
- ✓ Shrivastava (1989): Starches and protein content and enzyme level in developing cotyledons of chickpea CAB prompt series, chickpea, pigeon pea, 2: 150.
- ✓ Sharma, and R.C. Pant (1978): Myoinositol hexaphosphate as a potential inhibitor of α -amylase phytochemistry, 17: 201.
- Shrivastava; Mehta, S.L., Naik, M.S. and Eggum, B.O.C. (1981): Protein accumulation and protein quality of Bengal gram cotyledons during development. J. Agric. Food. Chem. 29: 24-27.
- ✓ Shantakumari, P. Sinha (1972) and Sinha, S.K. (1974): Variation in chlorophylls and photosynthetic rates in cultivars of Bengal gram (*C. arietinum* L.). Photosynthetica, 6: 189-194.
- Sheriff, D.W.; Fisher, M.J.; Rusitzka, G. and Ford, C.W. (1986): Aust. J. Plant Physiol. 13(3): 431-445.
- Singh, K.M. and R.S. Singh (1980): Response of pigeon pea to fertilizers and Rhizobium inoculation under rain-fed condition. Legume Research, 1(2): 87-91.

- Singh, B.K. and Pandey, R.K. (1980): Production and distribution of assimilate in chickpea (C. arietinum L.) Aust. J. Plant Physiol., 7: 727-735.
- Sinha, S.K. and D.J.D. Nicolas (1981): Nitrate reductase in 'Physiology and Biochemistry of Drought Resistance in Plants: Academic Press Australia. pp. 145-169.
- Sindhu, R.K. and H.V. Desai (1980b): Purification and characterization of diamine oxidase from groundnut embryos. Indian J. Biochemistry and Biophy., 17(3): 194-197.
- Sims, A.P.; B.F. Folks and A.H. Busey (1968); C.F. Darid and A.P. Sims (1979): Glutamine synthetase and the control of nitrogen assimilation in Lemna minor L. Academic Press, New York, pp. 501-519.
- Sindhu, R.K. and H.V. Desai (1980a): Inhibitor of diamine oxidase in cotyledons from groundnut embryos. Indian J. Biochemistry and Biophysics, 17(3): 194-197.
- Singh, V.; Raju, S.M. and Jambunathan, R. (1981c): Studies on desi and kabuli chickpea (C. arietinum L.) cultivars II. Seed protein fractions and amino acid composition. J. Food Sci. technol. 18: 86-88.
- Singh, T.P.; Soni, G.L.; Singh, H.R. (1977): Biochemical evaluation of common pulses of Punjab State. (1) Proximal analysis. J. Res. Punjab. Agric. Univ. 14: 318-322.
- Singh, U.; Jambunathan, R.G. (1981b): Studies on Desi and Kabuli chickpeas (C. arietinum L.) cultivars. Levels of protease

- inhibitor, levels of polyphenolic compounds and in vitro protein digestibility, J. Food Sci., 1364/
- ✓ Sinha, M.K. and T. Prasad (1978): The deterioration of sugar and organic and amino acids of arhar (*C. cajan*) seeds by *Aspergillus flavus*. Biol. Abstr. 66.
- Singh, V. and R. Jambunathan (1982): Qual. Plant Food. Human Nature, 31: 347.
- Sinha, S.K. and Rajgopal, V. (1965): Nitrogen assimilation and crop productivity, pp. 158-163.
- ✓ Sinha, S.K. (1973): Photosynthetica, 14(2): 8189-192. (Title of the paper)
- Slatyer, R.O. (1967) In "Plant Water Relations". Academic Press, New York.
- Smith, P.F. Reuther, W. and Specht, A.W. (1952): Proc. Am. Soc. Hortic. Sci., 59: 31-35.
- Smith, G.S., K.R. Middleton and A.S. Edmonds (1980): Na nutrition of pasture plants 2. Effect of NaCl on growth, chemical composition as the reduction of nitrate nitrogen New Phytol., 84(4): 613-622.
- Smith, D.L. (1973): Nucleic acid, protein and starch synthesis in developing cotyledons of *P. arvense* L. Ann. Bot. N.S., 37: 795-804.
- ✓ Smith, J.E. (1962): Mitochondrial transamination in sunflower cotyledons. Biochem. Biophys. Acta. 57: 183-185.
- ✓ Smith (1974): Nucleic acid, Protein and starch synthesis in developing cotyledons of Pisum arvense. L. Ann. Botany, 37: 795-804.

- Smart, J. (1976): Tropical pulses, Longman Group Limited 348.
- Spanwick and William (1964): Dunlop and Bowling (1971): Ansari and Bowling (1972).
- Spencer, Donald; Thomas, J.V.; Higgins, Susane, C. Button and R.A. Davey (1981): Protein synthesis in developing pea seeds. Bioch. Abstr., 71(5): 3555-33936.
- Srivastava, K.N.; Mehta, S.C. and Naik, N.S. (1978): RNA polymerase and protein synthesising activity in developing Bengal gram cotyledons. Indian J. Biochem. Biophys. 15: 269-272.
- Steward, C.R.; Boggers, S.F.; Aspinall, D. and L.G. Paleg (1977): Plant Physiology, 69: 930-932.
- Steward, F.C. and L.O., Chang (1963): J. Exp. Bot. 14: 379-386.
- Streeter, J.G. and M.E. Boster (1972): Comparison of in vivo and in vivo assays for nitrate reductase in Soybean leaves. Plant Physiology, 49: 448-450.
- Stewart, C.M. (1966): CSIRO Div. for Prod. Tech. Pap. 43.
- Storey, R.; L. Beevers (1977): Proteolytic activity in relation to senescence and cotyledonary development in P. sativum L. Planta, 137, 37-44.
- Stoy (1956): Photosynthesis, respiration and carbohydrate and mutation in spring wheat in relation to yield. Physiol. Plant. Suppl., 4: 1-125.
- Suzuki, Takeo (1984): Total nitrogen and free aminoacids in M. alba stems from autumn through spring. Physiol. Plant., 60(4): 473-478.

Suelter, C.H. (1970): Science 168: 789-795.

✓ Sun, Samuel (1978): Protein synthesis and accumulation in bean cotyledons during growth. Biol. Abst., 66(8): 37249-49664.

✓ Swamy, P.M.; Prakash, T.R.; Ramslakan, T. (1990): The control of proline accumulation in senescing kaf disease of cowpea (*V.unguiculata*) Crop. Physiol. Abst. Vol. 16(3): 793.

✓ Tempest, D.W.; J.L. Meers and C.M. Brown (1970): c.f. P.J. Lea and B.J. Miflin (1977): The assimilation of ammonia nitrogen by chlorophylls tissue in nitrogen assimilation of plants. Academic Press, New York, pp. 475-499.

Thimann (1970): Protein synthesis during water uptake by tuber tissue. Plant Physiology. 32: 324.

✓ Thibodene, P.S.; Jaworskiel (1975): The nutrients needed to support seed protein synthesis come from either new assimilation during seed development or redistribution from arhar tissues.

✓ Thomas, S.H. and J.L. Stoddort (1977): Leaf senescence, a review article. Ann. Rev. Plant Physiol., 31: 83-111.

✓ Toth, Prince A.L., Wallace A. and D.S. Michealson (1948): Rapid qualitative determination of 8 mineral elements in plant tissues by systematic producer involving use of a flame photometer. Soils Science, 66: 456-466.

✓ Turner (1982): Physiology of pea fruit IV. Changes in sugar in the developing seed. Aust. J. Biolog. Sci. 10, 407-413.

? Vavilos (1951): Cobely (1956); Barrua (1965): The origin, variation, immunity and breeding of cultivated plants: Transl. K.S.

- Chester Chorn. Bot. 13, 1-366.
- Vallee, J.C., Martin, C.V. and P.J. Gerard (1974): C.R. Held. Acad. Sci. Ser. D. 279.
- Vallee, J.C.; Wansuyt, G. and J. Prevost (1977): 485-488, Physical Plant, 40, 269-274.
- Van, D.W. (1990): Photosynthesis and nitrogen supply during the period of seed filling in legumes in comparison with cereals (Literature study) Crop Physical Abst., 16(4): 1170.
- Wallace, A. (1990): Plant responses to some hardly known trace elements and trace element composition and distribution in plants. Crop Physiol. Abst., 16(1): 145.
- West African Journal, Biol. Appl. (1967): Chemical Studies on some Nigerian pukes, Low pea (V. unguiculata) groundnut, Lima bean (Ph. Lunatus), Pigeon pea (C. cajan) and soybean. West Afri. Jour. Biol. Appl., 9 (52-55).
- Webb, J.R.; Obrogge, A.J. and Barber, S.A. (1954): The effect of magnesium upon the growth and the phosphorus content of soybean. Plant Proc. Soil Sci. Soc. Am., 18: 458-462.
- Whyte, R.O. and H.C. Trumble (1953): Legumes in agriculture, food and agricultural organization. Publ. Unleged Nations, Rome Italy.
- Williams, J.H. (1979): The Physiology of groundnuts (Arachis hypogaea) (V. Egret). Nitrogen accumulation and distribution. Rhod. J. Agric. Res., 17: 49-55.
- Winter, H.C.; G.K. Powell and E.E. Dekkar (1981)-4. Methylen glutamine in peanut plants, dynamics of formation, levels

- and turnover in relation to other free amino acids. Plant Physiol., 68: 588-593.
- Williams, J.H.; M. Dutta; P.T.C. Narobiar and V.M. Ramral (1987) Nitrogen fixation by groundnut In Abstracts National Seminar, 'Physiology and Biochemistry of Oil Seed Plants', organized by Deptt. of Botany, S.V. University, Tirupati (India), pp. 2-3.
- Wiewior Kowski, M.; J. Augustyniak; H. Skruzypinska; J. Przyloylska and Z. Kowki (1960): Protein Synthesis, Lupin Seeds. Chemical Abst., 54: 18695.
- Wilson, L.E. and R.A. Kennedy (1978): Photosynthetic carbon metabolism during leaf ontogeny in Zea mays L. Enzyme Studies. Plantalyz, 3: 269-274.
- Wong, K.C. (1975): The potential for four angled bean Psopocarpus tetragonolobus (D.C.) in Malaysia to increase food supply. In Proceeding of UMAGA/FAUM conference on Malaysian Food self-sufficiency, 21-23 Aug. Petaling Jaya.
- Wyn-Jones, R.G. Brandy, C.J. and Speirs, J. (1979): Academic Press, London, pp. 63-103.
- Wyn-Jones, R.G. Brandy, and Lunt, O.R. (1967): Bot. Rev. 22: 407-426.
- Yadav, J.S.P. (1981): Reclamation and crop production in Alkali soils. Current Sci. 50(a): 387-393.
- Yoshida, D.A., Cock, J.H. (1972): Laboratory manual for Physiological studies of rice. IHRI, Phillipines, Ind. J. Pt. Physiol.

30(1): 199-201.

Ziena, H.M., Schuta, A.M.; E. Tabey; Youssef (1987): Chemical Changes during the development of faba beans (*Vicia faba L.*) as affected by pod and seed position. Chem. Abstr. 110(7): 50494-59572.

Zosoldos, F. and Karvaly, B. (1978): Agron J. 74: 99-105.