

LIST OF FIGURES

Fig. No.	Contents
1	Light microscopic structure of gills of control bivalve, <i>Lamellidens marginalis</i> (X200)
2	Light microscopic structure of gills of control bivalve, <i>Lamellidens marginalis</i> (X600)
3	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 10 days (X200)
4	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 10 days (X600)
5	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 20 days (X200)
6	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 20 days (X600)
7	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 10 days (X200)
8	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 10 days (X600)
9	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 20 days (X200)
10	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 20 days (X600)
11	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to clay idol for 15 days (X200)
12	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to clay idol for 15 days (X600)
13	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to PoP idol for 15 days (X200)
14	Light microscopic structure of gills of bivalve, <i>Lamellidens marginalis</i> exposed to PoP idol for 15 days (X600)

15	Light microscopic structure of hepatopancreas of control bivalve, <i>Lamellidens marginalis</i> (X200)
16	Light microscopic structure of hepatopancreas of control bivalve, <i>Lamellidens marginalis</i> (X600)
17	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 10 days (X200)
18	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 10 days (X600)
19	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 20 days (X200)
20	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 400 ppm colour pigments for 20 days (X600)
21	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 10 days (X200)
22	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 10 days (X600)
23	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 20 days (X200)
24	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to 900 ppm colour pigments for 20 days (X600)
25	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to clay idol for 15 days (X200)
26	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to clay idol for 15 days (X600)
27	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to PoP idol for 15 days (X200)
28	Light microscopic structure of hepatopancreas of bivalve, <i>Lamellidens marginalis</i> exposed to PoP idol for 15 days (X600)