

List of Figures

Fig. No.	Title	Page
1.	<u>Parthenium hysterophorus</u> L. growing in natural habitat ..	3
2.	Phyllody disease of <u>P. hysterophorus</u> showing broom like appearance due to transformation of reproductive parts into leafy structure. ..	10
3.	Distribution of <u>P. hysterophorus</u> in world ..	15
4.	Distribution of <u>P. hysterophorus</u> in India ..	16
5.	A field showing healthy and MLO infected plants of <u>P. hysterophorus</u> ..	56
6.	A twig of healthy and MLO infected plant of <u>P. hysterophorus</u> ..	58
7.	Ray floret (♀) showing attachment of two lateral disc florets (♂) in <u>P. hysterophorus</u> . ..	59
8.	Microphotograph showing transformation of female flower and male flower into leaf like structure due to phyllody disease in <u>P. hysterophorus</u> . ..	60

Fig.No.	Title	Page
9.	Phyllody of <u>P. hysterophorus</u> showing conversion of inflorescence into a small seedling like structure ..	61
10.	Histochemical localization of MLO using Diene's stain in <u>P. hysterophorus</u> ..	66
11.	Phyllody of <u>Parthenium</u> showing +ve test for starch content ..	76
12.	Polyphenol oxidase activity in healthy and MLO infected plant leaves of <u>P. hysterophorus</u> ..	82
13.	Chromatogram showing separation of phenolic compounds of healthy and MLO infected <u>P. hysterophorus</u> ..	90
14.	Chromatography of standard phenolic compounds. ..	91-92
15.	Tracing of individual standard phenolic compounds separated on chromatographic paper ..	93
16.	Chromatographic separation of free amino acids from healthy and MLO infected <u>P. hysterophorus</u> ..	97
17.	Chromatography of standard amino acids ..	98-99

Fig.No.	Title	Page
18.	Tracing of individual standard free amino acids separated on chromatographic paper ..	100
19.	Microphotograph showing purified flakes of parthenin isolated from <u>P. hysterophorus</u> . ..	105
20.	Microphotograph showing radiating type crystals of parthenin ..	108
21.	Microphotograph showing trichome of <u>P. hysterophorus</u> present all over the plant body ..	108
22.	Absorption curve showing max of SL (parthenin) isolated from healthy and MLO infected <u>P. hysterophorus</u> ..	111
23.	Effect of different concentrations of SL isolated from healthy and MLO infected <u>P. hysterophorus</u> on germination per centage of wheat ..	113
24.	Effect of different concentrations of SL isolated from healthy and MLO infected <u>P. hysterophorus</u> on root length of wheat after 72 h germination ..	114
25.	Effect of different concentrations of SL isolated from healthy and MLO infected <u>P. hysterophorus</u> on shoot length of wheat after 72 h germination ..	115

Fig.No.	Title	Page
26.	Photograph showing effect of different concentration of SL isolated from healthy (a) and MLO infected (b) <u>P. hysterophorus</u> on wheat germination after 72 h	.. 116
27.	Photograph showing inhibitory effect of different concentration of SL isolated from healthy (a) and MLO infected (b) <u>P. hysterophorus</u> on root and shoot length of wheat after 72 h germination	.. 117
28.	Proposed pathogenetic mechanism of allergic contact dermatitis evoked by sesquiterpene lactone	.. 121
29.	NMR spectra showing structure and chemical formula of parthenin isolated from healthy <u>P. hysterophorus</u>	.. 124
30.	NMR spectra showing structure and chemical formula of parthenin isolated from MLO infected <u>P. hysterophorus</u>	.. 125
31.	HPLC analysis of SL isolated from healthy <u>P. hysterophorus</u> and visualised chromatogram of parthenin	.. 127

Fig.No.	Title	Page
32.	HPLC analysis of SL isolated from MLO infected <u>P. hysterophorus</u> and visualized chromatogram of parthenin.	.. 128