

-----  
**CHAPTER - VII**  
-----

**SUMMARY AND CONCLUSION**

## SUMMARY AND CONCLUSION

The dissertation includes the results of forestic studies on the fossil plant from the Kota formation in Chandrapur district of Maharashtra. These beds have been earlier studied by several workers. These results are given on pp. 1 in Introduction. These locality selected for the present work belongs to Kota deposits in Chandrapur district. They yielded impression as well as petrified coniferous woods.

The method employed in the investigation of the material are given in Chapter Two - Material and Methods on pp.

Chapter Three is Previous Work. It gives an account of contribution of several workers on different divisions of Upper Gondwanas in India. It gives on pp.

The geology of the area studied has been briefly given in Chapter Four Geology and Topography of Area. It is given on pp.

Chapter Five is Descriptive. It includes two parts. First part comparies descriptive of coniferous woods. It consists of two species of Podocarpoxyton; two species of Taxaceoxyton, One species of Araucarioxyton and one species of Planoxyton. Report of Planoxyton from Kota bed is a distinct feature of this work.

Part two describes the impression of Cycadophytes, Conifers, Pteridosperms and Pteridophytes. It is given on pp.

Chapter Six is General Consideration. It brings out the fact arising from the present data and are gives on pp.

It also includes the comparative account the age assigned to the beds by different workers as well as our own findings.

Chapter Seven the Summary and Conclusion. It gives the content of the different chapters mention above and gives on pp.

Chapter Eight is the references. It includes the literature cited in the present work. It is given on pp.

-----  
CHAPTER - VIII  
-----

REFERENCE

REFERENCES

- Bailey, I.W. (1933). The cambium and its derivative tissues VII. Problems in identifying the woods of Mesozoic Coniferae. Ann. Bot., Lond., 47(185) : 145-157.
- Bakshi, S.K. (1967). A new occurrence of Ginkgoites feistmantelli Bose and Sukh-Dev (1958) from the Coastal Gondwana of South India Curr.Sci. 36 : 580.
- \_\_\_\_\_ (1968). Fossil plants from Raghavapuram, Mudstone, West Gondwana Dist : A.P. India, The Palaeobotanist 16(3): 206-215.
- Banerji, J. (1982). Phlebopteris minutifollus sp.nov. from the Bhuj formation of Kachchh, India The Palaeobotanist 30 (3) : 310-315.
- Barnard, P.D.W. (1968). A new species of Masculostrobus Seward producing classopollis pollen from Jurassic of Iran. J. Linn. Soc. (Bot.) 61: (384) : 167-176.
- Bhardwaj, D.C. (1952). On a new species of Taxoxylon Unger from the Jurassic Rajmahal Hills, Bihar, India Lloydia, 15(4) : 234-240.
- \_\_\_\_\_ (1953). Jurassic woods from the Rajmahal Hills, Bihar The Palaeobotanist, 2 : 59-70.
- Biradar, N.V. (1967). Studies in fossil plants from Kota Maleri beds and embryology of genus Phoenix Linn. A Ph.D. Thesis submitted to University of Poona.

- \_\_\_\_\_ (1978). Studies in Monocotyledons : Micromegasporogenesis and Development of Female Gametophyte in Yucca gloriosa superba L. Biology vol. 4(1) : 33-43.
- \_\_\_\_\_ and Mahabale, T.S. (1978). Occurrence of Ginkgo In East Coast Gondwanas of India, 'Recent researches in Geology' 5
- \_\_\_\_\_ (1978). Occurrence of Ginkgo in East Coast Gondwana of India 'Recent researches in Geology' 5.
- Bose M.N. (1958). Morrisia a new genus of Cycadophytic fronds from the Rajmahal Hills, Bihar, India. The Palaeobotanist 7 (1) : 21-25.
- Bose M.N. (1958-a). The occurrence of cycadpteris zigno in the Mesozoic rocks of India, Palaeobotanist 6 (1-2): 113-114 (1957).
- \_\_\_\_\_ (1958 b). Cycadpteris sp. from the Rajmahal Hills, Bihar. Curr. Sci., 27 : 312.
- \_\_\_\_\_ (1960). The fossil flora of Jabalpur series-2. Filicales The Palaeobotanist (7) : 90-92.
- \_\_\_\_\_ (1966). A revision of Rajmahalia. The Palaeobotanist 14 : 85-88.
- \_\_\_\_\_ (1966). A revision of Rajmahalia The Palaeobotanist, 14 : 85-88.
- \_\_\_\_\_ (1966a). Significance of fossil plants in the Indian Gondwana stratigraphy Sci. and Cult. 32: 532-534.
- \_\_\_\_\_ (1966b). Fossil plant remains from the Rajmahal and Jabalpur series in the Upper Gondwana of India Symp. florist.strat. Gond.Land. B.S.I.P. Lucknow : 143-154.

- \_\_\_\_\_ (1974). Bennettiales. Aspects and Appraisal of Indian Palaeobotany Birbal Sahni Institute of Palaeobotany : Lucknow pp. : 189-200.
- \_\_\_\_\_ (1974). Bennetiales pp 189-200 in K.R. Surange et.al. (eds.) Aspect and Appraisal of Indian Palaeobotany, B.S.I.P. Lucknow.
- \_\_\_\_\_ and Banerji, J. (1981). Cycadophytic leaves from Jurassic Lower Cretaceous rocks of India. The Palaeobotanist, 28-29 : 218-300.
- \_\_\_\_\_ and Banerji, J. (1984). The fossil rocks of Kachchh-I Mesozoic Megafossils. The Palaeobotanist, 33 : 1-189.
- \_\_\_\_\_ and Jain, K.P. (1964). Cycadolepis sapotra from the Rajmahal Hills, Bihar, India. The Palaeobotanist, 12 (3) : 224-225
- \_\_\_\_\_ and Jain, K.P. (1967). Otozamites Vemavaramensis sp.nov. from the Upper Gondwana of the East coast of India. The Palaeobotanist, 15 (3) : 314-315.
- \_\_\_\_\_ and Kasat, M.L.(1972). The genus Ptilophyllum in India. The Palaeobotanist 19 (2) : 115-145.
- \_\_\_\_\_ Kutty, T.S. and Maheshwari H.K. (1982). Plant fossils from the Gangapur Formation. Palaeobotanist 30 : 121-142.
- \_\_\_\_\_ and Maheshwari, H.K.(1974). Mesozoic conifers In Aspects and Appraisal of Indian Palaeobotany, B.S.I.K., Lucknow, p.p. 212-223.
- \_\_\_\_\_ and Roy, S.K. (1964). Studies the Upper Gondwana of Kutch-2 Isoetaceae. The Palaeobotanist, 12 (3) : 226-228 (1963)

- \_\_\_\_\_ and Roy, S.K. (1968). On the occurrence of Pachypteris in the Jabalpur series of India. The Palaeobotanist, 16 (1) : 1-9.
- \_\_\_\_\_ and Sukh. Dev. (1958). Studies on the fossil flora the Jabalpur series from the South Rewa. Gondwana cycadpteris, Nipaniophyllum and Ginkgoites. The Palaeobotanist 7 (2) : 143-154.
- \_\_\_\_\_ and Sukh-Dev.(1972) Three new species of Pagiophyllum from Bansa, (M.P.) India. Geophytology (2) : 122-161.
- \_\_\_\_\_ and Sah, S.C.D. (1968). Some pteridophytic remains from the Rajmahal Hills, Bihar. The Palaeobotanist.
- \_\_\_\_\_ & Zeba bano (1978). Genus - Dictyozamites, oldham from India. The Palaeobotanist, 25 : 79-99 (1976).
- Cotter G. dep. (1917). A revised classification of the Gondwana system. Re.geol.surv. India. 48 : 23-33.
- Datta. et. al. 1978. Discovery of Early Jurassic micromammals from Upper Gondwana sequence of Pranhita-Godavari Valley, India, J geol. soc. India 19 : 4-68.
- Datta, P.M., Yadgiri,P. and Rao, B.R.J. (1978). Discovery of Early Jurassic micromammals from Upper Gondwana sequence of Pranhita-Godavari valley, India. J.Geol.Soc. India. 19 (2) : 64-68.
- Feistmantel, Ottókar (1876). Fossil flora of the Gondwana system. Jurassic (Oolitic) flora of the Kutch, Mem.Geol.Surv. India. Palaeont. India Ser. 11 2(1) : 1-60.



Feistmantel, O. (1876). Fossil flora of the Gondwana system.

Jurassic (Oolitic) flora of the Kutch Mem.Geol.Surv. Indi. Palaeont. Indica, Ser. 11 2 (1) : 1-60.

\_\_\_\_\_ (1877). Notes on the fossil flora of India XI. Notes on plant fossils from Barakar district (Barakar group) Ibid. 10 (2) : 73-74.

\_\_\_\_\_ (1877a). Jurassic (Liassic) Flora of Rajmahal group in the Rajmahal Hills, Mem.Geol.Surv.Ind.Pol. Ind. Ser-2, 1 (2) : 53-162.

\_\_\_\_\_ (1877b). Jurassic (Liassic) flora of Rajmahal group from Gollapalli (Near Ellore), South Godavari District II Ibid 1 (3) : 163-190.

\_\_\_\_\_ (1979). The Fossil flora of the Gondwana system; Lower Gondwana. The flora of the Talchir, Karharbari beds. Mem.Geol.Surv.India. Palaeont Indica ser. 12, 3 : 1-48. suppl. 1881 : 49-64.

\_\_\_\_\_ (1882). The fossil flora of the Gondwana system-II. Fossil flora of the south Rewa Gondwana Basin. Mem. Geol.Surv.India.Pol.Ind.ser. XII 4 (1) : 1-66.

Fox C.S. (1931). The Gondwana system and the related formations Mem.geol.Surv.India. 58 : 1-241.

Ganju, P.N. (1946). On a collection of Jurassic Plants from the Rajmahal Hills, Bihar. Jour.Indian.Bot.Soc. (Iyengar. Comm. Volume.) : 51-85.

\_\_\_\_\_ (1947a). On a collection of Jurassic plants from Rajmahal Hills, Bihar. J.Indian bot. Soc. Iyengar Commemoration volume : 51-85.

- Ganesan, T.M. and Bose, M.N. (1932). Plant remains of Mesozoic age from Lingshi Basin, Bhutan. Geophytology 12: 279-286.
- Gopal, V., J. Jacob, C. and Jacob, K. (1957). Stratigraphy and Palaeontology of the Upper Gondwana of the Ramnad District on the East Coast. Rec. Geol. Surv. India, 84(4): 477-496.
- Gothan, W. (1905). Anatomic Leber und Fossiler Gymnospermenholzer Abh. K. Preuss. Geo. Landesant. (N.F.), 44 : 108.
- Govindan A. (1975). Jurassic freshwater Ostracods from the Kota Limestone of India. Palaeontology, 18(1): 207-216.
- Gupta, K.M. (1955). Notes on some Jurassic plants from the Rajmahal Hills, Bihar, India Palaeobotanist 3 : 18-25.
- \_\_\_\_\_ (1955b). Hausmannia indica sp. nov. Gupta a Dipteridaceous leaf from the Jurassic of Rajmahal Hills, Bihar (India) Proc. Nat. Inst. Sci. India 21(3) : 147-148.
- \_\_\_\_\_ (1966). Significance of the study of the Cycadean fronds from the Upper Gondwana of India (Rajmahal Hills). In: Symposium on floristics and stratigraphy of Gondwana land pp. 137-142, Birbal Sahni Institute of Palaeobotany, Lucknow.
- \_\_\_\_\_ and Sharma, B.D. (1968 b). Investigations on the Jurassic flora of the Rajmahal Hills, India 2. On a new species of Ptillophyllum, P. sahnii from Amarjola in Amaropara region J. Palaeont. Soc. India 11:1-7.

- Gururaja, M.N. and Pant, S.C. (1970). A notes on the fossil plants from Rajmahal Hills, Bihar, Indian Minor, 24 (4): 336-388.
- Holden, R. (1913). Some fossil plants from Eastern Canada Annals of Bot. Vol. XXVII Page 243.
- Hughes, T.W.H. (1884). Southern Coalfields of Rewa Gondwana basin Ibid 21(3) : 72 (Munchen.Bot.Ztg.)
- Jain, S.L. (1959). Fossil fishes from the Kota formation of India. Proc.geol.Soc.Lond. 156 : 20-27.
- \_\_\_\_\_ (1973). New specimens of Lower Jurassic holostean fishes from India. Palacontology 16(1) : 149-177.
- \_\_\_\_\_ (1983). A review of the genus Lepidotes (Actinopterygii seminotofermes) with special reference to the species from Kota formation. (Lower Jurassic), India. Palacont. Soc.India, 28 : 7-42.
- Jain, K.P. (1968). Some plant remains from the Upper Gondwana of East Coast, India The Palaeobotanist, 16(2) : 151-155.
- Jeya Singh, D.E.P. and Sudharsan, C. (1989). Gymnospermic remains from the Sivaganga beds of the East-Coast Gondwanas, India. Proc.Spl.Geo.Con. Poona, 61-68.
- King, W. (1881). The geology of the Pranhita-Godavari valley Mem.geol.Surv.India 18(3) : 1-150.
- Krausel, R. and Jain, K.P. (1964). New fossil coniferous woods from the Rajmahal Hills, Bihar, India, The Palaeobotanist 12(1) : 59-67.

- Krishnan (1960). Geology of India.
- Kutty, T.S. (1969). Some contributions to the stratigraphy of the Upper Gondwana formations of the Pranhita-Godavari valley, Central India, The geol.soc.India, 10 (1): 33-40.
- Lele, K.M. (1955). Plant fossils from Parsora in the South Rewa, Gondwana basin, India, The Palaeobotanist 4:23-24.
- \_\_\_\_\_ (1962). Studies in the Middle Gondwana flora-I. On Dicroidium from the South Rewa Gondwana Basin, India, Ibid 10 (1-2) : 42-68.
- \_\_\_\_\_ (1964). The problem of Middle Gondwana of India Proc. 22nd Int.geol.Congr.India 9: 181-202.
- Lepekhina, V.G. (1972-1974). Woods of Palaeozoic Pycnoxylic Gymnosperms. with special reference to Noth Eurasia representative Palaeontographica, 133 B. 44-106.
- Mahabale, T.S. (1967). Mesozoic flora of India, The Kota-Maleri stage Ibid. 15 (3) : 308-313.
- \_\_\_\_\_ and Rao, S.V. (1973). Fossil flora of Rajahmundry area Proc. symp. Deccan. Trap. Country I.N.S.A.Bull. No.45 : 192-214.
- \_\_\_\_\_ and Satyanarayana, T. (1977). Two species of the Genus Taxaceoxylon Krausel and Jain from Deccan Intertrappean beds near Rajahmundry in Andhra Pradesh, India Biovigvanam 3(1) : 131-139.
- \_\_\_\_\_ (1979). Upper Gondwana plant fossils from East Godavari district Andhra Pradesh, India. Geophytology 9(1) : 65-82.

- Maheshwari, H.K. (1965). Studies in the Glossopteris flora of India - 22. On some species of the genus Glossopteris from the Raniganj stage of the Raniganj Coal field, Bengal. The Palaeobotanist 13 : 144-147.
- \_\_\_\_\_ (1966b). Studies in the Glossopteris flora of India-30. Remarks on the age of the Lower Gondwana beds of Bensoli valley, santhalparagahas, Bihar, India Syn.florist. strat. Gend. land B.S.I.P., Lucknow 110-120.
- \_\_\_\_\_ (1972). Permian wood from Antaretica and revision of some Lower Gondwana wood taxa, Palacontographica, 138B (1-4) : 1-43.
- \_\_\_\_\_ (1986). Thinnfeldia indica feistmantel and associated plant fossil from Tiruchirapalli Dist : Tamil Nadu, India The Palaeobotanist, 35(1) : 13-21.
- Medlicott and Blandford W.T. (1879-1887) A manual of Geology of India, Chiefly compiled from the observation of the Geological survey 1: 1-144 and 2: 445-817.
- Neeru Pandya, V.B. Srivastava and Sukh-Dev (1990). A new conifers fossil from Vemavaram (Early Cretaceous) Andhra India. Geophytology 20(1) : 74.
- Oldham R.D. (1893). A manual of Geology of India : 1-543.
- \_\_\_\_\_ and Morris, J. (1863). Fossil flora of the Gondwana system Bihar. In fossil flora in the Rajmahal series in the Rajmahal Hills Palacont.Ind.Ser. 1(1): 1-52.
- Pal, P.K. and Ghosh, A.L. (1987). Paralycopodium A new name for fossil shoots resembling modern Lycopodium L. earlier placed under Lycopodites Brongniart. Seventh Ind.Geophyto. Conf. Lucknow, (Abst.).

- Pant, D.D.(1955). On two new disaccate spores from the Bacchus Marsh Tillite, Victoria (Australia) Ann.Mag.nat.Mist. 12 (8) : 757-764.
- Prasad, M.N.V., (1978). Contribution to the knowledge of Fossil plants from Kamthi Beds of Maharashtra, India, Thesis, Lucknow University, 202 pp.
- \_\_\_\_\_ and Chandra (1978). First record of Toxopitys from the Indian Lower Gondwana Current Science, 47(10): 348.
- \_\_\_\_\_ (1978). Australoxylon from the Kamthi beds of Lower Gondwana, India, Curr.Sci., 47(16) : 597.
- Patra, B.P. (1973). Notes on the occurrence of Otozamites sp. in the Athgarh sand at Naraj District, Cuttack, Orissa, India Curr.Sci. 42(13) : 477-478.
- Rajanikanth, A. and Sukh-Dev (1989). The Kota formation : fossil flora and stratigraphy Geophytology 19(1) : 52-64.
- Rao, C.N. (1959). On the occurrence of Dicroidium (Thinnfeldia) feistmantelli in the east coast Gondwanas of Vemavaram, Andhra Pradesh, India. Proc.46th Indian Sc. Congr. 3 : 278.
- \_\_\_\_\_ and Shah S.C. (1959). Fossil insects from the Gondwana of India Indian Min. 12(1) : 3-5.
- \_\_\_\_\_ (1963). On the occurrence of Pterosaur from the Kota-Maleri beds, Chandrapur District, Maharashtra, Rec. geol.surv. India 92(2): 315-318.
- Rao, A.R. (1943). Jurassic spores and Sporangia from the Rajmahal Hills, Bihar Proc.Nath.Acad.Sci.India 13: 181-197.

- \_\_\_\_\_ (1964). Stachyotaxus sampathkumarani sp.nov., from Onthea In the Rajmahal Hills, Bihar, The Palaeobotanist 12 : 217-219.
- \_\_\_\_\_ and Bose M.N. (1971). Podostrobus gen. nov. a petrified podocarpaceae male cone from the Rajmahal Hills, India The Palaeobotanist 19 : 83-85
- Roy, S.K. (1965). Studies on the Upper Gondwana of Kutch-3 Otozamites Imbricatus, Feistmantel, The Palaeobotanist 13 (2) : 215-217.
- \_\_\_\_\_ (1967). Ptilophyllum horridum sp.nov. from Tambu, Kutch, India Curri.Sci. 36, (21): 581-582.
- \_\_\_\_\_ (1968). Pteridophytic remains from Kutch and Kathiwar India The Palaeobotanist, 16 (2) : 108-114.
- Sahni, B. (1928). Revision of Indian fossil plants pt. I Coniferales (Impressions and Incrustations) Palacont. Indica (nser) 11 : 1-49.
- \_\_\_\_\_ (1931). Revision of Indian fossil plants pt. II Coniferales (Petrifaction) Mem.Geol.Surv.Ind.Palacont.Indica (n.ser) 11: 53-124.
- \_\_\_\_\_ (1932). A petrified Willamsonia (W. sewardina sp.nov.) from the Rajmahal Hills, India Mem.Geol.Surv.India, Palacont. Indica (N.S.) 20 (3): 1-19.
- \_\_\_\_\_ (1936) The occurrence of Matonidium and Welchsilla in India, Rec. Geol.Surv. India 71(2): 156-165.
- \_\_\_\_\_ (1948). The Pentxyleae : A new group of Jurassic Gymnosperms from the Rajmahal Hills of India Bot.Gaz, 110 (1) : 47-80.

- \_\_\_\_\_ and Rao, A.R. (1933). On some Jurassic plants from the Rajmahal Hills, Bihar, Proc.Asiat.Soc.Beng. (N.S.) : 27 (2): 183-208.
- \_\_\_\_\_ (1934). Rajmahalia pradoxa gen.et.sp. nov. and other Jurassic plants from the Rajmahal Hills, Bihar Proc. Indian Acad.Sci. 1(6) : 258-269.
- Sah, S.C.D. (1965). Ginkgoites rajmahalense sp.nov. from the Rajmahal Hills, Bihar, India, Ibid 13(2): 155-157.
- \_\_\_\_\_ and Sukh-Dev (1957) Thinnfeldia chunakhalensis sp.nov. from Jurassic of Rajmahal Hills. The Palaeobotanist 6(i) : 22-24.
- \_\_\_\_\_ and Jain, K.P. (1964) some fossil woods from the Jurassic of Rajmahal Hills, Bihar, India, The Palaeobotanist 12(2): 169-180.
- Saksena, S.D. (1952). Correlation of the Gondwana based on the evidence of the plant fossils. Agra uni.Jour.Res. (Sc), 1 : 1-13.
- Satsangi, P.P. and Shah, S.C. (1973). A fish from Kota formation. Prannita-Godavari Basin India. (Abst). Proc. 60th Indian Sci. Cong. 3 : 193.
- Seward, A.C. (1917). Fossil plants, 3 i-XVII + 1-656. Cambridge (Uni.Press) and Sahni, B. (1920). Indian Gondwana plants A revision Palaeont Indica. 7(1) : 1-41.
- Seward, A.C.(1919). Fossil plants. Cambridge, 4 : 165-244.
- Seward, A.C. and Sahni, B.(1920). Indian Gondwana plants A revision Mem.Geol.Surv.Ind., Pal.Ind. N.S. 7 : 1-54.



Sharma, B.D. Petrified conifers with anomalous Secondary growth from the Rajmahal Hills, India, Geophytology 19 (1) : 49-51.

\_\_\_\_\_ (1971). On a Collection of Bennettitalean stems and fructifications from Amarjola in Rajmahal Hills, India Paleontograpica, 135. B : 48-52.

\_\_\_\_\_ (1989). Petrified conifers with Anomalous secondary growth from the Rajmahal Hills, India. Volume 19,(1) Geophytology.

Shah, S.C.D. and Jain, K.P. (1965). Ginkgoites rajmahalense sp.nov. from Rajmahal Hills, Bihar. The Palaeobotanist, 13 (2) : 155-157.

Singh. et.al. (1990). Equisetites sehorensis sp.nov. from Jabapur formation. Madhya Pradesh. Geophytology, 20(1) : 72-73.

Srivastava, B.P.(1945). Slicified plant remains from the Rajmahal series India. Proc.Natn.Acad.Sci.Ind., 15 (6) : 185-211.

Srivastava, R.N. and Shah, S.C. (1966). Ginkgo (Ginkgoites) digitata Brongn, from the Rajmahal Hills, santhal parganas. (Bihar), Rec.Geol.Surv. India, 94(2) : 305-312.

Stopes, M.C. (1916). An early type of the Abietineae (?) from the cretaceous of New Zealand. Ann.Bot., 30 (117) : 117-124.

Sukh-Dev.(1974). Mesozoic pteridophytic Aspects and Appraisal of Indian palaeobotanic Lucknow, pp. 73-76.

- Sukh-Dev and Rajanikanth, A. (1988). The Gangapur formation, fossil flora and stratigraphy. Geophytology 18 (1) : 1-27.
- Surange, K.P. (1966). Distribution of Glossopteris flora in Lower Symp.florist.Strat.Gend.Land. B.S.I.P. Lucknow 55-68.
- Vagyanl, B.A. (1984). Occurrence of Ginkgo crasipes (Feistmante) seaward from the Jurassic of Andhara Pradesh, India. Curr.Sci. 54 (19) : 705-706.
- \_\_\_\_\_ (1984). On the occurrence of Desmiphyllum indicum. Sahni from Vemavaram. (A.P.) Proc.5th Geophyto Conf. Lucknow : 362.
- \_\_\_\_\_ (1986). On the Occurrence of Pterophyllum footteanum, feistmantel, from Uppugunduru (A.P.) India. Bot. Report. 5 (2) : 212-213.
- \_\_\_\_\_ and Mahabale, T.S. (1972). A new specie of Fossil Gymnospermous wood. Plan.Oxylon, stopes from Adhari (M.S.) Palaeobotanist, 21 (2) : 211-215.
- \_\_\_\_\_ and Zutting, M.P. (1986). Occurrence of the Pterophyllum distans, Morris from Uppugundurau, Andhra Pradesh Geophytology. 16 (1) : 133.
- \_\_\_\_\_ and Jamane, M.R. (1987). On the occurrence of Elatocladus plana (feistmantel) Seward from Uppugunduru, Andhra Pradesh. India Curv. Sci. 56 (19) : 1023-1024.
- \_\_\_\_\_ and Mane, S.K. (1989). Pterophyllum incisum from Uppugunduru, Andhra Pradesh. Current Science, 58 (1) : 33.

- Vishnu-Mittre (1953). A male flower of Pentoxyleae with remarks on the structure of the female cones of the group. The Palaeobotanist, 2 : 75-84.
- \_\_\_\_\_ (1956). Masculostrobus Sahnii sp.nov. A petrified conifer male cone from the Jurassic of Rajmahal Hills, Bihar. Grana.Polynol 1 : 99-107.
- Vrendenburg E.W. (1910). Summary of Geology of India Calcutta.
- Wadia, D.N. (1953). Geology of India, London.
- \_\_\_\_\_ (1957). Geology of India, Macmillan and Co., London, 3rd Edn.
- Yadagiri, P. Satsangi, P.P. and Prasad, K.N. (1980). The Piscean fauna from the Kota formation of the Pranhita Godavari valley, Andhra Pradesh. Mem.Geol.Surv.India.Palaeont. Indica, 45 : 1-131.
- Zeba bano and Bose M.N. (1981). Motonidium cinglatum n.sp. from Kutch, India. The Palaeobotanist 27 (1) : 95-99.
- Zellier, R. (1902). Observations sur quelques plantes fossiles des Lower Gondwana. Mem.Geol.Surv.India. Palaeont Indica. N.S. : 2(1) : 1-40.