

## LIST OF FIGURES

Sr. No.	Title	Page No.
1.	Length of Shoot	
	a) Initial growth phase	34
	b) Growth rate after establishment	40
2.	Number of leaves per plant	
	a) Initial growth rate	35
	b) Growth rate after establishment	41
3.	Girth of stem	
	a) Initial growth rate	36
	b) Growth rate after establishment	42
4.	A phenogram records of phenological events of <u>Ipomoea carnea</u> spp. <u>fistulosa</u>	46
5.	Eury species exhibit awide tolerance zone	61
6.	T.S. of stem of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from arid habitat	62
7.	T.S. of root of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from arid habitat	62
8.	T.S. of leaf of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from arid habitat	62
9.	T.S. of stem of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from aquatic habitat	63
10.	T.S. of root of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from aquatic habitat	63
11.	T.S. of leaf of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> from aquatic habitat	63
12.	Allelopathic effects of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> on wheat	64
13.	Allelopathic effects of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> on Jowar	66
14.	Allelopathic effects of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> on Rice	67
15.	Allelopathic effects of <u>Ipomoea carnea</u> spp. <u>fistulosa</u> on <u>Phaseolus aconitifolius</u>	69