

BIBLIOGRAPHY

Nutraceutical evaluation of some wild edible tuberous plants.

- Aberoumand A. (2009). Preliminary assessment of nutritional value of plant based diets in relation to human nutrients. *Int. J. Food Sci. Nutr.*, **1**: 1-8.
- Aberoumand, A. and Deokule, S. S. (2008). Comparison of compounds of some edible plants of Iran and India. *Pak. J. Nutr.*, **7** (4): 582-585.
- Aberoumand, A. and Deokule, S. S. (2009). Proximate and mineral composition of wild Coco (*Eulophia ochreata* L.) tubers in Iran. *As. J. Food Agri. Industry*, **2** (2): 203-209.
- Aberoumand, A. and Deokule, S.S. (2008). Determination of elements profile of some wild edible plants. *Food Anal Methods.*, Doi.10.1007/s 12161-008-9038-z.
- Aberoumand, A. and Deokule, S.S. (2009). Studies on nutritional values of some wild edible plants from Iran and India. *Pak. J. Nutr.*, **8**(1): 26-31.
- Adeyeye, E. I. and Otokiti, M.K. (1999). Proximate composition and some nutritional valuable minerals of two varieties of *Capsicum annum*. *Discovery and Innovation*, **11**: 75-81.
- Agrahar-Murugkar, D. and Subbulakshmi, G. (2005). Nutritive value of wild edible fruits, berries, nuts, roots and consumed by the khasi tribes of India. *Eco. Food Nutr.*, **44**: 207-223.
- Alessandra, G. and Robert, H.C. (2005). The crucial role of metal ions in neuro Degeneration, the basis for promising therapeutic strategy. *Brit J. of Pharmaco*, **146**: 1041 - 1059.
- Alozie, Y., Akpanabiatus, M.I., Eyong, E.U., Umoh, I.B. and Alozie, G. (2009). Amino acid composition of *Dioscorea dumetorum* Varieties. *Pak.J.Nutr.*, **8** (2): 103-105.

Nutraceutical evaluation of some wild edible tuberous plants.

Annan, K., Kojo, A.L., Cindy, A., Samuel, A.N. and Tunkumgnen, B.M. (2010).

Profile of heavy metals in some medicinal plants from Ghana commonly used as components of herbal formulations. *Pharmacognosy Research*, 2(1): 41-44.

Ansari, M.Y. (1968). A new species of *Ceropegia* Linn. (Asclepiadaceae) from Western ghats, Maharashtra. *Bull. Bot. Surv. India*, 40: 95-97.

Ansari, M.Y. (1969). *Ceropegia media* (Huber) Ansari Stat. Nov .from Western ghats, Maharashtra. *Bull. Bot. Surv. India*, 11: 199-201.

Ansari, M.Y. (1984). Fasicles of flora of India. *Bull. Bot. Surv. India*, 16: 1-16.

Ansari, M.Y. and Kulkarni, B.G. (1980). A new species of *Ceropegia* Linn. (Asclepiadaceae) from the Western ghats in Maharashtra state, India. *Bull. Bot. Surv. India*, 22: 1-4.

Ansari, M.Y. (1971). *Ceropegia vincaeifolia* Hook. (Asclepiadaceae) from Maharashtra its history and identity. *Bull. Bot. Surv. India*, 13:187-191.

Ansari, M.Y.(1972). New *Ceropegia* Linn. (Asclepiadaceae) from sahyadri range in Maharashtra state. *J. Bomb. Nat. Hist. Soc.*, 69: 250-253.

AOAC, (1990). Official Methods of Analysis. *Association of Official Analytical Chemists*, Washington DC.

Arafat, S.M., Gaafar, A.M., Basuny, A.N. And Nassef, S .L. (2009). Chufa tubers (*Cyperus esculantus* L.): As a new source of food. *World Applied Sci. J.*, 7 (2):151-156.

Arekal, G.D. and Ramkrishna, T.M. (1981). External nectaries of *Calotropies gigantia* and *Waittakaka volubilis*. *Phytomorphology*, 36: 303-306.

Arntzen, J. C. and Ritter, M. E. (1984). Encyclopedia of Agricultural Science. Orlando,Academic Press.

Nutraceutical evaluation of some wild edible tuberous plants.

- Atawodi, S.E. (2005). Antioxidant potential of African medicinal plants. *Afr. J. Biotechnol.*, **4**: 128-133.
- Auti, B.K. and Jagdale, P.E. (2008). Medico-botanical studies on some angiosperms from Eastern part of Mula Dam Ahmednagar District, Maharashtra State. *Ad. Plant Sci.*, **21**(II): 603-605.
- Balentine,, D. A., Aibano, M. C. and Muraleedharan, G. N. (1999). Role of medicinal plants, herbs and spices in protecting human health. *Nutr. Reviews*, **57**: 41-45.
- Barminas, J.T., Charles, M. and Emmanuel, D. (1998). Mineral composition of non conventional leafy vegetables. *Plant Food Hum.Nutr.*, **53**: 29-36.
- Barua, U., Hore, D.K. and Sarma, R. (2007). Wild edible plants of Majuli Island and Darrang district of Assam. *Int. J. Trad. Knowledge*, **6**(1): 191-194.
- Bergmeyer, H.U. (1974). Methods of enzymatic analysis (Vol.2) (Publ.). Academic Press, Inc., New York, San Francisco, London: 673-684.
- Bhandari, M.R., Kesai, T. And Kawabata, J. (2003). Nutritional evaluation of wild yam (*Dioscorea spp.*) tubers of Nepal. *Food Chem.*, **82**: 619-623.
- Blatter, E. and McCann, C. (1933). Revision of the flora of the Bombay presidency (Asclepiad). *Ibid*, **36**: 524-537.
- Borah, S., Baruah, A. M., Das, A. K. and Borah, J. (2009). Determination of mineral content in commonly consumed leafy vegetables. *Food Anal. Methods*, **2**: 226-230.
- Brody, T. (1998). Nutritional Biochemistry San Diego Academic press: 11- 12.
- Brown, K. H. and Wuehler, S. E. (2000). Zinc and human health: results of human trials and implications for program interventions and research. *Micronutrient Initiative.*, Canada, International Research Centre.

Nutraceutical evaluation of some wild edible tuberous plants.

- Char, M.B.S. (1978). A new species of *Brachystelma* R.Br.(Asclepiadaceae) from India. *Curr. Sci.*, **47**(24): 965-966.
- Chatterjea, M.N. and Shinde, R. (1998). Text of medicinal biochemistry. Jaypee brothers medican publishers: 173.
- Chelikani, P., Ramana, T. and Radhakrishnan T.N. (2005). Catalase: A repertoire of unusual features. *Indian J. Clin. Biochem.* **20** (2): 131-135.
- Cheng, S. S., Chang, H. T., Vchang, S. T., Tsai, K. H. and Chen, W. J. (2003). Bioactivity of selected plant essential oils against yellow fever mosquito *Aedes aegypti* larvae. *Bioresource Tech.*, **89**: 99-102. Chanwitheesuk (2005).
- Cho, E.J., Yokozawa, T., Rhyu, D.Y., Kim, S.C., Shibahara, N., Park, J.C. (2003). Study on the inhibitory effects of Korean Medicinal plants and their main compounds on the 1, 1-diphenyl-2-picrylhydrazyl radical. *Phytomedicine* **10**: 544-551.
- Choudhary, K., Singh, M. and Pillai, U. (2008). Ethnobotanical survey of Rajasthan- An Update. *American-Eurasian J. Botany*, **1**(2): 38-45.
- Chuakul, W. and Boonpleng, A. (2004). Survey on medicinal plants in Ubon Ratchathani Province (Thailand). *J. Phytopharmacy*, **11**(1): 33-54.
- Cook, T. (1904). The flora of the presidency of Bombay.
- Dassana, M.D. and Jaysuriya, A.H.M. (1975). A new species of *Brachystelma* (Asclepiadaceae) from Srilanka. *Ceylon J. Bio.Sci.*, **11**: 39-41.
- Dey, P.M. (1990). (eds.) Methods in plant biochemistry, II. Carbohydrates (Publ.) Acad press London.

- Diaz-Gomez, N.M., Domenech, E., Barroso, F., Castells, S., Cortabarria, C. and Jimenz, A. (2003): The effect of zinc supplementation on linear growth body composition and growth factors in preterm infants. *Pediatrics*, **111** (5): 1002 - 1009.
- Dogan, S., Turan, P., Dogan, M., Arslan, O. and Alkan, M. (2007). Partial characterization of peroxidase from the leaves of thymbra plant (*Thymbra spicata* L.var. *spicata*). *Eur. Food Res. Tech.*, **225**: 865-871.
- Dogan, Y., Baslar, S., AY, G. and Mert, H.H. (2004). The use of wild edible plants in western and central Anatolia(Turkey). *Economic Botany*, **58**(4): 684-690.
- Duhan, A., Chauhan, B.M. and Punia, D. (1992). Nutritional value of some non conventional plant foods of India. *Plant food Human Nutr.*, **42**: 193-200.
- Dyer, R.A. (1977). New species of *Brachystelma* (Asclepiadaceae). *Bothalia* **12**: 55-57.
- Dyer, R.A. (1978). New records of *Brachystelma*. *J. South. Afr.*, **43**: 9-20.
- Dyer, R.A. (1979). Asclepiadaceae: Two new species of *Brachystelma*. *Bothalia*, **12**: 497-498.
- Edison, S., Unnikrishnan, M., Vimala, B., Pillai, S.V., Sheela, M.N., Sreekumari, M.T. and Abraham, K. (2006). Biodiversity of tropical tuber crops in India; NBA Scientific Bulletin 7. National Biodiversity Authority of India, Chennai, Tamilnadu, India: 60.
- Ejoh, R.A., Nkonga, D.V., Inocent, G. and Moses, M.C. (2007). Nutritional components of some non-conventional leafy vegetables consumed in Cameroon. *Pak. J. Nutr.*, **56**(6): 712-717.

Nutraceutical evaluation of some wild edible tuberous plants.

- Eruvbetine, D. (2003). Canine nutrition and health. A paper presented at the seminar organized by kensington pharmaceuticals Nig. Ltd.
- Etkin, N. L. and Ross, P. J. (1991). Should we set at place for diet in ethnopharmacology. *J. Ethnopharmacology*, **32**: 25-36.
- Etkin, N. L. and Ross, P. J. (1982). Food as medicine and medicine as food an adaptive framework for the interpretation of plant utilization among the Hansa of Northern Nigeria. *Social Medicine* **16**: 1559-1573.
- Falconer, J. (1990). The major significance of 'minor' forest products the local use and value of forests in the West African humid forest zone Community forestry note no 6.
- FAO/WHO (1973). Adhoc expert committee on energy and protein requirement. WHO Tech. Report Series (Genewa): 522.
- Fleuret, A. (1993). Dietary and therapeutic uses of fruit in three Talta communities in
- Etkin, N. L., ed., Plants in indigenous medicine and diet. Redgrave, Bedford Hills, NY.
- Folin, O. and Denis, W. (1985). A colorimetric estimation of phenols (and phenolic derivatives) in urine. *J. Biol. Chem.*, **22**: 305-308.
- Gaikwad, S.P. and Yadav, S.R. (2004). Endemic flowering plant species of Maharashtra and their possible utilization. Biodiversity in India vol.3 (Editor T.Pullaiah Regency publication): 28-58.
- Gamble, J.S. (1923): The flora of the presidency of Madras.

- Ghirardini, M.P., Carli, M., Vecchio, N.D., Rovati, A., Cova, O., Valigi, F. and Agnetti, G. (2007): The importance of a taste. A comparative study on wild food plant consumption in twenty-one local communities in Italy. *J.Ethnobiol. Ethnomed.*: 3-22.
- Ginnopolitis, C. N. and Ries, S.K. (1977). Superoxide dismutase: Occurrence in Higher Plants. *Plant physiol.*, **59**: 309-314.
- Glew, R. S., Vanderjagt, D. J. and Bosse, R. (2005). The nutrient content of three edible plants of the Republic of Niger. *J. Food Comp. Anal.*, **18**: 15–24.
- Goud, P.S. and Pullaiah, T. (1996). Ethnobotany of Kurnool district-some wild plants as Food. *J. Econ. Taxon. Bot.*, **12**: 224-227.
- Gupta, K. and Wagle, D.S. (1986). Nutritional and antinutritional factors of green leafy Vegetables. *J. Agric. Food Chem.*, **36**: 472-474.
- Hambiadge, M. (2000). Human zinc deficiency. *Tour of Nutrition Denver*, **130**: 1344-1349.
- Hammilton, E.M.N., Whitney, E.N. and Sizer, F.S. (1994). Nutritional: Concepts and controversies, St. Paul, M. N. (USA) 4th ed.: 223.
- Hanif, R., Iqbal, Z., Iqbal, M., Hanif, S. and Rashid, M. (2006). Use of vegetables as nutritional food: role in human health. *J.Agric. Biological Sci.*, **1**(1): 18-22.
- Hashmi, D.R., Ismail, S. and Shaikh, G.H. (2007). Assessment of the level of trace metals in commonly edible vegetables locally available market of Karachi city. *Pak. J. Bot.* **39**(3): 747-751.
- Hasling, C., Sondergaard, K. and Moselkilde, C.P. (1991). Calcium metabolism in

Nutraceutical evaluation of some wild edible tuberous plants.

postmenopausal osteoporotic woman is determined by dietary calcium and coffee intake. *Am. J. Nutr.*, **23**: 119 - 126.

Hassan, S. W., Umar, R.A., Ladan, M.J., Neyemilke, P., Wasagu, R. S. U., Lawal, M. and Ebbo, A.A. (2007). Nutritive value, phytochemical and antifungal properties of *Pergularia tomentosa* L. (Asclepiadaceae). *Int. J. Pharmacog.*, **3**(4): 334-340.

Hatano, T., Edmatsu, R., Mori, A., Fujita, Y., Yasuhara, E. (1989). Effects of tannins and related polyphenols on superoxide anion radical and on 1, 1-diphenyl-2-picrylhydrazyl radical. *Chem. Pharm. Bull.*, **37**: 2016-2023.

Halevy, S., Koth, H. and Guggenheim, K. (1957). The vitamin and mineral content of fruits and vegetables grown in Israel. *Nutrients in Israeli fruits and vegetables II*: 409-413.

Hawk, P.B., Oser, B.L. and Summerson, W.H. (1948). Practical physiological chemistry (Publ.). The Blockiston Co. USA.

Heaney, R.D. (1994). Thinking straight about calcium. *The New Eng. J. Med.*, **328** (7): 503 – 505.

Hedrick, U.P. (1919). Sturtevants edible plants of the world; New York, Dover Publication: 182-122.

Hemadri, K. (1968). A new *Ceropegia* Linn. (Asclepiadaceae) from Western ghats, Maharashtra. *Bull. Bot. Surv. India*, **10**: 123-125.

Hemadri, K. and Ansari, M.Y. (1971). *Ceropegia mahabale* A new species from Sahyadri range, Western ghats. *Indian forester*, **97**: 105-108.

Hemadri, K. (1968). A new *Ceropegia* Linn. (Asclepiadaceae) from Western ghats, Maharashtra. *Bull. Bot. Surv. India*, **10**: 123-125.

Hooker, J.D. (1983). The flora of British India.

Indrayan, A.K., Sharma, S., Durgapal, D., Kumar, N. and Kumar, M. (2005).

Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. *Curr. Sci.*, **89**(7): 1252-1255.

International Institute for Environment and development IIED (1995). The value of wild resources in agricultural systems. The Hidden Harvest, London.

Irawan, D., Hanny, C.H., Limin, S.H., Hashidoko, Y., Osaki, M. and Kulu, I. P. (2006). Ethnobotanical study and nutrient potency of local traditional vegetables in central Kalimantan. *Tropics* **15**(4): 441-448.

Irvine, F.R. (1952). Supplementary and Emergency food plants of West Africa. *Economic Botany*, **6**(1): 23-40.

Ishtiaq, M., Hanif, W., Khan, M.A., Ashraf, M. and Butt, A.M. (2007). An ethnomedicinal survey and documentation of important medicinal folklore food phytonyms of flora of Samahni valley, (Azad Kashmir) Pakistan. *Pak. J. Biol. Sci.*, **10** (13): 2241-2256.

Jagtap, S.D., Deokule, S.S. and Bhosale, S.V. (2008). Ethnomedicobotanical uses of endemic and RET plants utilized by the Korku tribe of Amaravati district, Maharashtra. *Ind. J. Trad. Know.*, **7**(2): 284-287.

Jagtap, S.D., Deokule, S.S., Pawar, P.K. and Harsulkar, A.M. (2009). Traditional ethnomedicinal knowledge confined to the Pawra tribe of Satpura hill, Maharashtra, India. *Ethnobotanical Leaflets*, **13**: 98-115.

Jain, A., Katewa, S.S., Galav, P. K. and Sharma, P. (2005). Medicinal plant diversity of Sitamata Wildlife sanctuary, Rajasthan, India. *J. Ethnopharmacol.*, **102**: 143-157.

Nutraceutical evaluation of some wild edible tuberous plants.

- Jain, S.K. (1963). Wild plant food of the tribals of Bastar (Madhya Pradesh). *Prod. Nat. Inst. Sci. India*, **30**(2): 56-64.
- Jaworska, G. and Kmiecik, W. (1999). Content of selected mineral compounds, nitrates and oxalates in spinach (*Spinacia oleracea L.*) and New Zealand spinach (*Tetragonia expansa*) from spring and autumn growing seasons. *Food Sci. Tech.*, **2**: 221-226.
- Johns, T and Duquette, M. (1991). Deficiency of phosphorus in man. *Am. J. clin. Nutr.*, **53**: 448 - 456.
- Kahkonen, M. P., Hopia, A. J., Vuorela, H. J., Raucha, J. P., Pithlaja, K. and Kulaja, T. S. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *J. Agri. Food Chem.*, **47**: 3954-3962.
- Kalra, E.K. (2003). Nutraceutical-Definition and Introduction. *AAPS Phar. Sci.*, **5** (2): 1-3.
- Kamble, M.Y and Yadav, S.R. (2005). Wild asclepiads of ornamental potential from Maharashtra. Plant Taxonomy: Advances and relevances (Eds. A. K. Pandey, Junwen and J. V. V. Dogra), CBS publisher and Distributors, New Delhi: 299-302.
- Kamble, M.Y. (2007): Taxonomic studies on asclepiadaceae and periplocaceae of Maharashtra. Ph.D.thesis (Shivaji University,Kolhapur)
- Kamble, S.Y., More, .N., Patil, S.R., Pawar, S.G., Bindurani, R. and Bodhankar, S.L (2008). Plants used by tribes of Northwest Maharashtra for the treatment of gastrointestinal disorders. *Ind. J. Trad. Knowledge*, **7**(2): 321-325.
- Kanodia, K.C. and Reddy, B.V. (1964). *Ceropegia fantastica* (Asclepiadaceae) an imperfectly known species. *Bull. Bot. Surv. India*, **6**: 311-312.

Nutraceutical evaluation of some wild edible tuberous plants.

- Khan, A.A., Shukla, K.M.L. and Khan, I.M. (2000). Enumeration of wild food plant of ethnobotanical significance in central India. *Ad. Plant Sci.*, **13** (1): 277-281.
- Khyade, M.S., Kolhe, S.R. and Deshmukh, B.S. (2009). Wild edible plants used by the tribes of Akole taluk of Ahmednagar district (MS), India. *Ethnobotanical Leaflets*, **13**: 1328-1336.
- King, S.R. and Gershoff, S.N. (1987). Nutritional evaluation of three underexploited Andean tubers: *Oxalis tuberosa* (Oxalidaceae), *Ullucus tuberosus* (Basellaceae) and *Tropaeolum tuberosum* (Tropaeolaceae). *Economic Botany*, **41**(4): 503-511.
- Kirk, J.O.T. and Allen, R.L. (1965). Dependence of chloroplast pigment on actidione Arch. *Biochem. Biophys. Res. Commun.*, **21**: 523-530.
- Kirtikar, K.R. and Basu, B. D. (1975). Indian medicinal plants Vol.II. Ed. Jayyed Press, New Delhi: 240.
- Kubmarawa, D., Andenyang, I.F.H. and Mogomya, A.M. (2008). Amino acid profile of two non-conventional leafy vegetables, *Sesamum indicum* and *Balanites aegyptiaca*. *Afr. J. Biotech.*, **7**(19): 3502-3504.
- Kubmarawa, D., Andenyang, I.F.H. and Mogomya, A.M. (2009). Proximate composition and amino acid profile of two non-conventional leafy vegetables, *Hibiscus cannabinus* and *Haematostaphis bartrri*. *African J. Food Sci.*, **3**(9): 233-236.
- Linder, K. and Manria, C. (1991). Nutritional biochemistry and metabolism with chemical application. *Appleton and Lange, Norwalk* **2**: 191-212.
- Lockett, C., Calvert, C.C. and Grivetti, L.E. (2000). Energy and micronutrient

Nutraceutical evaluation of some wild edible tuberous plants.

- composition of dietary and medicinal wild plants consumed during drought. Study of rural Fulani, Northeastern Nigeria. *Int. J. Food Sci. and Nutr.*, **51**: 195-208.
- Loliger, L. (1991). The use of antioxidants in food in Aruoma O.I. and Halliwell B. Editors. Free radicals and food additives, London; Taylor and Francis.
- Lowry, O.H., Rosenbrough, N.J., Furr, A.L. and Randall, R.J. (1951). Protein measurements with folin phenol reagent. *J. Biol. Chem.*, **193**: 262-263.
- Luck, H. (1974). In: Methods in Enzymatic Analysis II (ed.) Bergmeyer. (Publ.) Academic Press, New York: 885.
- Luximon-Ramma, A. and Bahorun, T. (2002). Antioxidant activities of phenolic proanthocyanidin and flavonoid components in extracts of *Cassia fistula*. *J. Agri. Food Chem.*, **50**(18): 5042-5047.
- Machly, A.C. (1954). Methods in biochemical analysis (Ed.) Glick D. (Publ.) Interscience publishers Inc. New York: 385-386.
- Mahekar, P.D. and Yadav, S.R. (2006). Medicinal plants of south western Maharashtra; Biodiversity in India Vol.4 (Edited by T.Pullaiah, Regency publication New Delhi): 75-599.
- Malhotra, V.K. (1998). Biochemistry for students 10th Edition Jaypee Brothers medical publishers (P) Ltd. New Delhi India.
- Marwah, R.G., Fatope, M. O., Mahrooqi, R. A., Varna, G. B., Abadi, H.A., Suad, K.S. and Burtamani, A.(2007). Antioxidant capacity of some wild edible and wound healing plants in oman. *Food Chem.*, **101**(2): 465-470.
- Mensah, J., Okoli, K., Ohaju-Obodo, R.I. and Eifediyyi, K. (2008). Phytochemical,

- nutritional and medicinal properties of some leafy vegetables consumed by
Edo people of Nigeria. *Afr. J. Biotech.*, **7**(14): 2304-2309.
- Mills, D.F. (1981). Symposia from the XII International congress on Nutrition. Prog.
Clin. Biol. Res. **77**: 165 - 171.
- Moon, J.K. and Shibamoto, T. (2009). Antioxidant assays for plant and food
components. *J. Agri. Food Chem.*, **57**: 1655-1666.
- Nag, A., Galav, P. and Katewa, S.S. (2007). Indigenous animal healthcare practices
from Udaipur district, Rajasthan. *Ind. J. Trad. Knowl.*, **6**(4): 583-588.
- Najafi, S. and Deokule, S.S. (2010). Pharmacognostic study of *Tylophora dalzellii*
Hook.F. *J. Medicinal plant Res.*, **4**(5): 403-406.
- Ndlovu, J. and Afolayan, A.J. (2008). Nutritional analysis of the South African wild
vegetable *Corchorus olitorius* L. *Asian J. Plant Sci.*, **7** (6): 615-618.
- Nelson, N. (1944). A Photochemical adaptation of the Somogyi method for the
determination of glucose. *J. Biol. Chem.*, **153**: 375-380.
- Nile, S.H. and Khobragade, C.N.N. (2009). Determination of nutritive value and
mineral elements of some important medicinal plant of western part of India.
J. medicinal Plant, **8**(5): 79-88.
- NRC (1989). Recommended Daily Allowance. Food and Nutition Board. 10th ed.,
National Research Council. National Academy of Science Press, Washington
DC, USA.
- Odhav, V.B., Beekrum, S., Akula, U. S. and Baijnath (2007). Prelimanery assement
of nutritional value of traditional leafy vegetables in Kwazulu Natal, *South*
Africa. J. Food Comp. Anal., **20**: 430-435.
- Odukoya, O.A., Inya-Igaha, S. I., Segum, F. I., Sofidiya, M. O. and Hori, O. O.

Nutraceutical evaluation of some wild edible tuberous plants.

- (2007). Antioxidant activity of selected Nigerian green leafy vegetables. *Amer. J. Food Tech.,* **2** (3):169-175.
- Ogbuagu, M.N. (2008). Nutritive and Antinutritive composition of the wild (In-Edible) species of *Dioscorea bulbifera* (Potato yam) and *Dioscorea dumetorum* (Bitter yam). *J. Food Tech.,* **6** (5): 224-226.
- Oguntibeju, O.O. (2008). The biochemical, physiological and therapeutical roles of ascorbic acid. *Afr. J. Biotech.,* **7** (25): 4700-4705.
- Oliveira, J.S. and carvalho, M. F. (1975). Nutritional value of some edible leaves used in Mozambique. *Economic Botany,* **29:** 255-263.
- Oyenuga, O.A. and Fetuga, B.L. (1975). Some aspects of the biochemistry and nutritive value of the watermelon seed (*Citrullus vulgaris* schrad). *J. Sci. Food Agric.,* **26:** 843-846.
- Padmaja, G., Moorthy, S. N., Nambisan, B., Babu, L., Sundaresan, S., Sajeev, M. S., Nanda, S. K., Susan, J. K., Rajalekshmi, L., Sudha, D. and Manikantan, N. M. (2005). Digestibility of Starch and Protein. In: Analytical Methodologies for Tropical Tuber Crops. Eds: Central Tuber Crops Research Institute, Kerala: 34-38.
- Pardo-de-Santayana, M., Tardio, J., Blanco, E., Carvalho, A.M., Lastra, J. J., Miguel, E. S. and Morales, R. (2007). Traditional knowledge of wild edible plants used in the northwest of the Iberian Peninsula (Spain and Portugal): a comparative study. *J. Ethnobi. Ethnomed.,* **3:** 27.
- Patil, V.N. (1990). Morphotaxonomic studies in asclepiadaceae genus: *Ceropegia* L. and *Brachystelma* R.Br.;M. phil. Dissertation, Shivaji University(Unpublished).

Nutraceutical evaluation of some wild edible tuberous plants.

- Poornima, G. N. and Ravishankar, R.V. (2009). Evaluation of phytonutrients and vitamin content in a wild yam, *Dioscorea belophylla* (Prain)Hains. *Afr. J. Biotech.*, **8** (6): 971-973.
- Powers, K.M., Oberley, L.W. and Domann, F.E. (2008). The adventure of superoxide dismutase in health and diseases: superoxide in the balance. G. Valacchi, Davis P.A.(eds.) *Oxidants in Biology*: 183-196.
- Prasad, A.S. (1982). Clinical, biochemical and nutritional aspects of trace elements, Prasad A.S. Ed., New York: Liss: 3.
- Prasad, A.S. (1993). Essential and toxic elements in human health and diseases: an update, New York: Wiley.: 46-51.
- Radhakrishnan, K., Pandurangan, A.G. and Pushpangadan, P. (1996). Ethnobotany of Wild edible plants of Kerala, India. Ethnobia. In Hum. Welfare, Deep Publications, New Delhi: 48-51.
- Raghvan, R.S. and Ansari, M.Y. (1975). Chromosome number in the genus *Ceropegia* Linn. *Curr. Sci.*, **44**: 863-864.
- Rajurkar, N.S. and Damame, M.M. (1998). Mineral content of medicinal plants used in the treatment of diseases resulting from urinary tract disorders. *Appl. Radiat. Isot.*, **49** (7): 773-776.
- Rajyalakshmi, P., Venkatalakshmi, K. Venkatalakshmamma, K., Joyshjna, Y., Balachandramani, D. K., Suneetha, K. (2001). Total carotenoid and beta carotene content of forest green leafy vegetables consumed by tribals of south India. *Plant Food Hum. Nutr.*, **56**: 225-238.

Nutraceutical evaluation of some wild edible tuberous plants.

- Ramarathnam, N., Ochi, H., Takeuchi, M. (1997). Antioxidants Defense system in Vegetable Extracts. In: Shahidi F (ed). Natural antioxidants: Chemistry, Health Effects and Applications. Aoca Press: Champaign, IL: 76-87
- Rashid, A., Anand V.K. and Serwar, J. (2008). Less known wild edible plants used by the Gujjar tribe of district Rajouri Jammu and Kashmir state –India. *Int. J. Bot.*, 4 (2): 219-224.
- Reddy, K.N., Pattanaik, C., Reddy, C.S. and Raju, V.S. (2007). Traditional knowledge on wild food plants in Andhra Pradesh. *Ind. J. Trad. Knowl.*, 6 (1): 223-229.
- Reddy, M.B., Chidambaram, M.V. and Bates, G.W. (1987). Iron transport in microbes, plants and animals, Winkelmann G., Helm D.V. and Neilands J.B. Eds., New York: 429.
- Sabnis, S.D. and Bedi, R. (1971). *Ceropegia odorata*, A little known plant of Western India. *Kew Bull.* 25: 57-60.
- Sadashivam, S. and Manikam, A. (1992). Biochemical method for agricultural sciences, Willey, Eastern Ltd.: 105.
- Sahu, T.R. (1996). Life support promising food plants among Aboriginals of Bastar(M.P.) India. Ethnobi. Hum. Welfare, Deep Publication New Delhi: 26-30.
- Samant, S.S. and Dhar, U. (1997). Diversity, endemism and economic potential of wild edible plants of Indian Himalaya. *Int. J. Sustain. Dev. World Ecol.*, 4: 179-191.
- Santapau, H. (1948). Miscellaneous notes 43. The genus *Ceropegia* further comment. *J. Bom. Nat. Hist. Soc.*, 47: 775-777.

Nutraceutical evaluation of some wild edible tuberous plants.

- Santapau, H. and Irani, N.A. (1958). The genus *Ceropegia* in Bombay. *Bull. Bot. Soc. Bengal*, **12**: 6-14.
- Santapau, H. and Irani, N.A. (1960). The asclepiadaceae and periplocaceae of Bombay.
- Scelig, M. (1989). Cardiovascular consequences of Mg deficiency and loss; pathogenesis, prevalence and manifestations. *Ame Tour of Cardio.*, **63**: 1101 – 1102.
- Schachter, M.B. (1996). The importance of magnesium to human nutrition, Schachter center for complementary medicine.
- Schoeninger, M.J., Bunn, H.T., Murry, S.S. and Marlett (2000). Compositional of tubers used by Hadza foragers of Tanzania. *J. Food Comp. Anal.*, **13**: 1-11.
- Seeling M.S. and Heggtveit, H.A. (1974). Magnesium interrelationship in ischemic heart Disease. *Amr. J. Clin. Nutr.*, **27**: 59-112.
- Sekeroglu, N., Ozkutlu, F., Deveci, M., Dede, O. and Yilmaz, N. (2006). Evaluation of some wild plants aspects of their nutritional values used as vegetable in eastern black sea region of Turkey. *Asian J. Plant Sci.* **5** (2): 185-189.
- Sekine, T., Sasakawa, T., Morita, S., Kimura, T. and Kuratom, K. (1965). A laboratory manual for physiological studies of Rice (Eds.)
- Shanthakumari, S., Mohan, V.R. and Britto, J.D. (2008). Nutritional evaluation and elimination of toxic principles in wild yam (*Dioscorea spp.*). *Tropical subtropical agroecosystem*, **8**: 319-325.
- Sheela, K., Nath, K.G., Vijayalakshami, D., Yankanchi, G. M. and Patil, R.B. (2004). Proximate composition of under utilized green leafy vegetables in southern Karnataka. *J. Hum. Ecol.*, **15** (3): 227-229.

Nutraceutical evaluation of some wild edible tuberous plants.

- Shretha, P.M. and Dhillon, S.S. (2006). Diversity and traditional knowledge concerning wild food species in locally managed forest in Nepal. *Agroforestry System*, **60**: 55-63.
- Simopoulos, A.P. (2004). Omega-3 Fatty acids and antioxidants in edible wild plants. *Biol. Res.*, **37**: 263-277.
- Singh, H.B. and Arora, R.K. (1978). Wild edible plants of India. ICAR, New Delhi.
- Smith, W.D. and Hammarsten, J.F. (1958). Serum mangnesium in clinical disorders. *South Mol. J.*, **51**: 1116-1117.
- Stajner, D., Igic, R., Popovic, B.M. and Malencic, D. (2008). Comparative study of antioxidant properties of wild growing and cultivated *Allium* species. *Phytother. Res.*, **22**: 113-117.
- Sundriyal, M. and Sundriyal, R.C. (2001). Wild edible plants of the Sikkim Himalaya: Nutritive values of selected species. *Economic botany*, **55**(3): 377-390.
- Sundriyal, M., Sundriyal, R.C. and Sharma, E. (2004). Dietary use of wild plant resources in the Sikkim Himalaya, India. *Economic Botany*, **58**(4): 626-638.
- Swaminathan, R. (2003). Magnesium metabolism and its disorders. *Clin Biochem. Rev.*, **24**(2): 47
- Swarnkar, S. and Katewa, S.S. (2008). Ethnobotanical observation on tuberous plants from tribal area of Rajasthan(India). *Ethnobotanical Leaflets*, **12**: 647-666.
- Talwar, G.P., Srivastava, L.M. and Mudgil, K.D. (1989). Textbook of biochemistry and human biology. 2nd ed. Hall of India private Ltd.
- Tayie, F. A. K. and Asibey-Berko, E. (2001). Mineral contents of some indigenous vegetables of Ghana. *Ghana J. Sci.*, **41**: 49-54.
- Thunus, L. and Lejeune, R. (1994). Handbook on metals in clinical and analytical

- chemistry, Seiler H.G., Sigel A. and Sigel H. Ed. New York: Dekker: 333-667.
- Toth, S.J., Prince, A.L., Wallace, A. and Mikkenlsen, D.S. (1948). Rapid quantitative determination of eight mineral elements in plant tissue by systematic procedure involving use of a flame photometer. *Soil Sci.*, **66**: 459-466.
- Tripathi, Y.C. (2000). Promising resources of Rajasthan forest. *Ad. Plant Sci.*, **13** (1): 31-37.
- Turan M., Kordali S., Zengin H., Dursun, A. And Sezen, Y. (2003). Macro and micro mineral content of some wild edible leaves consumed in eastern Anatolia. *Soil Plant Sci.*, **53**: 129-137.
- Ukpabi, U.J., Ukenye, E. and Olojede, A.O. (2009). Raw material potentials of Nigerian wild Polynesian arrowroot (*Tacca leontopetaloides*) Tubers and starch. *J. Food Tech.*, **7**(4): 135-138.
- Underwood, E.J. (1977). Trace elements in human and animal nutrition. 4th ed., Academic Press Inc., New York.
- Upadhye, A., Kumbhojkar, M.S. and Vartak, V.D. (1986). Observation on wild plants used in folk medicine in the rural areas of the Kolhapur district. *Ancient Sci. Life*, **VI** (2): 119-121.
- Varghese, E. (1996). Applied ethnobotany – A case study among the kharias of central India. Deep publication: 78.
- Varjravelu, E. (1987). A note on the *Brachystelma glabrum* Hook, (Asclepiadaceae) from South India. *J. Bomb. Nat. Hist.*, **84**: 262-263.

Nutraceutical evaluation of some wild edible tuberous plants.

- Venkatesh- Iyengar, G., Kawamura, H., Parr, R.M., Miah, F.K., Wang, J., Dang, H.S., Djojosubroto, H., Cho, S. Y., Akher, P., Natera, E. S. and Nguy, M.S. (2002). Dietary intake of essential minor and trace elements from asian diates. *Food Nutr.Bull.*, **23**(3): 124-128.
- Venkatreddi, B. (1968). *Ceropogia lawii* concepts; *J. Wiley and sons. Inc.*, New york.
- Victoria, R.G., Huanca, T., Vadez, V., Leonard, W. and Wilkie, D. (2006). Cultural, practical and economic value of wild plants: A quantitative study in the Bolavian Amazon. *Economic Botany*, **60**(1): 62-74.
- Vishwakarma, K. and Dubey, V. (2009). Nutritional potential of *Ipomea aquatica* FORSSK.: An ethnobotanically important plant. *J. Indian Bot. Soc.*, **88** (1&2): 159-162.
- Wadhawa, and Ansari, M.Y. (1968). A new species of *Ceropogia* Linn. (Asclepiadaceae) from Western ghats, Maharashtra. *Bull. Bot. Surv. India*, **10**: 95-97.
- Walter, W.M., Croom, E.D., Catignani, G.L. and Thresher, W.C. (1986). Compositional study of *Apios priceana* tubers. *J. Agric. Food Chem.*, **34**: 39-41.
- Wehmeyer, A.S. (1986). Edible wild plants of southern Africa. *NFRI*: 20-68.
- WHO/FAO/UNU (1985). Report: Energy and protein Requirement: WHO technical report series No.724: 220(WHO Geneva).
- Yadav, S.R., Salunkhe, C.B. and Dixit, G.B. (1989). Two new records of asclepiadaceae from Maharashtra. *J. Bombay Nat. Hist. Soc.*, **86**: 480-482.
- Yildirim, E., Dursun, A. and Turan, M. (2001). Determination of the nutrition

Nutraceutical evaluation of some wild edible tuberous plants.

- contents of the wild plants used as vegetable in upper Coruh valley. *Turk. J. Bot.*, **25**: 367-371.
- Zennie, T. M. and Ogzewalla, C. D. (1977). Ascorbic acid and vitamin A content of edible wild plants of Ohio and Kentucky. *Economic Botany*, **31**: 76-79.
- Zweig, G. and Whitkar, J.R. (1971). Paper chromatography and electrophoresis. (Publ.) Academic Press, Inc. New York.