



Bibliography

-
- Aderhold, D.; Williams, C.J. and Edyvean, R.G.J. (1996). The removal of heavy-metal ions by seaweeds and their derivatives. *Biores. Technol.* **58**: 1-6.
- Alexander, A. (1985). Optimum timing of foliar nutrient sprays. In: Alexander, A., (ed.) Foliar Fertilization. Publ Leiden, The Netherlands, Martinus-Nijhoff, pp 44-60.
- Arasaki, S. and Arasaki, T. (1983). Vegetables from the Sea. Japan Publication International, Tokyo, pp: 196.
- Armisen, R. (1995). World-wide use and importance of *Gracillaria*. *Journal of Appl. Phycol.* **7**: 231-243.
- Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts, polyphenol oxidase in *Beta vulgaris*. *Plant Physiol.*, **2**: 1-15.
- Atzmon, N. and Van Staden, J. (1994). The effect of seaweed concentrate on the growth of *Pinus pinea* seedlings. *New Forests.* **8** (3): 279-288.
- Banaimoon, S.A. (1992). Fatty acids in marine macroalgae from Southern Yemen (Hadramout) including occurrence of Eicosatetraenoic (20:4) and Eicosapentaenoic (20:5) acids. *Bot. Mar.* **33**: 165-168.
- Beckett, R.P. and Van Staden, J. (1989). The effect of seaweed concentrate on the growth and yield of potassium stressed wheat. *Plant and Soil*, **116** (3): 29-36.
- Beckett, R.P.; Mathegka, A.D.M. and Van Staden J. (1994). Effect of seaweed concentrate on yield of nutrient stressed tepary bean (*Phaseolus acutifolius* Gray). *J Appl Phycol.*, **16**: 429-430.
- Bensky, D. and Barolet, R. (1990). Chinese Herbal Medicine: Formulas and Strategies, rev. ed., Eastland Press, Seattle, WA.
- Bligh, E. G. and Dyer, W. J. A (1959). A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, **37**: 911-917.
- Blunden, G. (1977). Cytokinin activity of seaweed extracts. In: Faulkner D.L. and W.H. Fenical (ed.) Marine Natural Products Chemistry. (Publ.) Plenum, New York. pp-337-343.
- Blunden, G. (1991). Agricultural uses of seaweeds and seaweed extracts. In: Guiry, M. D. and Blunden, G. Seaweed Resources in Europe: Uses and Potential. (Publ.) John Wiley and Sons, Chichester. pp- 65-81.
-

-
- Blunden, G. and Wildgoose, P. B. (1977). The effects of aqueous seaweed extract and kinetin on potato yields. *J. Sci. Food Agri.*, **28** (2):121-125.
- Blunden, G.; and Woods, D.L. (1969). Effects of carbohydrates in seaweed fertilizers. In: *Proc. 6th Inter. Symp. on Seaweed Research*. Santiago de Compestela, Spain, pp. 647.
- Blunden, G.; Jenkins, T. and Liu, Y. (1997). Enhanced leaf chlorophyll levels in plants treated with seaweed extract. *J. Appl. Phycol.*, **8**:535-543.
- Bokil, K.K.; Mehta, V.C. and Datar, D.S. (1974). Seaweeds as manure: II pot culture manorial experiments on wheat. *Phykos.*, **13** (1): 1-5.
- Boynton, D.(1954). Nutrition of foliar application. *Annu. Rev. Plant Physiol.* **5**:31-54.
- Brain, K.R.; Chalopin, M.C.; Turner, T..D.; Blunden, G. and Wildgoose, P.B. (1973). Cytokinin activity of commercial aqueous seaweed extract. *Plant Sci. Letters*, **1**:241-245.
- Button, E.F. and Noyes, C. F. (1964). Effect of seaweed extract upon the emergence and survival of seedling of creeping red fescue. *Agron. J.* **60**: 324-326.
- Chakraborty, S. and Santra, S.C. (2008). Biochemical composition of eight benthic algae collected from Sundarban. *Ind J Marine Sci.*, **37**(3): 329-332.
- Chalker-Scott, L. (2005). Seaweed Extracts Reduce Disease, Improve Production, and Increase Stress Resistance in Landscape Plants. Available online at [www.puyallup.wsu.edu/~linda%20chalker scott/Horticultural%20Myths_files/Myths/ Seaweed%20extracts.pdf](http://www.puyallup.wsu.edu/~linda%20chalker%20scott/Horticultural%20Myths_files/Myths/Seaweed%20extracts.pdf)
- Chapman, V.J. and Chapman, D.J. (1980). Seaweeds and their uses, (3rd edition.). (Publ.) Chapman and Hall, London, pp: 334.
- Chidambaram, K. and Unny, M.M. (1953). Note on the value of seaweeds as manure. *I Intl. Seaweed Symposium.*, pp: 67-68.
- Choudhary, N.K.; Guru, R.C. and. Dashm, M.C (1976). Observation on the relationship between leaf pigment concentration, moisture and dry weight before i.e. of fall in *D. sisso*. Roxb. *Comp. Physiol. Ecol.*, **1**(4): 147-149.
- Clark, R. B. (1984). Physiological aspects of calcium, magnesium and molybdenum deficiencies in plants. *Agronomy Monography* (2nd edition) Madsion, U.S.A. pp. 99-168.
-

-
- Clarkson, D.T. and Hanson, J.B. (1980). The mineral nutrition of higher plants. *Annu Rev Plant Physiol.* **31**: 239-298.
- Crouch, I.J. and Van Staden, J. (1993). Evidence for the presence of plant growth regulators in commercial seaweed products. *Pl. Growth Regul.*, **13**(1): 21-29.
- Crouch, I.J. and Van standen, J. (1992). Effect of seaweed concentrate on the establish and yield of green house tomato plants. *J. App. Phycology*, **4**: 291-296.
- Csizinszky, A.A. (1984). Response of tomatoes to seaweed based nutrient sprays. *Proc. Fla. State Hort. Soc.*, **97**:151-157.
- Dank, M.L.; Fletcher, J.S. and Rice, E.L. (1975). Effects of phenolic inhibitors on growth and metabolism of glucose-UL-¹⁴C. in Poul's scarlet rose cell suspension cultures. *Am. J. Bot.*, **62**: 311-317.
- Dave, M.J. and Parekh, R.G. (1975). Protein content of green seaweeds from the Sourashtra coast. *Salt Res. and Ind.*, **11**(2): 41-44.
- Dave, M.J.; Parekh, R.G.; Ramavat, B.K.; Doshi, Y.A. and Chauhan, V.D. (1987). Protein content of red seaweeds from Gujarat coast. *Seaweed Res. and Util.*, **10**(1): 17-20.
- Demir, N.; Dural, B. and Yildirim, K. (2006). Effect of seaweed suspensions on seed germination of tomato, pepper and aubergine. *J. Biol. Sci.*, **6**:1130–1133.
- Demos, E.K.; Woolwine, M.; Wilson, R.H. and McMillan, C. (1975). The effects of ten phenolic compounds on hypocotyl growth and mitochondrial metabolism of mung bean. *Amer. J. Bot.*, **62**: 97-102.
- Dhargalkar, V.K. (1979). Biochemical studies on *Ulva reticulata* Forsskal. *Proc. Int. Symp Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, pp: 40.
- Dhargarkar, V.K. (1986). Biochemical studies in *Ulva reticulata* Forsskal, *Mahasagar Bull of the National Institute of Oceanography*, **19** (1):45-57.
- Durand, C.; Manuel, M.; Boudouresque, C.F.; Meinesz, A.; Verlaque, M. and Le Parco, Y. (2002). Molecular data suggest a hybrid origin for the invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta) in the Mediterranean Sea. *J. of Evol. Biol.*, **15**: 122–133.
-

- El-Baroty, G.S.; Moussa, M.Y.; Shallan, M.A.; Ali, M.A.; Sabh, A.Z. and Shalaby, E.A. (2007). Contribution to the aroma, biological activities, minerals, protein, pigments and lipid Contents of the red alga: *Asparagopsis taxiformis* (Delile) *Trevisan J. of App. Sci. Res.*, **3**(12): 1825-1834.
- El-Sheekh, M. M. and El-Saled, Ed-D. (2000). Effect of crude seaweed extract on seed germination, seedling growth and some metabolic processes of *Vicia faba* L. *Cytobios*, **101**: 23-35.
- Epstein, E. (1972). Mineral Nutrition of Plants: Principles and Perspectives. (Publ.) John Wiley. & Sons, New York, pp. 412.
- Eris, A.; Sivritepe, H.O. and Sivritepe, N. (1995). The effects of seaweed (*Ascophyllum nodosum*) extract on yield and quality criteria in peppers. *Acta Hort.*, **412**:185-192.
- Esteves, B.S.; Taogil, A. and Suzuki, M.S. (2005). Nutrient composition of microalgae and macrophytes of the Acu lagoon, *Acta Limno. Bras.*, **17**(3):233-244.
- Evans, H. J. and Sorger, G. J. (1966). Role of mineral elements with emphasis on the univalent cations. *Annu. Rev. Plant Physiol.*, **17**:47-76,
- FAO (2006). Year book of fishery statistics, vol 98 (1-2). Food and Agricultural Organisation of the United Nations, Rome.
- Featonby-Smith, B.C. and Van -Staden , J. (1983). The effect of seaweed concentrate and fertilizer on the growth of *Beta vulgaris*. *Z. Pflanzenphysiol. Bd.*, **112**:155-162.
- Feibert, E. B. G.; Shock, C. C. and Saunders, L. D. (2003). Non-conventional additives leave onion yield and quality unchanged. *Hort Science*. **38**(3):381-386.
- Fleurence, J. (2004). Seaweed proteins. In : Proteins in food processing. (eds.). Yada, R.Y. (Publ.) Woodhead, Cambridge, pp 686.
- Fleurence, J. (1999). Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends Food Sci. Technol.*, **10**: 25-28.
- Folin, O. and Denis, W. (1915). A colorimetric estimation of phenols (and phenolic derivatives) in urine. *J. Biol. Chem.*, **22**: 305 – 308.

-
- Fuller, W. H. and Roger, R. N. (1952). Utilization of the phosphorus of algal cells as measured by the Neubauer technique. *Soil Sci.*, **74**: 417-429.
- Galls, A.D.M. and Dunlap, J. (1974). Influence of phenolic acids on ion uptake. IV. Depolarization of membrane potentials. *Plant Physiol.*, **56** : 801-806.
- Ganesan, M. and Kannan, L.(1994). Seasonal variation in the biochemical composition of some green algae from Mandapam coast. *Seaweed Res. and Util.*, **12**(1-2): 37-40.
- Genard, H.; Le Saos, J.; Billard, J. P.; Tremolieres, A. and Boucaud, J. (1991). Effect of salinity on lipid composition, glycine betaine content and photosynthetic activity in chloroplasts of *Suaeda maritima*. *Plant Physiol Biochem.*, **29**: 421–427
- Goatley, J.M. and Schmidt, R.E. (1990). Seedling Kentucky bluegrass growth responses to chelated iron and biostimulator materials. *Agron. J.*, **82** (5):901-905.
- Gupta, U.C. (1979). Copper in agriculture crops. In: Copper in the environment Part-I. Ecological cycling (Eds.) J.O. Nriagu.. (Publ.) John wiley and Sons, New York. Pp. 255-288.
- Hall, J.R. (1991). Growth regulators research in Kentucky bluegrass sod production. Proc. Virginia Turfgrass Landscape Conference. **31**:69-71.
- Haroon, A.M., (2000). The biochemical composition of *Enteromorpha* spp. from the Gulf of Gdansk coast on the southern Baltic Sea. *Oceanologia*, **42**: 19-28.
- Hawk, P. B.; Oser, B. L. and Summerson, W. H. (1948). Practical physiological chemistry. (Publ.) The Blackiston Co. USA. pp-1323.
- Hawker, J. S.; Marschner, H. and Downtown, W.J.S.(1974). Effects of sodium and potassium on starch synthesis in leaves. *Aust J Plant Physiol.* **1** : 491- 501.
- Henriques, F. S. (1989). Effects of copper deficiency on the photosynthetic apparatus of sugarbeet (*Beta vulgaris* L.). *J Plant Physiol.*, **135**: 453-458.
- Hong, D.D.; Hien, H.M. and Son, P. N. (2007). Seaweeds from Vietnam used for functional food, medicine and biofertilizer. *J. Appl. Phycol.*, **19**:817–826.
-

-
- Honya, M.; Kinoshita, T.; Ishikawa, M.; Nori, H. and Nisizawa, K. (1993) Monthly determination of alginate, M/G ratio, mannitol and minerals in cultivated *Laminaria japonica*. *Bull Jap Soc Sci Fish*, **59**:295–299
- Hsiao, T.C. and Lauchli, A.(1986). Role of potassium in plant water relationship. In : Advance in plant nutrition Vol. 2 (Eds.) B. Tinkler and A. Lauchli.. (Publ.) Parger New York. pp 281- 311
- Indegaard, M. and Minsaas, J.(1991). Animal and human nutrition. In: Seaweed resources in Europe: Uses and potential. (Eds.), M. D. Guiry and G. Blunden. (Publ.) John Willey and Sons London, pp.21–64).
- Ito, K. and Hori, K. (1980). Seaweed: Chemical composition and potential uses. *Food Review Intl.*, **5**: 101-144.
- Jennings, R. C. (1968). Gibberellins as endogenous growth regulators in green and brown algae. *Planta*, **80**: 34-42.
- Jensen, A. (1969). Tocopherol content of seaweed and seaweed meal III. Influence of processing and storage on the content of tocopherols, carotenoids and ascorbic acid in seaweed meal. *J. Sci. Food Agr.*, **20**:622-626.
- Jensen, A. (1972). The nutritive value of seaweed meal for domestic animals. In: *Proc. of the Inter. Symposium on Seaweed Research*. Aug. 8-12, 1971. Sapporo, Japan.
- Jiménez-Escrig, A. and Goni, C. I. (1999). Nutritional evaluation and physiological effects of edible seaweeds. *Arch Latinoam Nutr.*, **49** : 114-120.
- Joko, S.; Lestari, O. A. and Anugrahati, N. A. (2006). Substitution of wheat flour with seaweed flour to increase dietary fiber and iodine content in dry noodle". *J. of Food Sci. and Tech.*, **4** (2): 131 - 145.
- Jones, A. L. and Harwood, J. L. (1993). Lipids and lipid metabolism in the marine alga, *Enteromorpha intestinalis*. *Phytochemistry*, **34**(4):969–972.
- Jurkovic, N.; Kolb, N. and Colic, I. (1995). Nutritive value of marine algae *Laminaria japonica* and *Undaria pinnatifida*. *Nahrung*, **1**:63-66.
- Kennish R. and Williams, G. A. (1997). Feeding preferences of the herbivorous crab *Grapsus albolineatus*: The differential influence of algal nutrient content and morphology. *Mar. Ecol. Progr. Ser.*, **147**:87–95.
-

-
- Khaleafa, A. F.; Kharboush, M. A. M.; Metwalli, A.; Mohsen, A. F. and Serwi, A. (1975). Antibiotic (fungicidal) action from extracts of some seaweeds. *Bot. Mar.*, **18**: 163 - 165.
- Kirk, J. O. T. and Allen, R. L. (1965). Dependence of chloroplast pigment on actidione *Arch. Biochem. Biophys. Res. Commun.*, **21**: 523-530.
- Kolb, N.; Vallorani, L.; Milanovi, N. and Stocchi V. (2004). Evaluation of Marine Algae Wakame (*Undaria pinnatifida*) and Kombu (*Laminaria digitata japonica*) as Food Supplements. *Food Technol. Biotechnol.*, **42** (1) 57–61.
- Krishnamurthy, V. and Joshi, H.V. (1970). A checklist of Indian marine algae. C.S.M.R.I., Bhavnagar. : 1-36.
- Kuiper, D. and Staal, M. (1987). The effects of exogenously applied plant growth substances on the physiological plasticity in *Plantago major* spp. Pleiosperma: Responses of growth shoot to root ratio and respiration. *Physiol. Plant.*, **69**:651-658.
- Kumar, V. (1993). Biochemical constituents of marine algae from Tuticorin coast. *Indian J. of Mar. Sci.*, **22**: 138-140.
- Lean, D. R. and Nalewajko, C. (1976). Phosphate exchange and organic phosphorus excretion by freshwater algae. *J. Fish. Res. Bd. Can.*, **33**: 1312-1323.
- Loliger, J. (1991). The use of antioxidants in food. In: Free radicals and food additives. (eds.) Aruoma, O. I. and Halliwell B., (Publ): Taylor London, pp. 129–150.
- Lourenço, S.; Barbarino, E.; Nascimento, A.; Freitas, J. and Diniz, G. (2006). Tissue Nitrogen and Phosphorus in seaweeds in a tropical eutrophic environment: what a long term study tells us. *J. of Appl. Phycol.*, **18**(3-5): 389-398.
- Lowry, O. H.; Rosebrough, N. J.; Farr, A L. and Randell, R. J. (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, **193** : 265-275.
- Lynn, B. (1972). The chelating properties of seaweed extract *Ascophyllum nodosum* vs. *Macrocystis periferia* on the mineral nutrition of sweet peppers, *Capsicum annuum*. M. Sc. Thesis, Clemson University, Clemson, S.C.
- Mabeau, S. and Fleurence, J.(1993.) Seaweed in food products : Biochemical and nutritional aspects. *Trends in Food Sci. Technol.*, **4**: 103-107.
-

-
- Machold, O. and Stephan, W. W. (1969). Function of iron porphyrin in chlorophyll bio-synthesis. *Phytochem.*, **8** : 2189-2192.
- Manivannan, K.; Karthikai Devi, G.; Thirumaran, G. and Anantharaman, P. (2009). Mineral composition of marine macroalgae from Mandapam coastal regions; southeast coast of India. *Am. Eur. J. of Bot.*, **2** (1): 42-51.
- Marinho-Soriano, E and Bourret, E. (2005) Polysaccharides from the red seaweed *Gracilaria dura* (Gracilariales, Rhodophyta). *Bioresource Technology*, **96**:379–382.
- Marinho-Soriano, E.; Fonseca, P.C.; Carneiro, M.A.A. and Moreira, W.S.C. (2006). Seasonal variation on the chemical composition of two tropical seaweeds. *Biores. Technol.*, **97**: 2402-2406.
- Marschner, H. (1986 and 1995). Mineral nutrition of higher plants. (Pub). Academic Press, New York.
- Mc Dermid, K. J. and Stuercke, B. (2003). Nutritional composition of edible Hawaiian seaweeds. *J. of Appl. Phycol.*, **15**:513–524.
- Mc Dermid, K.J. and Stuercke, B. (2004). A comparison of the nutritional composition of Hawaiian *Gracillaria* species. In: Taxonomy of Economic Seaweeds. (eds), Abbott , I. A., and Mc- Dermid, K.J. (Publ.) Hawaii. Sea Grant College University of Honolulu, pp-241-246.
- Mercer, J.P.; Mai, K.S. and Donlon, J. (1993). Comparative studies on the nutrition of two species of abalone, *Haliotis tuberculata-coccinea* Linnaeus and *Haliotis discushannai* :1.Effects of algal diets on growth and biochemical composition. *Invertebr. Reprod.Dev.*, **23**:75–88.
- Mohan, V.R.; Venkataraman Kumar, V.; Murugeswari, R. and Muthuswami, S. (1994). Effect of crude and commercial seaweed extracts on seed germination and seedling growth in *Cajanus cajan* L. *Phykos.*, **33** (1-2): 47–51.
- Moller, M. and Smith, M. L. (1999). The effects of priming treatments using seaweed suspensions on the water sensitivity of barley (*Hordeum vulgare* L.). *Annal Appl. Biol.*, **135** : 515 – 521.
- Mooney, P.A. and Van Staden, J. (1985). Effect of seaweed concentrate on the growth of wheat under conditions of water stress. *South African J. Sci.*, **81**:632-633.
-

-
- Morgan, K. T. and Tarjan, A.C. (1980). Management of sting nematode on centipede grass with kelp extracts. *Proc. Fl. St. Horti. Soc.* **93**: 97-99.
- Murthy, M.S. and Radia, (1978). Eco-biochemical studies on some economically important Intertidal algae from Port Okha (India). *Bot. Mar.*, **21**(7): 417-422.
- Muthuraman, B. and Ranganathan, R. (2004). Biochemical studies on some green algae of Kanyakumari coast. *Seaweed Res. and Util.*, **26**(1& 2): 69-71.
- Myers, D.J. and Perry, M.W. (1986). Organic materials applied as seed treatment or foliar sprays fail to increase grain yield of wheat. *Australian J. Exper. Agric.*, **26**: 367-373.
- Nabati, D.A.; Schmidt, R.E. and Parrish, D.J. (1994). Alleviation of salinity stress in Kentucky bluegrass by plant growth regulators and iron. *Crop Sci.*, **34** (1):198-202.
- Nada, K.; Vallorani, L.; Nada, M. and Vilberto, S. (2004). Evaluation of Marine Algae Wakame (*Undaria pinnatifida*) and Kombu (*Laminaria digitata japonica*) as food supplements. *Technol. Biotechnol.*, **42** (1) 57-61.
- Nagai, T. and Yukimoto, T. (2003). Preparation and functional properties of beverages from sea algae. *Food Chem.*, **81**:327-332
- Naik, V.N. (1998). Flora of Marathwada. (Publ.) Amrut, Aurangabad, pp646.
- Neela, M.V. (1956). Analysis of seaweeds. *Home Sci Bull Women's Christian Coll. Madras.* pp. 46.
- Nelson, W.R. and Van Staden, J. (1984). The effect of seaweed concentrate on growth of nutrient-stressed, greenhouse cucumbers [*Cucumis sativa* Ecklonia maxima]. *Hort Sci.* **19** (1): 81-82.
- Nelson, W.R. and Van Staden. J. (1986). Effect of seaweed concentrate on the growth of wheat. *South African J. Sci.*, **82** (4):199-200.
- Nisizawa, K.; Noda, H.; Kikuchi, R. and Watanabe, T.
(1987.) The main seaweed foods in Japan. *Hydrobiologia*, **151**: 5-29.
- Norrie, J. and Keathley, J.P. (2006). Benefits of *Ascophyllum nodosum* marine-plant extract applications to 'Thompson seedless' grape production. *Acta Hort.*, (ISHS) **727**: 243-248
-

-
- Osawa, T. (1994). Novel natural antioxidants for utilization in food and biological systems. In: Postharvest chemistry of plant food- materials in the tropics (eds.), Uritani, I.; Garcia, V. and Mendoza, E. M. (Publ.) Japan Scientific Societies Press, Tokyo, Japan, pp. 241–251.
- Parekh, R.G.; Maru, L.V. and Dave, M.J. (1977). Chemical composition of green seaweeds of Saurashtra Coast. *Bot. Mar.*, **20**(6): 359-362.
- Passam, H.C.; Olympios, C.M. and Akoumianakis, K.(1995). The influence of pre and postharvest application of seaweed extract on early production and storage of cucumber, *Acta Hort.*, **379**:229
- Pattama, R. and Chirapart, A. (2006). Nutritional evaluation of tropical green seaweeds *Caulerpa lentillifera* and *Ulva reticulate*. *Kasetsart J. Nat. Sci.*, **40** (1) : 75 – 83.
- Pillai, V.K. (1956). Chemical studies on Indian seaweeds: Mineral constituents. *Proceeding of Indian Academic Science*, **B45**: 3-29.
- Reed, M. (1907). The economic seaweeds of Hawaii and their food value. *Annual Report of the Hawaii Agricultural Experimental Station* , **1909**: 61-88.
- Reitz, S. R. and Trumble, J. T. (1996). Cytokinin-containing seaweed extract does not reduce damage by an insect herbivore. *Hort. Sci.*, **31**: 102-105
- Riekert*, H , (1972). *Drugs from the Sea*, Govt Printer, RSA.
- Rodrigo, R. and Bosco, C. (2006). Oxidative stress and protective effects of polyphenols: Comparative studies in human and rodent kidney. A review. *Comparative Biochem. Physiol. Part C*, **142**:317-327.
- Sadasivam, S. and Manickam, A. (1992). *Biochemical method for agricultural science*, Willey, Eastern Ltd. pp. 105.
- Sanderson, K.J. and Jameson, P.E. (1986). The cytokinins in a liquid seaweed extract: Could they be the active ingredients? *Acta Horticulturae*, **179**: 113-116.
- Sandmann , G. and Boger, P. (1983). The enzymatological function of heavy metals and their role in electron transfer process of plants. In “Encyclopedia of Plant Physiology, New Series,” (Eds.) A. Lauchli and R. Bialeski , Vol. 15A, pp. 563-596. Springer-Verlag, Berlin and New York.
-

-
- Schmidt, S.P. and Osborn, T.G. (1993). Effects of endophyte-infected tall fescue on animal performance. *Agric. Ecosyst. Environ.*, **44**: 233-262.
- Sekine, T.; Sasakawa, T.; Morita, S.; Kimura, T. and Kuratom, K. (1965). cf. labrotory manual for physiological studies of Rice (Eds.) Yoshida, S.; Forno, D.; Cook, J.B. and Gomez, K. A. (Pub.) International Rice Research institute, Manila, India. 1972.
- Senn, T.L., (1987). Seaweed and Plant Growth. (Publ.)Clemson University, Clemson, U.S.A. pp.192.
- Shing, H.B. and Joshi, B. S. (1960). Ministry of food and Agric. New Delhi. (Publ.) *Farm Bull.* No-56.pp 235.
- Singh, V. P. and Purohit, S. (2003). Research Methodology in Plant Sciences. (Publ.) Scientific Pub., Jodhpur. pp.167.
- Singh, H.B. and Joshi, B.S. (1960). Farm Bull. No. 56. Ministry of food and Agric. New Delhi.
- Sitakara Rao, V. and Tipnis, U.K. (1967). Chemical constituents of marine algae from Gujarat coast. *Proceeding and Seminar on Sea. Salt and Plants*, CSMCRI, Bhavanagar, pp: 277-288.
- Sivasankaria, S.; Venkatesalu, V. and Chandrasekarana, M. (2006). Effect of seaweed extracts on the growth and biochemical constituents of *Vigna sinensis*. *Biores. Technol.*, **97**:1745–1751.
- Stephenson, W. A. (1974). Seaweeds in agriculture and horticulture. Mexico, USA pp. 241. (publ). Faber London.
- Stephenson, W. M. (1966). The effect of hydrolyzed seaweed on certain plant pests and diseases. *Proc. Int. Seaweed. Symp.* **5**: 405-415.
- Stirk, W.A. and Staden, J.V. (2002). Desorption of cadmium and the reuse of brown seaweed product as biosorbents. *Botanica Marina*, **45**: 9-16.
- Subba Rao, P.V. (2007). Mineral composition of edible seaweed *Porphyra vietnamensis*. *Food chem.*, **102**: 215-218.
- Takahashi , M. and Asada, K. (1977). Manganese binding to sodium cyanide- treated chloroplast: effects of light and redox potentials on the binding. *Plant cell Physiol.*, **18** : 807-814.
-

-
- Tarjan, A.C. (1977). Kelp derivatives for nematode-infected citrus trees. *J. Nematology* 9:287.
- Tay, S.A.B.; Palni, L.M.S and Macleod, J.K. (1987). Identification of cytokinin glucosides in a seaweed extract, *J. plant Growth Reg.*, 5 : 133-138.
- Taylor, I.E.P. and Wilkinson, A.J. (1977). The occurrence of gibberellins and gibberellins-like substances in algae. *J. Phycologia.*, 16: 37-42.
- Taylor, J.S.; Harker, K.N.; Robertson, J.M. and Foster, K.R. (1990). The effect of a seaweed extract containing cytokinin on the growth and yield of barley. *Canadian J Plant Sci.*, 70 (4): 1163-1167.
- Temple, W.D. and Bomke, A.A. (1989). Effects of kelp foliar applications on bean crop growth. *Plant and Sci.*, 117:85-92.
- Torrey, J G. (1976). Root hormones and plant growth. *Annu. Rev. Plant Physiol.*, 27: 435-459.
- Toth, S.J.; Prince, A.L.; Wallace, A. and Mikkelsen, D.S. (1948). Rapid quantitative determination of eight mineral elements in plant tissue by systematic procedure involving use of flame photometer. *Soil Sci.*, 66: 459-466.
- Tourte, L.; Bugg, R.L. and Shennan, C. (2000). Foliar-applied seaweed and fish powder do not improve yield and fruit quality of organically grown processing tomatoes. *Biological Agr. and Hort.*, 18(1):15-27.
- Tressler, D. K. and Lemon, J. M. (1951). Marine products of commerce, 2. ed., Reinhold Publ. Corp., New York. Pp- 782.
- Untawale, A.G. and Dhargalkar, V.K. (1975). Report on the seaweed resources of the Goa coast. *Tech. Rep.* (National Institute of Oceanography., Goa) : 1-10.
- Untawale, A.G.; Dhargalkar, V.K. and Agadi, V.V. (1983). Checklist of marine algae from Indian coast *Tech. Report-* National Institute of Oceanography., Goa. Pp. 42..
- Van Steveninck, R.F.M. (1965). The significance of calcium on the apperent permeability of cell membranes and the effects of substitution with other divalent ions. *Physiol Plant.*, 18 : 54-69.
- Verkleif, F.N. (1992). Seaweed extracts in agriculture and horticulture: a review. *Biol-Agr. and Hort. Int. J.*, 8 (4):309-324.
-

- Vernieri, P.; Borghesi, E.; Ferrante, A. and Magnani, G. (2005). Application of biostimulants in floating system for improving rocket quality. *J Food Agric Environ*, 3:86–88.
- *Vicente, N.; Moreteau, J.C. and Escoubet, P. (1980). Etude de l'évolution d'une population de *Pinna nobilis* L. (Mollusque Eulamelibranche) au large de l'anse de la Palud (Parc Nacional de Port-Cros). *Trav. Sci. Parc. Natl. Port-Cros*, 6: 39-67.
- Viera, M.P.; Gomez, J.L.; Pinchetti, G.; Vicoose, C.; Bilbao, A.; Suarez, S.; Haroun, R.J. and Izquierdo, M.S. (2005). Solubility of three red macro algae as a feed for the abalone *Haliotis tuberculata coccinea* Reeve. *Aquaculture*, 248: 75-82.
- Whapham, C. A.; Blunden, G.; Jenkins, T. and Hankins, S. D. (1993). Significance of betaines in the increased chlorophyll content of plants treated with seaweed extract. *J Appl Phycol.*, 5:231–234.
- Wilczek, C.A. and Ng, T. (1982). The promotion of seed germination in table beet by an aqueous seaweed extract. *Hort Science*, 17: 629-630.
- Wong, K.H. and Cheung, P.C.K. (2000). Nutritional evaluation of some subtropical red and green seaweeds: Part I—Proximate composition, amino acid profiles and some physico-chemical properties. *Food Chem.*, 71: 475–482.
- *Yan, J. (1993). Influence of plant growth regulators on turfgrass polar lipid composition, tolerance to drought and saline stress, and nutrient efficiency. Ph.D. Dissertation, Dept. of Crop and Soil Environmental Sciences, Virginia Polytechnic Institute and State University, VA.
- Yan, X.; Nagata, T. and Fan, X. (1998). Antioxidant activities in some common seaweeds. *Plant Foods Human Nutr.*, 52 : 253–262.
- Yang Y. (2002). Chinese Herbal Medicines: Comparisons and Characteristics (Publ.) Churchill Livingstone, London, pp.223.
- * Yoshida (2001), *Mizujoho*, Vol.21, No.7, pp. 12-14 (in Japanese).
- Zahid, P. B. (1999). Preparation of organic fertilizer from seaweed and its effect on the growth of some vegetable and ornamental plants. *Pakistan J. of Biol. Sci.*, 4 (2):1274-1277.

-
- Zaragoza, E. S.; Lopez, D.G. and Diaz, G.P. (2002). Gross chemical composition of three common macroalgae and a sea grass on the Pacific coast of Baja California, Mexico. *Hydrobiologia*, 12 (2): 113-118.
- *Zhang, X. (1997). Influence of growth regulators on turfgrass antioxidant status, drought tolerance, and growth response. Ph.D. Dissertation, Dept. of Crop and Soil Environmental Sciences. Virginia Polytechnic Institute and State University, VA.
- Zhang, X. and Schmidt, R.E. (1997). The impact of growth regulators on the α -tocopherol status in water-stressed *Poa pratensis* L. *Int Turfgrass Res J.*, 8:1364–1373.
- *Zodape S.T.; Mukherjee, S.; Reddy, M.P. and Chaudhary, D. R. (2009) .Effect of *Kappaphycus alvarezii* (Doty) Doty ex silva. extract on grain quality, yield and some yield components of wheat (*Triticum aestivum* L.) *Int. J. of Pl. Prod.*, 3 (2): 97-101.
- Zubia, M.; Payri, C.E.; Deslandes, E. and Guezennec, J. (2003). Chemical composition of attached and drift specimens of *Sargassum mangarevense* and *Turbinaria ornata* (Phaeophyta: Fucales) from Tahiti, French Polynesia. *Bot. Mar.* , 46 : 562–571.

* originals not seen

<http://www.alcoorganic.org/pdf/soil>.

<http://www.shorganic.com/page33.html>.

<http://www.googleearth.com>.