

Chapter VI
Bibliography

- 1) Anandhi S., and Ramanujam M. P., (1992) Effect of salicylic acid on black gram (*Vigna mungo*) cultivars. *Indian J. Physiol.*, Vol. 2(2) 135 – 141.
- 2) Arjunan. A. and Gopalakrishnan S. (1980). Effect of Calcium-sodium interaction on the degree of hydration in groundnut.
- 3) Arya, A. and Chauhan, R. (1995). Seed mycoflora of chickpea. *Botanica Indica*. 23(2) : 293 – 295.
- 4) Bhowal, S., Boran, P. K., Senapoty, D. and Rabha, B. K. (2006). Seed-borne micoflora in groundnut and their management. *Indian phytopathology*. 59(3) : 398.
- 5) Dawar, S., Ghaffar, A., Shaukat, S. S. and Rasheed, S. (2004). Seed borne mycoflora of groundnut. *Pakistan Journal of Botany*. 36(1) : 199-202.
- 6) Deepali Sabale, R. B. Thoke and B. A. Kore (2008). Seed Mycoflora of Groundnut. *Bioinfolet* 5(3) : 263 – 266.
- 7) Dutta R. K., Deka, N. c. and Neog. K. (2007) effect of different levels of sulphur on groundnut (*Arachis hypogaeae* L.) in sandy loam soil of Assam.
- 8) Dwivedi, S. N. and Shukla, T. N. (1990). Mycoflora of Gram seed in different agro-climatic regions and their pathology. *Indian phytopathology* 43(1) : 98 – 99.
- 9) Frank. A. B. (1885). Cf. Principles of seed pathology. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and distributors. Delhi. 3p, 1993.
- 10) Girisha¹, S. T. and Raju² N. S. (2007). Effect of sewage water on seed germination and vigour index of different varieties of groundnut (*Arachis hypogaea* L.)

- 11) Gupta, V. K. and Chouhan, J. S. (1970). Seed-borne fungi and seed health testing in relation to seedling diseases of groundnut. *Indian phytopathology*. **23** (4) : 622 – 625.
- 12) Gurbaksh Singh, Nirmal Sekhon Maniit K. (1980). Effect of Phenolic compounds on the yield potential of Gram (*Cicer arietinum* L.) *Indian Journal of Plant Physiology*. Vol. XXIII.1.
- 13) ISTA. (1966). International Rules for Seed Testing. *Seed Sci. Technol.* **24** : 335 – 337.
- 14) Jadhav D. N. (2008) Studies on Indigenous Arbuscular mycorrhizae (Am. fungi) Inoculated to groundnut (*Arachis hypogae* L. Cv. JL. 24) under unsterilized soil condition. *Bioinfolet* **5**(2) : 116 – 119.
- 15) Jain R., Chowdhary L. and Chatterjee C. (1997) Influence of calcium on growth, composition and yield of black gram. *Indian J. Plant Physiol*; Vol. **2**(3) 221 – 224.
- 16) Jat, R. G., Goyal J. P. and Jain S. C. (2004). Effect of fertilizers on leaf blight disease and pod yield of groundnut. *J. Mycol. Pl. Pathol.*, Vol **34** (1).
- 17) Krishnasastry; K. S., Malathi Chari, Prasad T. G., Udaykumar, M. and Sashidhar, V. R., Marh (1985). Flowering pattern and Pod development in bunch types of ground. *Indian J. Plant Physica*; Vol. XXVIII No. **1** pp., 64 – 71.
- 18) Lalithakumari, D., Govindaswamy, C. V. and Vidhya Sekaran, P. (1971). Cf. *Seed Pathology*. Vol. I, Neergaard, P. The MacMillan Press Ltd.; London, **24p**, 1977.
- 19) Mathur, R. L. and Sharma, L. c. (1971). Field assay of Seed dressing fungicide for groundnut seeds. *Madras Agri. J.* **58** : 199 – 202.

- 20) Menthe, P. R., Singh, B., Mathur, S. C. and Singh, S. B. (1953). Cf. Seed Pathology. Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 18p, 1977.
- 21) Mittal, R. K. and Mathur, S. B. (1993). Pathology. Part I, ICAR, New Delhi, India and Danish Government Institute of Seed Pathology, Denmark, 178P.
- 22) Muley, S. M. and Baig, M. F. (2007). Study on mycoflora of oil seeds. Bioinfolet. 4(2) : 146 – 150.
- 23) Neergaard, P. (1977). Seed Pathology. Vol. I, The MacMillan Press Ltd., London, pp. 53 – 60.
- 24) Nobel, M., De Tempe, J. and Neergard, P. (1958) Cf. *Principles of Seed Pathology* Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and Distributors. Delhi. 3p. 1993.
- 25) Orton, C. R. (1931). Cf. *Principles of Seed Pathology*. Agarwal, V. K., Sinclair, J. B., Vol. I, CBS Publisher and Distributors, Delhi. 3P. 1993.
- 26) Patale, S. S. & Mali, V. S. (2009). Efficacy of trichoderma in management of collar rot of groundnut. Bioinfolet 6(2) : 154 – 156.
- 27) Portor, R. H. (1949). Cf. *Principles of Seed Pathology*. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and Distributors, Delhi. 3p. 1993.
- 28) Prasad, V. V. S., Pandey, R. K. and Saxena M. C. (1978) Physiological analysis of yield variation in Gram (*Cicer arietinum* L.) *Indian Journal of Plant Pathology*, Vol. XXI (3).
- 29) Prevost, B. (1807). Cf. *Principles of Seed Pathology*. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and Distributors, Delhi. 2p. 1993.
- 30) Reddy, S. H. Sugunakar, M. and Reddy, B. M. (1998). Effect of seed treatment with fungicides and insecticides on seed borne fungi. storage insect

pest, seed viability and seeding vigour of groundnut seed research. 26(1) : 62
– 72.

- 31) Richardson, M. J. (1979). Cf. *Principles of Seed Pathology*. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and Distributors, Delhi. 3p. 1993.
- 32) Shaikh, A. A., Kumbhar, S. G. and Jawale S. M. (2007). Integrated nutrient management under black gram – wheat cropping sequence. *Bioinfolet* 4(2) ; 119 – 121.
- 33) Sheoran I. S. and Garg O. P. (Dec. 1983). Effect of different types of Salinities on Gram during germination seedling growth and water relations. *Indian J. Plant Physiol.* Vol. XXVII, No. 4, PP. 363 – 369.
- 34) Shinde M. R. and S. B. Bhamburdekar (2008). Effect of aluminium toxicity on the activity of enzyme lipase during groundnut seed germination 5(4) : 362 – 363.
- 35) Singh A. L. and Chaudhari Vidya (1996). Interaction of sulphur with phosphorus and potassium in Groundnut nutrition in calcareous soil, *Indian J. Plant Physiol.*, Vol. 1(1) pp. 21 – 27.
- 36) Singh, K. and Singh, A. K. (2004). Detection of Seed mycoflora of gram (*Cicer arietinum*). *Indian Phytopathology*. 57(3) : 361.
- 37) Singh, S. D., Rawal, P. and Bhargava, N. K. (2004). Pathogenic Potential and control of seed mycoflora of Groundnut (*Arachis hypogae*). *Journal of mycology and plant pathology*. 34(2) : 687 – 690.
- 38) Sonwalkar S. N., Thobre V. K., Bhondve T. S. and Shaikh A. A. (2006) Effect of various methods of weed control and planting layouts on nutrient uptake by weeds in groundnut (*Arachis hypogae*). *Bioinfolet* 3(4) : 319 – 322.

- 39) Thoke, R. B. (1989). Fungi of Maharashtra. Ph.D. Thesis. Pune University. Poona.
- 40) Tillet, D. M. (1775). Cf. *Principles of Seed Pathology*. Vol. I, Agarwal, V. K., Sinclair, J. B., CBS Publisher and Distributors, Delhi. 2p. 1993.
- 41) Toorray, N. K., Verma, K. P., Thakur, M. P. and Sinha, A. K. (2005). Evaluation of chickpea accessions by standard blotter method. *Advances in plant Sciences*. 18(1) : 31 – 37.
- 42) Verma, V. (1984). A Text Book of Economic Botany. Emkay Publications Delhi. 1 – 196.
- 43) Wadikar M. S. and Shinde A. A. (2008) Effect of different fungicides on root rot fungi of chickpea. *Bioinfolet* 5(2) : 98.
- 44) Wallen, V. R. (1965). Cf. *Seed Pathology*. Vol. I, Neergaard, P. The MacMillan Press Ltd., London, 832p. 1977.
- 45) Zaidi., B. I. S., Khan, M. I. and Saxena, S. K. (1991). Effect of fungicide on mycoflora of Chickpea seeds. *Indian Phytopathology* 44(3) : 394 – 395.

580
YAD



T16292