

Table No.	List of Tables
4.1	Bulk density values of Newara samples.
4.2	Weight percent frequency in grams for the sediments from different levels of Newara beach.
4.3	Cumulative weight percent frequency for the sediments from different levels of Newara beach.
4.4	The comparative study of grain size parameters in the southern, central and northern parts of the Newara beach having low - tide, high - tide and dune samples.
4.5a	Population percentage of different minerals in Newara sand from low - tide zones.
4.5b	Population percentage of different minerals in Newara sand from high - tide zones.
4.5c	Population percentage of different minerals in Newara sand from base of dunes.
4.5d	Population percentage of different minerals in Newara sand from dunes.
4.6a	The frequency variation of different

heavy minerals in Newara beach sand
(Low - tide zones).

4.6b The frequency variation of different
heavy minerals in Newara beach sand
(High - tide zones).

4.6c The frequency variation of different
heavy minerals in Newara beach sand
(Base of dunes).

4.6d The frequency variation of different
heavy minerals in Newara beach sand
(Dunes).

Plate No.

List of Plates

- 2.1 Field photograph - southern and central part of Newara beach showing broad surf zone and also gentle slope.
- 2.2 Field photograph - alternate ripples of marine origin indicating characteristic presence of heavy dark coloured sand with whitish light sand.
- 2.3 Field photograph - ripples of eolian origin indicating characteristic presence of whitish light sand and absence of dark heavy sand.
- 3.1 Field photograph - conglomeratic laterite consisting of subrounded to subangular pebbles.
- 3.2 Photomicrograph - porphyritic texture within massive basalt indicating plagioclase laths.(12 x).
- 3.3 Photomicrograph - euhedral and subheral phenocrysts of augite. (12 x).
- 3.4 Photomicrograph - characteristic presence of opaque oxides - ilmenite and magnetite in the form of minute dusty grains in basalt. (12 x).

- 3.5 Photomicrograph - steel black coloured ilmenite grains with brownish borders in reflected light (12 x).
- 3.6 Photomicrograph - formation of leucoxene i.e. 3rd stage of ilmenite alteration under ore microscope. (40 x).
- 3.7 Photomicrograph - columnar tourmaline grain with other opaque minerals. (12 x).
- 3.8 Photomicrograph - euhedral crystal of zircon in association with other opaque minerals. (12 x).
- 4.1 Field photograph - heavy mineral rich zone at the base of the dune.