

VIII. REFERENCES

- Agashe, S.N. (1977). Prototaxopitys andrewsii a new combination for Prototaxoxylon andrewsii. Agashe and Chitnis. Geophytology, 7(2) : 278-279.
- Agashe, S.N. & Chitnis S.R. (1971). Studies on the fossil gymnosperms Part-3 : Prototaxoxylon andrewsii, a new species of taxinean wood from the Lower Gondwana Strata. Palaeontographica, 133 B (1-3) : 52-60.
- Agashe, S.N. & Gowda, P.R. (1979) : Anatomical study of gymnospermous wood from Lower Gondwana of Maharashtra. Phytomorphyology, 28 : 269-275.
- Agashe, S.N. & Gowda, P.R. (1982). Studies of fossil gymnospermous wood Part-IV : Two new species of petrified woods from the Lower Gondwana (Permian) Strata. Indian J. Bot., 4 : 73-78.
- Arbor, E.A.N. (1905). Catalogue of the fossil plants of the Glossopteris flora in the Department of Geology, British Museum, Natural History, Longman & Co., London.
- Archangelsky, S.(1958). Estudio Geologico of Palaeontologico del. Bajo de la Leone (Santa Cruze). Acta Geol. Lilloana, 2 : 5-133.

- Bajpai, U. (1985). A gangamopteroid leaf from Raniganj formation
Geophytology, 15 : 60-63.
- Banerjee, Manju (1987). A study of the frequency of megaplant
fossil in Raniganj formation (Upper Permian) of Raniganj
Coalfield. Geo.Surv.India.Spl.Publ., 1 : 261-288.
- Banerjee, M.& D. Rozario, A.(1987). Palynostratigraphy and
Environment of deposition in the Lower Gondwana
sediments of chuperbhita coalfield, Rajmahal Hills.
Workshop on : Concepts, limits & extension of the
Indian Gondwana, Lucknow, (Abstracts) : 10-11.
- Bharadwaj,D.C.(1953). Jurassic wood from the Rajmahal Hills,
Bihar. Palaeobotanist, 2 : 59-69.
- Bharadwaj,D.C.(1966). Distribution of spores and Pollengrains
dispersed in the Lower Gondwana formation of India.
In : Symposium on Floristics and Stratigraphy. Gondwana
Land (Special Session), Birbal Sahni Institute of
Palaeobotany, Lucknow, PP. 66-84.
- Bharadwaj,D.C.(1971). Palynostratigraphy of Lower Gondwana
succession in India. Annals.Geol.Dept., A.M.U.,Aligarh,
5-6 : 390-419.
- Bharadwaj,D.C.(1974). Palaeobotany of Talchir and Karharbari
formations and Lower Gondwana glaciation in : Surange,
K.R. et al (Eds.). Aspects and appraisal of Indian
Palaeobotany, Birbal Sahni Institute of Paleobotany,
Lucknow, PP. 369-385.

- Biradar, N.V. (1967). Studies in fossil plants from Kota Maleri beds and embryology of genus Phoenix Linn. A Ph.D. Thesis submitted to University of Poona.
- Biradar, N.V. & Bonde, S.N. (1981). Nandorixylon saksenae gen. et. sp. nov. - A new gymnospermous wood from Lower Gondwana Strata. - Indian J. Bot., 4: 73-78.
- Blanford, W.T. (1877). The Palaeontological relations of the Gondwana system : a reply to Dr. Feistmantel, Rec. Geol. Surv. India. 11(1) : 1-160.
- Blanford, W.T. (1879). Manual of the Geology of India.
- Bose, M.M. (1966-a). Significance of fossil plants in the Indian Gondwana stratigraphy. Sci. and Cult., 32 : 532-534.
- Bose, M.M. (1966-b). Fossil plant remains from the Rajmahal and Jabalpur series in the Upper Gondwanas of India. Symp. Florist. Strat. Gond. Land., B.S.I.P., Lucknow : 143-154.
- Brongniart, Adolphe (1822). Sur la classification et al distribution des vegetaux fossiles en general, et sur ceux des terrains de sediment superior on particulier. Mus. Histoire nat., Paris Mem. 8 : 203-348.
- Bunbury, C.J.F. (1861). Notes on collection of fossil plants from Nagpur, Central India, Q. Jl. Geol. Soc. Lond., 17 : 325 - 346.
- Chandra, S. & Chandra, A. (1988). Vegetational changes and their climatic implications in coal bearing Gondwana in : Venatachala B.S. & Maheshwari, H.K. (eds.). Concepts, limits and extension of Indian Gondwana, Palaeobotanist, 36 : 000-000.

- Chandra, S. and Prasad, M.N.V. (1979-a). Two new fossil woods from Kamthi beds of Chandrapur District, Maharashtra. Palaeobotanist, (In Press).
- Chandra, S. and Prasad, M.N.V. (1979-b). Two new fossil woods from Kamthi beds of Chandrapur District, Maharashtra. Palaeobotanist (In Press).
- Chandra, S. & Prasad, M.N.V. (1981). Fossil plants from the Kamthi formation of Maharashtra and their biostratigraphic significance. Palaeobotanist, 29-29 : 99-121.
- Chandra, S. & Rigby, J.F. (1981). Lycopoid, Sphenopsid and cycadaceous remains from the Lower Gondwana of Handappa, Orissa, Geophytology, 11 : 214-219.
- Chandra, S. and Rigby, J.F. (1983). The Filicales from the Lower Gondwanas of Handappa Orissa. Palaeobotanist, 31 : 143-147.
- Chandra, S & Surange, K.R. (1976). Cuticular studies of the reproductive organs of Glossopteris Part-1. Dictyopteridium feistmanteli sp.nov. attached on Glossopteris tenuinervis. Palaeontographica, 156-B (4-6) : 87-102.
- _____ " _____ (1977-a). Cuticular studies of the reproductive organs of Glossopteris Part-2. Cistella type fructification Plumsteadia elliptica gen. et.sp. nov., attached on Glossopteris taenioides Feistmantel. Palaeobotanist, 23(3) : 161-174.

- _____ „ _____ (1977-b). Cuticular studies of the reproductive organs of Glossopteris Part-3. Jambadostrobus and Venustostrobus borne on Glossopteris leaves.
Palaeontographica, 164 B (4-6) : 127-152.
- _____ „ _____ (1977-c). Cuticular studies of the reproductive organs of Glossopteris Part-4. Venustostrobus indicus sp. nov.
Palaeobotanist, 24(3) : 149-160.
- _____ „ _____ (1977-d). Some scale leaves and sporangia from the Raniganj coalfield, India. Palaeobotanist, 24(3) : 245-253.
- _____ „ _____ (1979). Revision of the Indian species of Glossopteris Monograph No.2. Birbal Sahni Institute of Palaeobotany, Lucknow, India.
- Chitale, S.D. (1949). On a new species of Dadoxylon, D. chandensis sp. nov. from the District Chandra C.P.J.
- Cotter, G. de P. (1917). A revised classification of the Gondwana system. Rec. geol. surv. India, 48 : 23-33.
- Dasgupta, H.C. (1915). On the correlation of Kamthi beds. Proc. Indian Assoc. Sci., 1 : 5-19.
- Dasgupta, H.C. (1928). Batrachian and reptilian remains found in the Panchet beds at Deoli, Bengal. Jour. Assiat. Soc. Bengal, 24 : 473-479.
- Evans, W.P. (1934). Microstructures of New Zealand lignite-III. Sci. Tech., 15(6) : 365-385.
- Evans, W.P. (1937). Note on the flora which yielded the Tertiary lignite of Centerbury otago and Southland. N.Z. Sci. Tech., 19 : 190.

- Feistmantel,O. (1870). The fossil flora of Gondwana System-I.
The flora of the Talchir-Karharbari beds. Palaeont.
Ind.Ser., 12 3 : 1-64.
- Feistmantel,O. (1876). Notes on the age of some fossil floras
of India. Rec.Geol.Surv.India, 9(2) : 28-42.
- Feistmantel,O. (1877). Notes on the fossil flora of India-XI.
Notes on Plant fossils from Barakar district (Barakar
group). Mem.Geol.Surv.Indi.Palaeont.Indica,Ser., 11.2(1)
: 1-80
- Feistmantel,O.(1879). The fossil flora of the Gondwana system
: Lower Gondwana. The flora of the Talchir -
Karharbari beds. Mem.Geol.Surv. India.
Palaeont. Indica,Ser. 12, 3 : 1-48.Suppl.1881 : 49-64.
- Feistmantel,O. (1880). The fossil flora of the Lower Gondwana
- 1. The flora of the Damuda and Panchet divisions.
Mem.geol. Surv.India, Palaeontindica, Ser. 12,3 :1-77.
- Feistmantel,O. (1881). The fossil flora of the Gondwana System
II. The flora of Damuda Panchet Division. Mem.geol.
Surv.India, Palaeont.indica, Ser.12, 3(3) : 78-149.
- Feistmantel,O.(1882). The fossil flora of the Gondwana System.
The fossil flora of the South Rewa Gondwana Basin.
Mem.Geol.Surv. India Palaeontindica,Ser.12, 4(1):1-52.
- Feistmantel,O. (1886). The fossil flora of the Gondwana system.
Fossil flora of some of the coal fields in West Bengal.
Mem.geol.Surv.India Palaeont.indica, Ser.12,4(2):1-66.
- Feistmantel,O. (1889). Geological & Palaeontological relations
of the coal and Plant bearing beds of Palaeozoic and
Mesozoic age in the Eastern Australia and Tasmania with
reference to fossil flora. Mem.Geol.Surv.N.Sw.(Palaeonf).
3 : 1-183.

- Fox, C.S. (1931). The Gondwana system and the related formations. Mem.geol.Surv.India, 58 : 1-241.
- Ganguli, S.K. (1959). Palaeontological study of Lower Gondwana rocks including the coal seams around Chirmiri of Surguja district, Madhya Pradesh.
J.Geol.Min.Metall.Soc.India, 31 : 155-166.
- Ganju, P.N. (1943). On a new species of Psygmyphyllum (P. sahani) from the Lower Gondwana beds of Kashmir.
J.Indian Bot.Soc., 22 : 201-207.
- Greguss, P. (1955). Identification of living gymnosperms on the basis of Xylotomy. 1 : 263 Budapest.
- Halle, T.G. (1908). Zur Kenntnis mesozoischer Equisetales, Schweden : Kgl. Svenska Vetenskapsakad. Handlingar, 43(1) : 1-40.
- Halle, T.G. (1911). On the geological structure and history of the Falkland Islands, Bull.geol.Instn.Univ. Upsala, 11 : 1-117.
- Hartig, Th. (1848). Beiträge Geschichte Pflanzen und Kenntniss der norddeutschen Braun Kohlen flora. Bot.Ztg. 10 : 185-190.
- Hoeg, O.A. and Bose, M.N. (1960). The Glossopteris flora of Belgian Congo with a note on some fossil plants from the Zambezi basin. (Mozambique). Ann.Mus.Congo.Dolge. (Sci. Geol.) 32 : 1-106.
- Hughes, T.W.H. (1881). Notes on the South Rewa Gondwana basin. Mem.Geol.surv.Ind., 14 : 126-138.
- Hughes, T.W.H. (1884). Southern coalfields of Rewa Gondwana basin.

- Kapoor, H.M. (1979). Gondwana of Kashmir - a reappraisal in : Laskar, B. & Raja Rao, C.S. (eds.). Proc.IV Inf.Gondw. Symp., Calcutta, 2 : 443-462.
- Kar,R.K.(1966). Palynology of the Barren Measures sequence from Jharia coalfield, Bihar, India.1. Summary and Discussion. Sym.Florist.Stratigr. Gondwanaland, Sahni, Inst. Palaeobot. 121-127.
- Kar,R.K. (1968). Studies in Glossopteris flora of India. 36. Plant fossils from Barren measures succession of Jharia coalfield. The Palaeobotanist, 16(3) : 243-248.
- Kar,R.K. (1970-b). Palynological distinction between Upper Permian and Lower Triassic in Raniganj coalfield, Bengal, India.
- King, W.(1881). Geology of Pranhita Godavary valley. Mem. Geol.Surv.Ind., 18 : 151-311.
- Krausel,R. and Dolianti,E.(1958). Gymnospermen holzer ausdem Palaeozoikum Brasilliens. Palaeontographica, 104 B(4-6) : 115-137.
- Krausel, R. & Jain, K.P.(1964). New fossil coniferous woods from the Rajmahal Hills, Bihar, India. The Palaeobotanist 12(1) : 59-67.
- Kulkarni,S.(1971). Barakaroxylon monocanalosum sp.nov. from the Jharia coalfield. Palaeobotanist, 18(3) : 309-311.
- Lele,K.M. (1955). Plant fossils from Parsora in the South Gondwana basin, India. The Palaeobotanist, 4 : 23-24.

- Lele, K.M. (1956). Plant fossils from Parsora in the South Rewa Gondwana Basin, India. Palaeobotanist, 4 : 23-24.
- Lele, K.M. (1962). Studies in the Indian Middle Gondwana flora. 1. On Dicroidium from the South Rewa Gondwana basin. Palaeobotanist, 10(10-2) : 48-68.
- Lele, K.M. (1964). The problem of Middle Gondwana of India. Proc. 22nd Int. Geol. Congr., India 9 : 182-202.
- Lele, K.M. (1966). Quest for the early straces and subsequent development of Glossopteris flora in the Talchir Stage in : Symposium of floristics and stratigraphy of Gondwana land. Birbal Sahni Institute of Palaeobotany, Lucknow, PP. 85-97.
- Lele, K.M. (1969). Studies in the Glossopteris flora of India. 39. Alatocarpus gen. nov. - a new Platyspermic seed from the Singrauli coalfield. The Palaeobotanist, 12(1) : 7-17.
- Lele, K.M., Swarup, P. and Singh, J.N. (1968). Occurance of Plant fossils in the Lower Gondwana succession of Singrauli coalfield, U.P. J. Palaeont. Soc., India, 11:8-17.
- Lepekhina, V.G. (1972-a). Woods of Palaeozoic pycnoxylic gymnosperms with special reference to North Eurasia representatives. Palaeontographica, 138 B(1-4) : 44-106.
- Mahabale, T.S. (1964). The Kota Maleri stage and Indian Mesozoic floras. Proc. Xth. Inter. Bot. Congr. Edinburgh, (Abst.) : 19-20.

- Mahabale, T.S. (1966). Flora of the Deccan : Past and Present. Presidential Address, 53rd Ind.Sci.Congr., Chandigarh:1-30.
- Mahabale, T.S. & Rao, S.V. (1968). Fossil flora of Rajahmundry area, Bull.Ind.Nat.Sci.Acad., New Delhi. Symposium of Deccan Trap Country held at Poona, 45 : 192-214.
- Mahabale, T.S. & Vagyani B.A. (1980) : On the occurrence of Kamthioxylon a new genus of gymnospermous wood from Adhari (Maharashtra). Biovigyanam, 6 : 131-134.
- Maheshwari, H.K. (1965-a). Studies in Glossopteris of India 22. On some species of genus Glossopteris from the Raniganj Stage of East Raniganj Coal field, Bengal, India. The Palaeobotanist, 13(2) : 129-143.
- , (1965-b). Studies in Glossopteris flora of India. On two fructifications from the Raniganj stage of Rajmahal coalfield, Bengal, India. The Palaeobotanist, 13(2) : 144-147.
- , (1965-c). Studies in the Glossopteris flora of India-24. On two new species of fossil woods from Raniganj Stage of Raniganj coalfield, Bengal, India. The Palaeobotanist, 13(2) : 144-147.
- , (1966). Studies in the Glossopteris flora of India -28. On some fossil woods from the Raniganj Stage of Raniganj coalfield, Bengal, India. The Palaeobotanist, 15(3) : 243-257.
- Maheshwari, H.K. (1966-b). Studies in the Glossopteris flora of India-30. Remarks on the age of the Lower Gondwana beds of Bensoli valley, Santhal Paraganas, Bihar, India. Sym.Florist.Strat.Gond.Land., B.S.I.P.Lucknow:110-120.

- Maheshwari, H.K. (1967). Studies in the Glossopteris flora of India-28. On some fossil woods from the Raniganj Stage of Raniganj coalfield, Bengal. Palaeobotanist, 15(3) : 243-257.
- Maheshwari, H.K. (1972). Permian wood from Antarctica and revision of some Lower Gondwana wood taxa. Palaeontographica, 138 B : 1-43.
- Maheshwari, H.K. and Prakash, Gyan. (1965). Studies in the Glossopteris flora of India-21. Plant megafossils from the Lower Gondwana Exposures along Bansoli river in Rajmahal Hills, Bihar. Palaeobotanist, 13(2): 113-128.
- Maheshwari, H.K. & Tiwari, R. (1986). Maheshwariella spinicornuta, a new gymnospermous seed from the Karharbari formation.. Palaeobotanist, 35(1) : 69-72.
- Maithy, P.K. (1965-a). Studies in the Glossopteris flora of India-17. On the genus Rubidgea Tate. Palaeobotanist, 13(1) : 42-44.
- Maithy, P.K. (1966-a). Palaeobotany and stratigraphy of Karharbari stage with particular reference to Giridih coalfield, India. Symp. Florist. Strat. Gond. Land., B.S.I.P. Lucknow : 102-109.
- Maithy, P.K. (1968). Studies in the Glossopteris flora of India-35. On two new fossil plants from the Ganjra Nalla beds, South Rewa Gondwana basin. Palaeobotanist, 16(3) : 219-221.

- Maithy, P.K. (1969-a). Studies in the Glossopteris flora of India-39. On some new plant fossils from the Karharbari beds, Giridih coalfields, Bihar, India. Palaeobotanist, 18(2) : 167-172.
- Maithy, P.K. (1969-b). The contemporaneity of the Karharbari flora in Gondwana in : Santapau, H. et.al. (eds.). J.Sen.Mem.Vol.Bot.Soc.Bengal, Calcutta, 274-298.
- Maithy, P.K.(1971-a). Fossil flora of the Barakar Stage in Auranga coalfield. The Palaeobotanist, 19(1) : 70-75.
- Maithy, P.K.(1971-b). The Lower Gondwana Plants of India and other stratigraphic significance. Septime congress International de stratigraphic et. de. Geologic du carbonifere, Kerfald : 385-390.
- Maithy, P.K. (1974). Gymnospermous woods from the Lower Gondwana. In Aspects and Appraisal of Indian Palaeobotany. B.S.I.P. Lucknow, 163-169.
- Maithy, P.K. (1977). Revision of some fossil plants from the Karharbari formation, Giridih coalfield, Bihar, India. Palaeobotanist, 23(3) : 220-222.
- Maithy, P.K. & Mishra, P.K.(1984). A new Cyanophycean remain from the Karharbari formation, Giridih Coalfield, Bihar. Palaeobotanist, 32(2) : 130-133.
- Margurier, J.(1971). Le genera Australoxylon Dansta flora Gondwaniene. 96 Congr.nat.soc.saratentes. Joulouse, 5 : 99-105.

- Mc Elroy C.T. (1969). Comparative lithostratigraphy of Gondwana sequences : Eastern Australia and Antarctica. Gondwana stratigraphy, Unesco, Paris. PP. 441-466.
- Medlicott, H.B. & Blanford, W.T. (1879-1887). A manual of Geology of India, Chiefly compiled from the observation of the Geological Survey. 1 : 1-144 & 2 : 445-817.
- Oldham, T. (1869). The fossil flora of the Rajmahal hills, Bihar. Mem. Geol. Surv. Ind., Pal. Ind., 1(1) : 1-52.
- _____, (1972-b). Late Palaeozoic gymnosperms of the Soviet Union from Palaeobotanical data. Palaeontol., 2 : 122-129 (In Russian).
- Oldham, R.D. (1893). A manual of Geology of India. 1-543.
- Pant, D.D. (1958). The structure of some leaves and fructi.
- Pant, D.D. & Gupta, K.L. (1968). Cuticular structure of some Indian Lower Gondwana species of Glossopteris. Bgt. Pt.1. Palaeontographica, 124-B : 45-81.
- Pant, D.D. & Singh, Sudha (1976). Cuticular structure and affinities of Cheirophyllum lacerata (Feistmantel) n. comb. Palaeobotanist, 25 : 353-359.
- Pareek, H.S. (1969). Some observations on the lithology of coal bearing beds and the nature of coal of the major Gondwana basins of India. In : Gondwana Stratigraphy I.U.G.S., Unesco, Paris, 883-904.
- Patton, R.J. (1958). Fossil wood from victorien brown coal. Proc. Roy. Soc. Viet. N.S., 70, : 129-143.
- Prasad, M.N.V. (1982). An annotated synopsis of Indian Palaeozoic gymnospermous woods. Rev. Palaeobot. Polynol. 38 : 119-156.

- Prasad, M.N.V. (1986). Xylotaphoflora of the Kamthi formation, Indian Lower Gondwana with remarks on the Bistratigraphic importance of its Taphoflora. Palaeontographics, Abt. Bd. 201 : 111-134.
- Prasad, M.N.V. (1978). Australoxylon from the Kamthi beds of Lower Gondwana, India. Curr.Sci., 47(16) : 597.
- Prasad, M.N.V. & Chandra, S. (1979-a). Some species of Dadoxylon from Kamthi beds of Kanhargaon Chandrapur District, Maharashtra, India. Prof. A.K. Ghosh. Commemoration volume, Calcutta (In Press).
- Prasad, M.N.V. & Chandra, S. (1979-b). Taxinean and Ginkgolean fossil woods from Kamthi formation. Palaeobotanists. (In Press).
- Prasad, M.N.V. & Chandra, S. (1979-c). On two species of Taxopitys from Kamthi formation. Palaeobotanist, (In Press).
- Prasad, M.N.V. & Chandra, S. (1979-d). Two species of Australoxylon from the Kamthi formation. Palaeobotanist (In Press).
- Prasad, M.N.V. & Chandra, S. (1981). Two species of Australoxylon from the Kamthi formation of Chandrapur District, Maharashtra. Geophytology, 11(1): 1-5.
- Rawat, M.S. & Jain, A.K. (1985). Marine Leiosphaerids and associated Palyno-fossils from the Talchir formation of Pranhita-Godavari Graben. Petroleum Asia. J1.8(11) : 168-173.

- Sahni, B. (1931). Revision of Indian fossil plants. Pt. II. Coniferales (Petrifications). Mem. Geol. Surv. India. Palaeont. Indica (n-ser), 11 : 53-124.
- Saksena, S.D. (1952). Correlation of the Gondwana based on the evidence of the plant fossils. Agra Univ. Jour. Res. (Sc). 1 : 1-13.
- Seward, A.C. (1907). Permocarbiniferous plants from Kashmir. Rec. Geol. Surv. India, 36(1) : 57-61.
- Seward, A.C. (1912). Lower Gondwana Plants from the Golabgarh Pass. Mem. Geol. Surv. India Palaeont. Indica, n-ser. 4(3) : 1-10.
- Seward, A.C. & Sahni, B. (1920). Indian Gondwana Plants. A revision. Mem. Geol. Surv. Ind., Pal. Ind. N.S., 7:1-54.
- Singh, T. (1979). Palynostratigraphy of the Permian rocks of Siang District, Arunachal Pradesh. In : Verma P.K. (Ed.) Metamorphic rock sequences of the eastern Himalaya. K.P. Bagchi & Co., Calcutta.
- Singh, et al. (1982). Upper Palaeozoic flora of Kashmir Himalaya, Palaeobotanist, 30(2) : 185-232.
- Srivastava, P.N. (1957). Studies in the Glossopteris flora of India-4. Glossopteris, Gangamopteris & Palaeovittaria, from the Raniganj coalfield. Palaeobotanist, 5(1):1-45.
- Srivastava, S.C. (1968). Two new species of Glossopteris from the Triassic of Nidpur, M.P. India, is Son Commomoration Volume, Bull. Bot. Soc. Beng., 200-303.
- Srivastava, S.C. (1973). Palynostratigraphy of the Giridih Coalfield. Geophytology, 3(2): 184-194.

- Srivastava, S.C. (1988). Palynology of Kamthi formation in Gondwana Graben. Palaeobotanist, 36 :
- Srivastava & Jha (1987). Palynology of Kamthi formation in Gondwana Graben. Palaeobotanist, 36 :
- Sternberg, G.K. (1825). Versuch einer geognostischen Botanischen Dowstellung der flora der voruedd. 1(4) : 1-48. Leipsig and Prague.
- Surange, K.R. (1957). Studies in the Glossopteris flora of India-6. Plant fossils from Talchir beds of South Rewa Gondwana basin. The Palaeobotanist, 5 : 82-90.
- Surange, K.R. (1966). Distribution of Glossopteris flora in Lower Gondwana formation of India. Symp. florist. Strat. Gond. Land., B.S.I.P., Lucknow, 55-68.
- Surange, K.R. (1974). Other Lower Gondwana gymnospermous plants. An Aspects and appraisal of India. Palaeobotany, B.S.I.P., Lucknow, 170-178.
- Surange, K.R. & Chandra, S. (1973-a). Dictyopteridium sporiferum Feistmantel female cone from the Lower Gondwana of India. Palaeobotanist, 20(1) : 127-136.
- Surange, K.R. & Lele, K.M. (1956). Studies in the Glossopteris flora of India-6. Plant fossils from Talchir beds of South Rewa Gondwana basin. Palaeobotanist, 5(2):82-90.
- Surange, K.R. & Saxena, Y.N. (1958). Studies in the Glossopteris flora of India-10. Dadoxylon barakarensis sp.nov. from the Jharia coalfield, India.
- Tasch, P. et al. (1975). Estheriids of the Indian Gondwanas significance for continental fit. in : Campbell, K.S.W. (Ed.)-Gondwana Geology, Australian National Univ. Press, PP. 443-452.

- Thomas, H.H. (1952). A Glossopteris with whorled leaves.
Palaeobotanist, 1 : 435-438.
- Tiwari, R.S. (1975). Palynological composition of the basal Gondwana in India. Bull.Soc.belge.Geol. 84: 11-17.
- Tiwari, R.S. et al (1987). Palynological dating of Olive - green shales Underlying the Atgarh Sandstone in Mahanadi Basin. Curr.Sci., 56 : 1150-1153.
- Tiwari, R.S. & Tripathi, A. (1988). Palynological zones and their climatic inference in the coal bearing Gondwana of the peninsular India. Palaeobotanist, 36.
- Trivedi, B.S. & Mishra, J.P. (1970). Triassic miospore assemblage from Nidhpur, district Sidhi, M.P. J. Palaeont.Soc., India, 14 : 14-46.
- Vagyani, B.A. & Mahabale, T.S. (1974). A new species of fossil wood, Planoxylon stopes from Adhari, M.S. Palaeobotanist, 21(2) : 211-215.
- Vagyani, B.A. & Jamane, M.R. (1986). A new species of Agathioxylon from the Kamthi formation of Chandrapur district, Maharashtra.
Proc.Spl.Ind.Geo.Con., Poona, PP. 181-183.
- Vagyani, B.A. & Jamane, M.R. (1989). A new species of Agathioxylon from the Kamthi formation of Chandrapur district, Maharashtra.
Proc.Spl.Ind.Geo.Con., Poona, PP 181-183
- Vagyani, B.A. & Raju, A.V. (1982). A new species of fossil gymnospermous wood. Araucarioxylon Kraus from Nandori, Maharashtra State, Biovigyanam, 7 : 11-13.

- Vagyani, B.A. & Raju, A.V.V. (1986). Australoxylon maheshwari
A petrified coniferous wood from Kamthi beds of
Chandrapur Dist. (M.S.) India. J.Bot., 9(1): 54-56.
- Varadpande, V.G. (1977). Dadoxylon satnauriense, a new species
of petrified gymnospermous wood from the Lower
Gondwana of India. J.Uni.Poona Sci.Tech., 50:159-162.
- Venkatachala, B.S. (1987). Concepts, limits and extension of the
Indian Gondwana - an introduction to theme. The
Palaeobotanist, 36 : 6-13.
- Venkatachala, B.S. & Rawal, M.S. (1973). Occurance of Permian
Palynofossils in the Chingelpul area, Palar Basin. Bull.
Oil.Nat.Gas.Comm., 10 : 105-108.
- Vrendenburg, E.W. (1910). Summary of Geology of India, Calcutta.
- Wadia, D.N. (1953). Geology of India : 1-336, London.
- Wadia, D.N. (1957). Geology of India, 3rd Edn., Revised Mac
Millan, London.
- Walton, J. (1925). On some South African fossil woods.
Ann.S.A.fr.Mus., 22 : 1-26.
- Zeiller, R. (1896). Etude Sur quelques Plantes fossils, en
particul ier Vertibraria Glossopteris des environs de
Johannesburg (Transvaal). Bull.Soc.geol.France, Ser.3,
24 : 349-379.
- Zeiller, R. (1902). Observations sur quelques plantes fossils des
Lower Gondwana. Mem.Geol.Surv.India., Palaeont indica,
N.S., 2(1) : 1-40.