

A C K N O W L E D G E M E N T

I wish to record my sincere thanks to my research guide Dr. B. K. Chougule, Department of Physics, Shivaji University, Kolhapur for his keen interest and able guidance. This work could not have been possible in the absence of his continued encouragement.

I consider it my pleasant duty to offer my thanks to Dr. R. N. Patil, Head of the Department of Physics, Shivaji University, Kolhapur for his support, encouragement and co-operation. The help and co-operation rendered by Dr. S. A. Patil is also gratefully acknowledged.

The XRD and IR work reported in this thesis was carried out at the Materials Research Laboratory, Indian Institute of Science, Bangalore and the SEM work at the University of Mysore. I thank both these Institutes for kindly extending these facilities to me.

The assistance given by Mr.G.D.Mahadik and Mr.M.M.Chougule of the Materials Science Laboratory, during the experimental work is worth appreciating. I thank both of them. I take this opportunity to also thank my research colleagues Mr. R.S. Shinde, Mr. D.S. Revadkar, Mr. D.I. Jadhav, Ms. L.D.Kulkarni and Mr. R.B. Pujar.

My thanks are also due to the authorities of Shivaji University, Kolhapur for making me available the necessary facilities to carry out this work. I thank the Principal, Netaji Subhashchandra Junior College, Yelgaon and the Rayat Shikshan Sanstha, Satara for their encouragement and help.

Finally, I reserve a special word of thanks to Mr. Mohan Kamat of M/s Data Acrobats, Kolhapur, for word processing facility used in typing this thesis and Mr. Ravan for neat and clean tracings.

KOLHAPUR

S. R. PATIL